



Jorge Lepikson Neto

**“Influência da suplementação por flavonóides na
formação da madeira em Eucalipto”**

*“Influence of Flavonoid supplementation on
Eucalyptus Wood formation”*

Campinas, 2013



Universidade Estadual de Campinas
Instituto de Biologia

Jorge Lepikson Neto

**“Influência da suplementação por flavonóides na
formação da madeira em Eucalipto”**

Orientador: Prof. Dr. Gonçalo Amarante Guimarães Pereira

*“Influence of Flavonoid supplementation on
Eucalyptus Wood formation”*

Tese de doutorado apresentada ao Programa de Pós Graduação em Biologia Funcional e Molecular do Instituto de Biologia da Universidade Estadual de Campinas para obtenção do título de Doutor em Biologia Funcional e Molecular, na área de Bioquímica.

Ph.D. thesis submitted to the Graduate Program in Functional and Molecular Biology of the Institute of Biology from the State University of Campinas to obtain the title of Doctor of Functional and Molecular Biology in Biochemistry.

Este exemplar corresponde à versão final da tese

Defendida pelo aluno Jorge Lepikson Neto e

orientada pelo Prof. Dr. Gonçalo Amarante Guimarães Pereira

Assinatura do Orientador:

Campinas, 2013

Ficha catalográfica
Universidade Estadual de Campinas
Biblioteca do Instituto de Biologia
Mara Janaina de Oliveira - CRB 8/6972

L557i Lepikson-Neto, Jorge, 1980-
Influência da suplementação por flavonóides na formação da madeira em Eucalipto / Jorge Lepikson Neto. – Campinas, SP : [s.n.], 2013.

Orientador: Gonçalo Amarante Guimarães Pereira.
Tese (doutorado) – Universidade Estadual de Campinas, Instituto de Biologia.

1. Eucalipto. 2. Lignina. 3. Flavonóides. I. Pereira, Gonçalo Amarante Guimarães, 1964-. II. Universidade Estadual de Campinas. Instituto de Biologia. III. Título.

Informações para Biblioteca Digital

Título em outro idioma: Influence of flavonoid supplementation on Eucalyptus wood formation

Palavras-chave em inglês:

Eucalyptus

Lignin

Flavonoids

Área de concentração: Bioquímica

Titulação: Doutor em Biologia Funcional e Molecular

Banca examinadora:

Gonçalo Amarante Guimarães Pereira [Orientador]

Carlos Alberto Labate

Fabio Marcio Squina

Isabel Rodrigues Gerdhardt

Jorge Mauricio Costa Mondego

Data de defesa: 02-07-2013

Programa de Pós-Graduação: Biologia Funcional e Molecular

Campinas, 02 de julho de 2013

BANCA EXAMINADORA

Prof. Dr. Gonçalo Guimarães Amarante Pereira
(Orientador)


Assinatura


Prof. Dr. Carlos Alberto Labate


Assinatura


Dr. Fabio Marcio Squina


Assinatura

Dra. Isabel Rodrigues Gerhardt


Assinatura

Dr. Jorge Mauricio Costa Mondego


Assinatura

Dr. José Antônio Bressiani

Assinatura

Prof. Dr. Cláudio Chrysostomo Werneck

Assinatura

Dra. Odalys Garcia Cabrera

Assinatura

Agradecimentos:

Gostaria de agradecer primeiramente à minha enorme família em Salvador, pai, mãe, irmãos, tios, tias, primos e primas;

Ao Professor Gonçalo pela oportunidade e motivação;

A todos os amigos do Laboratório de Genômica da Unicamp, em especial à Eliane, Ana Deckmann, Zezé, Marcela e Leandro que contribuíram diretamente para esta tese; e aos demais que muitas vezes contribuíram sem perceber apenas pela convivência, me considero um privilegiado por poder trabalhar em um local com tantos cientistas excepcionais;

Ao Professor José Carlos Rodrigues do Instituto de Investigação Científico Tropical de Lisboa, pela orientação no estágio de doutorado sanduíche.

A International Paper do Brasil, parceira desde o mestrado, fornecendo todo o suporte técnico na área florestal; Adriano, Carla, Tião, Beto e Danielle, obrigado!

Aos amigos: Marcão, Osório, Alessandro, Marcela, Zapas, Raissa, Pedro, Egito, Barrão, Bispo, Santoro, Jlia, Camila, Angela, Bolinha, Thanuci e tantos outros...

A todos os membros da banca de defesa e qualificação;

Agradeço ainda a Capes e ao CNPQ pelo apoio financeiro.

Meu sincero obrigado!

Jorge.

Resumo:

O *Eucalyptus* é o gênero florestal mais plantado no Brasil e no mundo. É composto por mais de 700 espécies, com numerosas subespécies, variedades botânicas e híbridos. O Brasil é atualmente o primeiro produtor de celulose de fibra curta do mundo, e décimo produtor de papel em 2010, além de estar na vanguarda do conhecimento de produção de bicompostíveis, sendo seu setor florestal altamente desenvolvido. Uma das principais limitações para o uso da biomassa lignocelulósica de eucalipto para a produção de energia e seu melhor aproveitamento na produção de papel vem da sua estruturação da parede celular, com a lignina impedindo o acesso a celulose. Para aperfeiçoar este processo é necessário estudar as características da parede celular e dos genes que controlam sua formação. Os flavonóides narigenin-chalcona e narigenin sintetizados respectivamente pelos genes chalcone synthase (CHS) e isomerase (CHI), tem comprovada influência na lignificação de diversas espécies vegetais e expressão diferencial observada em diferentes espécies de Eucalipto, sendo desta forma o estudo da sua influência na formação da madeira em Eucalipto um potencial ponto para o desenvolvimento de novos produtos para o setor florestal brasileiro visando o melhoramento do rendimento da espécie. Desta forma a presente tese de doutorado analisou a influência destes flavonóides na formação da madeira em um híbrido comercial de *Eucalipto grandis* x *Eucalipto Urophylla*, caracterizando quimicamente madeira e avaliando as alterações no perfil transcricional de mudas suplementadas, estando assim seus resultados diretamente ligados a ambição do país, hoje maior produtor mundial de celulose, em permanecer como referencia no setor.

Abstract:

The genus *Eucalyptus* is the most planted forest in Brazil and worldwide. It comprises more than 700 species, with numerous subspecies, botanical varieties and hybrids. Brazil is currently the largest producer of hardwood pulp in the world, and tenth paper producer in 2010, in addition to being at the forefront of knowledge of biofuel production, with its highly developed forestry sector. A major limitation to the use of lignocellulosic biomass for the production of eucalyptus energy and its better use in paper production comes from its structure of the cell wall with lignin preventing access to cellulose. To optimize this process it is necessary to study the characteristics of the cell wall and the genes that control their formation. Flavonoids naringenin-chalcone and naringenin respectively synthesized by genes chalcone synthase (CHS) and isomerase (CHI), has proven influence on lignification in various plant species and differential expression observed in different species of *Eucalyptus*, and thus the study of their influence on wood formation in eucalyptus a potential for the development of new products for the Brazilian forest sector aimed at improving the performance of the species. Thus the present doctoral thesis examined the influence of these flavonoids in wood formation in a commercial hybrid *Eucalyptus grandis* x *Eucalyptus Urophylla*, characterizing its wood chemically and evaluating changes in the transcriptional profile of seedlings supplemented with flavonoids. Results are therefore directly linked to the ambition of the country, today world's largest producer of pulp, to remain as a reference in the sector.

Sumário

1- Introdução	1
1.1. O Eucalipto e sua importância econômica	1
1.2. A parede celular	4
1.3 A Lignina e a via de fenilpropanóides	7
1.4 Estresse, flavonóides e lignificação	11
2. Objetivos	14
2.1. Objetivos específicos	14
3. Material e métodos	15
3.1. Material Biológico e delineamento experimental	16
3.2. Análises químicas e histológica	16
3.3. Extração de RNA e sequenciamento	19
3.4 Hidrólise enzimática	20
4. Apresentação dos trabalhos:	22
Capítulo 1: Caracterização química da parede celular de mudas de eucalipto suplementadas com flavonóides	24
Capítulo 2: Análise da expressão gênica e hidrólise enzimática	36
5. Discussão	67
6. Conclusões	70
7. Referências	71
Anexo 1. Déposito patente INPI	80
Anexo 2 Genes diferencialmente expressos	82

□

1- Introdução

1.1. O Eucalipto e sua importância econômica

O gênero *Eucalyptus* é composto por cerca de 500 espécies e 200 híbridos, sendo endêmico da Austrália e Tasmânia, com poucas espécies ocorrendo naturalmente fora destes locais (Ladiges, Udovicic, & Nelson, 2003).

Os eucaliptos crescem rápido, suportam solos pobres em nutrientes e possuem madeira de boa qualidade, motivos pelos qual o *Eucalyptus* é o principal gênero cultivado nas plantações industriais existentes no mundo: cerca de 20 milhões de hectares plantados em mais 90 países (Paiva et al. 2011).

A maioria das espécies de *Eucalyptus* se caracteriza por serem árvores típicas de floresta altas, atingindo alturas que variam entre 30 e 50 metros. Algumas espécies podem atingir excepcionalmente 100 metros de altura, sendo consideradas as árvores florestais latifoliadas mais altas do mundo (Boland et al. 2006).

Outras espécies comuns de *Eucalyptus* são aquelas típicas de florestas abertas, com árvores menores que atingem de 10 a 25 metros (Boland et al. 2006)). Cerca de 30 a 40 espécies são arbustivas (conhecidas como "Malees") e se caracterizam por possuírem diversos troncos oriundo de um único núcleo lenhoso subterrâneo. Apesar da grande variedade em relação ao habitat e aparência, todas as espécies de eucalipto devem possuir características comuns para serem classificadas dentro do gênero como, por exemplo, a presença de glândulas de óleo nas folhas, tipo de flores e frutos, entre outros descritores (Boland et al. 2006). Entretanto, a característica morfológica mais importante do eucalipto seja possivelmente a sua casca, sendo que os diversos tipos distintos de casca são capazes de caracterizar grupos de espécies estreitamente relacionadas (Boland et al. 2006).

Devido à grande quantidade de espécies, o gênero *Eucalyptus* possui uma alta gama de adaptações edafoclimáticas, podendo ser encontradas espécies que se adaptam bem tanto em temperaturas negativas (até -18 °C) quanto em climas quentes (+35°C), e em solos que apresentam uma variação desde arenosos a encharcados (Silva, 2001).

No Brasil, o eucalipto é responsável pelo abastecimento da maior parte do setor industrial de base florestal. Atualmente, existem cerca de 4,5 milhões de hectares de florestas plantadas de *Eucalyptus* no Brasil (Bracelipa, 2011). O segmento de celulose e papel é composto por aproximadamente 220 empresas distribuídas em 450 municípios, de 17 estados, e utiliza madeira exclusivamente de florestas plantadas. Segundo a Sociedade Brasileira de Silvicultura, em 2010 a produção de

papel alcançou 9,8 milhões de toneladas (aumento de 4,4% em relação ao ano anterior) e a de celulose 14 milhões de toneladas (aumento de 6,4% em relação ao ano anterior). Isso tornou o Brasil o maior produtor mundial de celulose de eucalipto (Bracelpa, 2011). No setor de carvão vegetal, a produção atinge anualmente a marca de 18,8 milhões de metros cúbicos (70% da produção nacional). Outro setor importante é o de chapa de fibra que tem uma produção anual de 558 mil m³, dos quais a madeira de eucalipto é responsável por 100% da produção nacional (Bracelpa, 2011). No setor de chapas de fibra aglomerada, a produção perfaz 500 mil m³, representando quase 30% da produção nacional (Bracelpa, 2011).

As receitas de exportação desses produtos florestais atingiram US\$ 7,4 bilhões em 2007 e, apesar de apresentar grande queda devido à crise de 2009, em 2010 o setor deu sinais de recuperação com aumento de 33% em relação ao ano anterior nas exportações, totalizando US\$ 6,7 bilhões. A receita de exportações de celulose registrou 41,2% de aumento em 2010, chegando a US\$ 4,7 bilhões, contribuindo expressivamente na economia do país, uma vez que o setor de base florestal emprega, direta e indiretamente, 6,5 milhões de pessoas em todos os segmentos (Bracelpa 2011).

Outra característica importante da espécie é a alta capacidade de produção de biomassa - até 100 m³/ha/ano de madeira rica em fibra de celulose (Bracelpa, 2011) - e uma considerável habilidade no seqüestro de carbono (10-14 ton/ha/ano), sendo que este saldo se mantém positivo mesmo quando considerada a emissão de CO₂ proveniente da produção de carvão e papel (Bracelpa, 2011).

Nos últimos anos a demanda por fontes de energia alternativa vem aumentando cada vez mais, principalmente devido à necessidade da substituição da plataforma petroquímica, com um desejo crescente por parte da sociedade, dos governos e da comunidade científica em se desenvolver fontes de energia sustentáveis (Pu et al. 2008). Dentre as principais alternativas para a produção de biocombustíveis, a biomassa lignocelulósica assume posição de destaque como uma das fontes de energia sustentável mais promissora, principalmente a madeira e fibras produzidas em áreas de silvicultura e em áreas agrícolas marginais (Mansfield 2009). A partir da biomassa é possível utilizar seus principais componentes (celulose, hemicelulose e lignina) como fonte de carbono para produção de biocombustíveis e, assim, substituir a utilização de combustíveis fósseis, complementando as culturas já existentes e aumentando a diversidade de matéria prima produtora de combustíveis (Zhu and Pan 2010; Yáñez-S et al. 2012).

Espécies arbóreas de rápido crescimento como *Eucalyptus* e *Populus* tem enorme potencial para a aplicação na produção de combustível de segunda geração

(Carroll and Somerville 2009; Hinchey et al. 2009; Mansfield 2009; Richard 2010; Somerville et al. 2010; Mizrachi et al. 2011; Séguin 2011). *Eucalyptus* e seus híbridos estão entre os líderes mundiais de produção de biomassa e suas ótimas propriedades da madeira, alta produtividade e ampla adaptabilidade poderão permitir a produção sustentável e eficiente de etanol de segunda geração.

A principal limitação para o uso da biomassa lignocelulósica está na sua recalcitrância à degradação, ligada à estrutura e composição das paredes celulares lignificadas (Séguin 2011). A sua complexa estrutura e composição impedem o aproveitamento total dos açúcares fermentáveis, sendo o processo de lignificação justamente um dos principais fatores que determinam tal complexidade. Sendo assim, vários esforços vêm sendo concentrados para melhor se compreender este processo e produzir árvores com menos lignina ou que sejam mais susceptíveis à degradação química (Weng et al. 2008; Mansfield 2009).

Atualmente a matéria prima utilizada para produção de biocombustíveis provém das culturas de cana de açúcar, milho e soja (Chisti 2007). Apesar de já estabelecida, esta parece não ser a melhor estratégia energética a longo prazo, uma vez que a atual capacidade de produção agrícola não é suficientemente capaz de suprir sustentavelmente a demanda, além de já existir uma competição por terras agriculturáveis para produção de alimentos (Mansfield et al. 2012).

Além disso, o Brasil possui altos níveis de produtividade de eucalipto comparados à média mundial devido, principalmente, às condições climáticas aliadas aos investimentos em pesquisa e desenvolvimento no setor, especialmente nas técnicas convencionais de melhoramento genético. Em 2010, a produtividade média da madeira proveniente de *Eucalyptus* alcançou a marca de 44 m³/ha/ano, enquanto outros produtores, como Chile, África do Sul e Portugal, apresentaram níveis abaixo de 30 m³/ha/ano (Bracelapa, 2011).

Sendo o Brasil o maior produtor mundial de celulose de eucalipto (Bracelapa, 2011) além de ser um dos maiores produtores mundiais de bioetanol - na safra de 2008/2009 produziu 27,5 bilhões de litros de etanol a partir do caldo da cana-de-açúcar (Unica, 2010) - é oportuno que a pesquisa científica do país direcione seus esforços a juntar e ampliar esses dois grandes mercados para continuar na vanguarda do conhecimento mundial. Para isso, é imprescindível a compreensão dos mecanismos de formação da parede celular e sua influência na composição final da madeira.

1.2. A parede celular

A parede celular do eucalipto é composta aproximadamente por 50% celulose, 25% hemicelulose e 25% lignina. Essa constituição química varia entre espécies do mesmo gênero e, em menor escala, entre os exemplares da mesma espécie (Kirst et al. 2004). Esses compostos são formados a partir da combinação de alguns elementos químicos, considerada como obedecendo à seguinte proporção geral: 50% de carbono, 44% de oxigênio, 6% de hidrogênio, 0,1% de nitrogênio e 0,3% de cinzas (material inorgânico) (Ragland et al. 1991).

A parede celular tem várias funções: regula a expansão celular, contribui para a adesão entre as células, funciona como barreira para potenciais patógenos e determina as propriedades físicas da planta, bem como a resistência mecânica que permite que algumas plantas alcancem grandes alturas (Boudet 2000; Grima-pettenati et al. 2012). A parede celular também atua como exoesqueleto, controlando a forma da árvore, e é necessária para regular as relações hídricas das plantas, pois determina a relação entre turgor e volume celular, sendo o crescimento vegetal limitado principalmente pela capacidade de expansão da parede celular. As células do xilema, tais como as encontradas na madeira, são notáveis por possuir parede secundária altamente espessada e reforçada por lignina (Fukuda 1996; Plomion et al. 2001).

O desenvolvimento do xilema secundário pode ser dividido em cinco eventos sequenciais: divisão celular, alongação, biossíntese da parede celular, lignificação e morte celular programada. Em cada fase, a expressão ordenada de famílias de genes é regulada por sinais ambientais e do desenvolvimento da planta, que influenciam a composição e a morfologia da parede celular das células do xilema, sendo assim o fator mais importante na determinação das propriedades da biomassa lignocelulósica (Chaffey et al. 2002).

As microfibrilas de celulose são estruturas rígidas que contribuem para a resistência e a disposição estrutural da parede celular. Os glucanos individuais que constituem as fibrilas estão firmemente alinhados e ligados entre si, formando uma fita altamente ordenada que exclui água e é relativamente inacessível ao ataque enzimático. Como resultado, a celulose é muito forte, muito estável e resistente à degradação (Fukuda 1996; Plomion et al. 2001; Mansfield 2009; Mansfield et al. 2012).

As diferentes funções da parede celular refletem sua composição entre os diferentes tipos celulares e processos de diferenciação celular. Acredita-se que até 15% dos genes contidos no genoma vegetal possam estar envolvidos com síntese da parede celular e sua regulação (Darley et al. 2001). Um grande número de genes também é necessário para a síntese de polímeros da parede celular, como lignina,

celulose, xilano, xiloglucano, pectinas, proteínas estruturais e outros componentes, bem como o envolvimento de genes relacionados a hormônios, citoesqueleto e sua regulação (Somerville et al. 2010). Figura 1.

Assim, para entender como as plantas constroem seus diferentes tipos de parede celular, é necessário conhecer os genes que participam das vias metabólicas de síntese e aqueles que participam da sua regulação, uma vez que a formação da parede celular secundária é finamente regulada por uma cascata de fatores de transcrição. (Demura and Fukuda 2007; Zhong et al. 2007; Zhong and Ye 2007; Zhong et al. 2008; Zhong et al. 2010; Zhang et al. 2011; Zhao and Dixon 2011).




Figura 1: A) Parede celular contendo microfibrilas de celulose, hemicelulose, lignina, pectina e proteínas solúeis; B) Celulose sintáse em forma de complexos roseta; C) Lignificação ocorre nas camadas S1, S2 e S3 da parede celular; retirado de: Sticklen 2008.

1.3 A Lignina e a via de fenilpropanóides.

A lignina é um termo genérico para um grupo grande de polímeros aromáticos resultantes do acoplamento oxidativo combinatorial de fenilpropanóides específicos (4-*hydroxyphenylpropanoids*) (Boerjan et al. 2003; Ralph et al. 2004).

A lignina é encontrada na parede celular de vários tipos de tecidos de sustentação e vascular, especialmente traqueídes e elementos de vaso, correspondendo entre 15 a 35% do peso seco da madeira. Ela é depositada, sobretudo, no espessamento da parede secundária, mas também pode ocorrer na parede primária e na lamela média em íntimo contato com a celulose e hemicelulose presentes (Vanholme et al. 2010).

A composição química elementar da lignina inclui carbono, hidrogênio e oxigênio, mas a contribuição relativa destes elementos pode variar, principalmente se a lignina for obtida de coníferas ou de folhosas, além de depender do método de isolamento da mesma. A rigidez mecânica da lignina fortalece os caules e o tecido vascular, permitindo o crescimento ascendente e possibilitando que a água e os sais minerais sejam conduzidos através do xilema sob pressão negativa, sem haver colapso do tecido (Weng et al. 2010).

A sua estrutura química consiste de três monômeros que diferem entre si pelo grau de metilação, sendo eles: álcool trans-coniferílico, álcool trans-sinapílico e álcool para-trans-cumárico. Estes três álcoois levam a formação dos diferentes tipos de lignina: guaicílica (G), sinapílica (S) e cumarílica (H), respectivamente. Em angiospermas, a lignina S é predominantemente encontrada em fibras enquanto a G está mais presente em vasos, e a última lignina, H, é encontrada em quantidade muito inferior, sendo mais presente em gramíneas (Besseau et al. 2007; Vanholme et al. 2010; Grima-pettenati et al. 2012;).

A diferença na composição das ligninas e a quantidade de cada um dos tipos de lignina influem nas características da madeira (Kirst et al. 2004). Taxas elevadas de S/G são vantajosas para a produção de celulose, papel e bioenergia, devido a maiores taxas de deslignificação, gerando menor consumo de produtos químicos e rendimentos mais elevados (Rodrigues et al. 1999; Gomes et al. 2008;).

Uma grande quantidade de genes é atribuída como participante da formação da lignina e, apesar de sua maioria já ter sido caracterizada, a via é constantemente revisitada e ainda é motivo de grande debate em algum nível de sua síntese (Boerjan et al. 2003; Grima-pettenati et al. 2012).

A síntese de lignina envolve a via geral de fenilpropanóides, começando pela desaminação da fenilalanina levando a produção de ésteres de *hydroxycinnamoyl-CoA* (Grima-pettenati et al. 2012). As principais enzimas que participam desta cascata de

reações são: *phenylalanine ammonia-lyase* (PAL), *cinnamate 4-hydroxylase* (C4H) e *4-coumarate-CoA ligase* (4CL) e posteriormente, *ferulate 5-hydroxylase* (F5H), *cinnamoyl-Co-A NADP reductase* (CCR), *cinnamyl alcohol dehydrogenase* (CAD), que participam das reações da etapa que antecede à polimerização da lignina (Boudet 2000; Darley et al. 2001; Boudet et al. 2003;).

Apesar do esforço realizado na pesquisa, especialmente nos últimos anos, a regulação fisiológica das taxas de biossíntese e a disposição espacial e temporal das unidades monoméricas que compõem o polímero de lignina ainda continua pouco entendida, principalmente devido a existência de diversos grupos separados de enzimas relacionados com a sua produção, e a grande intersecção de vias metabólicas a estes relacionados. (Ferrer et al. 2008).

Além da deposição natural destes compostos, sua biossíntese pode também ser induzida por várias condições de estresses bióticos e abióticos, tais como infecção, ferimento, patógenos, estresse metabólico e perturbações na parede celular (Caño-Delgado et al. 2003; Tronchet et al. 2010).

As vias de flavonoides e lignina estão intimamente relacionadas, tendo enzimas e intermediários em comum. Fazem parte da via de fenilpropanóides, sintetizados em organismos vegetais a partir do aminoácido fenilalanina (Besseau et al. 2007) (ver Figura 2). O nome "fenilpropanóide" é derivado da junção do grupo fenila (anel aromático) e uma cadeia lateral de três carbonos (grupo propila).

Os flavonóides *narigenin-chalcone* e *naringenin* são intermediários do metabolismo de fenilpropanóides em plantas, sendo estes sintetizados pelas enzimas *chalcone sintase* (CHS) e *chalcone isomerase* (CHI), respectivamente. Existem estudos mostrando que estes flavonoides influenciam o crescimento de diferentes plantas inibindo a síntese de lignina (Chen et al. 2004; Deng et al. 2004; Chen et al. 2005; Chen et al. 2011), e estes efeitos parecem residir na inibição de enzimas chave da via de ligninas, principalmente a *4-coumarate CoA Ligase* sintetizada pelo gene 4CL. O seu produto, *p-coumaroyl – CoA*, está situado na junção das rotas metabólicas levando a produção de flavonóides ou lignina *sensu stricto*, sendo o substrato comum a ambas (Voo et al. 1995; Campbell and Sederoff 1996; Besseau et al. 2007).

Estudos anteriores demonstraram que a quantidade e a composição monomérica da lignina pode ser reduzida e modificada pela inibição do gene 4CL (Lee et al. 1997; Tamagnone et al. 1998). Também foi demonstrado que *narigenin-chalcone* e *naringenin* inibem a atividade de 4CL *in vitro* e promovem a inibição do crescimento de diversas gramíneas, como milho (*Zea Mays*), arroz (*Oryza sativa*) e *Echinochloa oryzicola*, uma erva daninha (Deng et al 1999). Também foi demonstrado que, em

Pinus, a supressão do gene 4Cl afetou o fénótipo das plantas, resultando em plantas anãs (Wagner et al. 2009). Por outro lado, em *Aspen* ocorreu o efeito inverso, resultando em plantas com crescimento acelerado (Hu et al. 1999).

Dessa forma, fica claro que estes compostos contribuem para a formação e o desenvolvimento de diversos vegetais e que o estudo da sua suplementação em *Eucalyptus* é de fundamental importância no entendimento da lignificação do mesmo, permitindo uma análise ampla e integrada da influência destes flavonóides na composição química da madeira e na sua expressão gênica.




Figura 2: Vista conjunta das vias de flavonoides e lignina. Genes representados: ALDH, aldehyde dehydrogenase; C3H, *p*-coumarate 3-hydroxylase; C4H, *C*4-hydroxylase; F5H, ferulate 5-hydroxylase; CAD, cinnamyl alcohol dehydrogenase; CCoAOMT, caffeoyl-CoA O-methyltransferase; CCR, cinnamoyl-CoA reductase; CHI, chalcone isomerase; 4CL, 4-coumaroyl-CoA ligase; COMT I, caffeic acid O-methyltransferase; F3H, flavanone 3-hydroxylase; F39H, flavonoid 39-hydroxylase; FLS, flavonol synthase; HCT, hydroxycinnamoyl-CoA shikimate; PAL, phenylalanine ammonia lyase; SGT, sinapate UDPglucose sinapoyltransferase; SMT, sinapoylglucose malate sinapoyltransferase; UGTs, UDP sugar glycosyltransferases (Retirado de Besseau, 2007).

1.4 Estresse, flavonóides e lignificação

Todos os fenilpropanóides são formados a partir do ácido cinâmico (cinamato) que, por sua vez, é formado a partir do aminoácido fenilalanina pela ação da enzima PAL (*phenylalanine ammonia lyase*) sendo o ponto de encontro entre o metabolismo primário (via do shikimato) e secundário (Dixon and Paiva 1995).

Muitos fenilpropanóides simples (com o esqueleto de carbono C6-C3 derivado da fenilalanina) são produzidos a partir do cinamato através de varias reações de hidroxilação, metilação e ou desidratação, ai se incluindo os precursores da lignina: álcool trans-coniferílico, álcool trans-sinapílico e álcool para-trans-cumárico (Dixon and Paiva 1995; Vanholme et al. 2010).

Estresse e a via de fenilpropanóides estão intimamente ligados com um grande número de fenilpropanóides sendo produzidos em resposta ao estresse (figura 3), sendo derivados do esqueleto C15 dos flavonóides e sintetizados via CHS (*chalcone synthase*) pela condensação de *p-coumaroyl-CoA* e três moléculas de *malonyl-CoA* (Harbone 1988).

A principal função deste fenilpropanóides é de proteção, estresses abióticos, como a seca, salinidade, temperaturas extremas, toxicidade química e estresse oxidativo são ameaças sérias para a agricultura sendo a principal causa de perda de culturas em todo o mundo, reduzindo o rendimento médio para a maioria das principais plantas de cultura por mais de 50% (Wang et al. 2004). Alguns compostos são induzidos em resposta a ferimentos ou alimentação causada por herbívoros, antocianinas e flavonas aumentam em resposta a altos níveis de luz e irradiação ultravioleta induz a produção de flavonóides e de sinapato em *arabidopsis* (Dixon and Paiva 1995). Este ultimo é um intermediário na biossíntese da lignina siringil (S) em angiospermas, e em alguns taxóons serve como um precursor para a metabolitos secundários solúveis (Ruegger et al. 1999).

Também é sugerido que, em *arabidopsis*, o acúmulo de um ou mais intermediários da via de flavonóides promove a repressão de toda a via de fenilpropanoides ou de ramos da via, levando à produção de compostos específicos (Li et al. 1993). Esse mecanismo serviria para balancear a via e existem precedentes que demonstram que produtos da via podem regulá-la como, por exemplo, o ácido trans-cinâmico, produto de PAL, bloqueando a ação de CHS e da própria PAL (Bolwell et al. 1988; Mavandad et al. 1990), além da existência de *motifs* comuns aos promotores dos dois genes (Dixon and Harrison 1990). Esta flexibilidade seria importante para

promover alternativas biossintéticas diversas em resposta a diferentes estímulos ambientais e de desenvolvimento (Dixon and Paiva 1995).




Figura 3: Exemplos de fenilpropanóides induzidos por estresse. Retirado de (Dixon and Paiva 1995)

Os flavonóides são um grupo grande e diversificado de importantes metabólitos, envolvidos com inúmeras funções no desenvolvimento e crescimento das plantas. Muitas dessas funções estão envolvidas com atração de vetores animais, polinização, dispersão de sementes e reabsorção de nutrientes minerais de folhas senescentes. Outras funções incluem promover vantagens competitivas para o crescimento em circunstâncias estressantes, como por exemplo, aumentar a tolerância a fatores abióticos, defesa contra herbívoros e patógenos. Devido a sua importância, esse grupo de metabólitos encontra-se presente em todo o reino vegetal, sendo investida grande energia metabólica para a sua produção (Andersen & Markham 2006).

Chalconas, dihidrochalconas e auronas são três grupos distintos que constituem mais de 900 flavonóides descritos e são mais conhecidos como pigmentos de coloração amarela e laranja em flores de plantas, mas que se distinguem de outros flavonóides devido a sua estrutura de cadeia aberta de três carbonos. Esses flavonóides não se restringem apenas às flores estando presente em diversos tecidos vegetais e são convertidos a sua respectiva flavona pela atuação da enzima *chalcone isomerase* (CHI). Central para estas reações e para o metabolismo dos flavonóides esta a formação da narigenin-chalcona através da condensação de três moléculas de malonyl-CoA e uma de p-coumaroyl-CoA catalisada pela enzima *chalcone synthase* (CHS) (Andersen & Markham 2006).

Flavonóides são considerados reguladores naturais do fluxo de auxina em plantas, atuando junto a complexos de proteínas de membrana, e foi demonstrado em mutantes de *arabidopsis* que a acumulação de flavonóides era dependente das condições de luz e controlada pela expressão de CHS, com sua acumulação resultando em redução do crescimento da planta devido à inibição do transporte de auxina (Besseau et al 2007).

A expressão de genes que codificam enzimas da via de flavonóides também é dependente de luz, no entanto se acumulam em raízes de plantas sem acesso a luz se seus brotos estiverem expostos, podendo assim serem transportados pela planta. Da mesma forma, foi demonstrado que após a aplicação de flavonóides na raiz da planta estes podem ser transportados por longas distâncias para tecidos vasculares através de transporte de célula à célula.

Outro exemplo dessa flexibilidade é a via metabólica da raafinose, que é uma família de oligossacarídeos que atuam como transporte de carboidratos no floema, compostos de armazenamento e agentes metabólicos que combatem o estresse. Estudos recentes em *Populus thichocarpa* demonstram que a super-expressão dos genes dessa via *galactinol synthase* (Gals) e *raafinose synthase* (RaFs) resultam em formação de madeira de tensão, maiores quantidades de celulose e aumento da biomassa total (Unda et al. 2012; Unda 2012).

Em suma, apesar do grande potencial de utilização do material lignocelulósico de origem florestal para a produção de energia, este processo ainda está longe de ser viável do ponto de vista econômico, existindo pouco conhecimento sobre os fatores genéticos envolvidos na formação destes tecidos (Plomion et al. 2001), principalmente os envolvidos na sua regulação transcricional (Legay et al. 2010).

Conforme mencionado anteriormente, a complexa estrutura e composição da parede celular impedem o aproveitamento total dos açúcares fermentáveis. Uma vez que o processo de lignificação constitui um dos principais fatores que determinam tal dificuldade, e a deslignificação da biomassa constitui um processo caro, vários esforços vêm sendo concentrados para melhor se compreender este processo e produzir árvores com menos lignina ou que sejam mais susceptíveis à degradação química (Weng et al. 2008; Mansfield 2009;).

Os principais esforços nesse sentido constituem na construção de plantas transgênicas com super-expressão ou repressão dos genes envolvidos com a formação da parede celular (Baucher et al. 2003; Goicoechea et al. 2005; Foucart et al. 2009). Entretanto, esta estratégia, além de ser demorada e trabalhosa, envolve um componente legal que muitas vezes dificulta a comercialização de produtos transgênicos, além de analisar apenas um conjunto pequeno de genes de cada vez.

Assim, é de interesse o desenvolvimento de estratégias alternativas para a obtenção de plantas otimizadas para os processos industriais como, no caso de *Eucalyptus*, seu uso na produção de celulose e papel e, também, como biomassa para geração de bioquímicos de segunda geração. Entre estas estratégias alternativas, metodologias de manejo sobressaem-se como poderosas ferramentas quando aliadas ao conhecimento da genética da espécie, especialmente aqueles genes que se traduzem efetivamente em fenótipos desejáveis.

Diante do que já se sabe sobre os efeitos dos flavonoides sobre o crescimento e desenvolvimento vegetal e seu potencial na regulação de eventos específicos à parede celular, como lignificação, o presente estudo buscou adotar a suplementação de *Eucalyptus* por flavonoides visando o melhoramento da qualidade da madeira para uso industrial.

2. Objetivos

Os objetivos da presente tese de doutorado compreendem o estudo do efeito da suplementação química dos flavonóides narigenin e narigenin-chalcona na formação da parede celular de Eucalipto.

2.1. Objetivos específicos:

- Caracterizar fenotipicamente a parede celular de mudas de Eucalipto, avaliando os efeitos da suplementação por flavonóides na sua composição química;
- Avaliar o efeito da suplementação por flavonóides no perfil de expressão gênica das mudas de Eucalipto;
- Associar as mudanças físico-químicas na composição da madeira de eucalipto ao perfil de expressão gênica;
- Verificar a susceptibilidade das plantas suplementadas à hidrólise enzimática, através da quantificação dos açúcares fermentescíveis presentes no hidrolisado;

3. Material e métodos

3.1. Material Biológico e delineamento experimental.

Mudas de um clone comercial de um híbrido *E. urophylla* x *E. grandis* (*Eucalyptus urograndis*) com 6 meses de idade foram fornecidos pela International Paper (Mogi-Guaçu, Brasil) e cultivados em uma estufa (figura 4). As plântulas foram divididos em 5 grupos, de acordo com as condições de suplementação (para além da solução padrão nutricional fornecido para todos os grupos), como se segue: grupo controle (CT), grupo experimental 1 (CH), suplementado com 0,1 mmol de naringenina-chalcona durante 5 meses; grupo experimental 2 (NAR), suplementado com 0,1 mmol de naringenina, durante 5 meses, grupo experimental 3 (CHSTOP), suplementado com 0,1 mmol de naringenina-chalcona apenas para o primeiro mês, e grupo experimental 4 (NARSTOP), suplementado com 0,1 mmol de naringenina apenas o primeiro mês (figura 5). Todas as soluções foram administradas por aplicação na raiz a aproximadamente 100 a 150 mL por dia. Os tratamentos foram realizados durante 5 meses. A solução padrão nutricional foi composta como descrito previamente por (Sarruje 1975).

Todas as amostras foram analisadas de 5 meses após o início da experiência, independentemente da sua suplementação.

Para as análises químicas as hastes de madeira foram descascadas e moídas num moinho de Thomas-Wiley modelo ED-5 para passar de uma peneira de 1 mm e rastreada em um aparelho de crivagem vibratório, e de 40 a 60 mesh fracção farinha de madeira foram retidos para análise. Para análise histológica os caules foram conservados em solução de F.A.A (formalin:aceticacid: ethanol 50%, 1:1:18 v/v) e para as análises de expressão gênica, conservados imediatamente a -80° C.

Naringenina (4'-, 5 -, 7-trihidroxiflavanona, 95%) e naringenina-chalcona (1,3 - difenil-2-propen-1-ona, 97%) foram adquiridos da Sigma-Aldrich Co. (Tóquio, Japão) e AcrosOrganics Co. (Tóquio, Japão), respectivamente.

3.2. Análises químicas e histológica.

3.2.1. Histologia.

Três amostras por grupo após a colheita foram fixados em FAA (formalina: aceticacid: 50% de etanol, 1:01:18 v / v) durante pelo menos 24 horas (Johansen 1940). Todos os materiais foram desidratados utilizando o álcool butílico terciário série (Johansen, 1940), embebidos em parafina (paraplast Plus ® - Fischer) e seccionados a 12-14µm de espessura em micrótomo rotativo. Secções desparafinadas foram duplamente coradas com 1% de solução alcoólica safranina-O e 1% de Astral-blue (Gerlach 1969). As secções foram observadas com um microscópio Olympus BX51 sob luz branca, e as imagens foram obtidas com DP-72 da câmara digital e Image Pro Plus 6,3 software. Devido a limitação de tamanho de grupo, apenas os grupos de tratamento prolongado (CT, CH e NAR) foram analisados.



Figura 4. Mudas de *Eucalypto urograndis* na casa de vegetação. Ao fim dos 5 meses de suplementação as plantas alcançavam aproximadamente três metros, tocando no teto.




Figura 5. Representação ilustrativa do delineamento experimental da suplementação por flavonóides das mudas de *Eucalypto urograndis* em casa de vegetação. CT - suplementação apenas por solução nutritiva; CH - suplementação 5 meses narigenin-chalcona; NR - suplementação 5 meses narigenin; CHSTOP - suplementação 1 mês narigenin-chalcona, 4 meses solução nutritiva; NARSTOP - suplementação 1 mês narigenin, 4 meses solução nutritiva.

3.2.2 Medição do teor de extrativos.

Entre duas e oito amostras por grupo (0,3g, a 1,5 g) foram mantidos individualmente em sacos de filtro (Ankon Technology, Nova Iorque, EUA) e extraiu-se sequencialmente durante 16 horas com etanol a 95%, seguido por um outras 16h em água destilada em um aparelho de Soxhlet de 125ml. Depois de cada extracção as amostras foram deixadas para secar *overnight* a temperatura ambiente e em seguida, secos a 60°C *overnight*, seguido por 2 horas a $102 \pm 2^\circ\text{C}$. Os teores de extrativos foram avaliados pela perda de peso após cada passo (Alves et al. 2012). Apenas uma determinação por amostra foi possível devido à baixa quantidade de amostra disponível.

3.2.3. Médição dos teores de Lignina.

Para a análise dos teores de lignina, um *pool* de cinco indivíduos escolhidos aleatoriamente foram preparados por grupo de tratamento. Foram medidas a lignina Klason (lignina insolúvel em ácido) e lignina total (lignina insolúvel em ácido + lignina solúvel), valores da lignina Klason foram determinados de acordo com a normativa TAPPI T 222 om-02 de acordo com as modificações descritas por por Schwanninger e Hinterstoisser (Schwanninger, Manfred / Hinterstoisser 2002), valor de lignina solúvel foi determinado de acordo com a normativa TAPPI UM 157 250.

3.2.4.Pirólise Analítica.

Uma alíquota de 30 mg de cada amostra previamente extratada, foi moída num moinho vibratório (Mixer MM Mill, Retsch) durante 5 min e mantido em dessecador antes da análise por pirólise analítica. A pirólise analítica foi realizada utilizando uma Pyroprobe CDS 1000 com uma sonda de filamento conectado a um GC (Agilent 6890), com um detector de ionização de chama (FID), através de uma interface aquecida (270°C). A pirólise foi realizada a 600°C durante 5 segundos, utilizando-se 75-77 µg de cada amostra. Coluna capilar: DB1701 (60 mx 0,25 mm, 0,25 película um, J & W Scientific), Condições GC: injetor de 270°C, detector de 270°C, temperatura de programa de 45°C, 4 min isotérmica, seguida de aquecimento a taxa de 4°C min⁻¹ para 250 ° C e 6°C min⁻¹ até 270°C mantido por 8min (Alves et al. 2011; Rodrigues et al. 2001; Rodrigues et al. 1999).

Os picos produzidos pela pirólise foram analisados com Chemstation Software (Agilent Technologies, Palo Alto, EUA) as relações S/G e H/G, foram calculadas a partir da soma das áreas dos picos dos produtos de pirólise atribuídos a fenóis de tipo siringil (S), guaiacil (G) e p-hidroxifenil (H). A proporção cP/cH foi calculado a partir da soma das áreas dos picos dos produtos de pirólise características de pentoses (cP) e hexoses (CH) e representa uma estimativa grosseira da proporção relativa de hemicelulose e celulose. Py-lignina foi calculada como a razão entre a soma das áreas dos picos a partir de produtos de lignina dividida pela soma da área de todos os picos utilizados (lignina e polissacarídeos, 75% da área total), multiplicado por 100% (Alves et al. 2006).

3.3. Extração de RNA e sequenciamento.

3.3.1. Extração de RNA

O RNA foi extraído de acordo com o protocolo descrito por (Zeng and Yang 2002) com as modificações propostas por (Le Provost et al. 2003), concentrações e qualidade foram medidas com Nanodrop 2000 (Thermo Scientific).

3.3.2. Sequenciamento de mRNA

O sequenciamento de RNA mensageiro foi feito na facility de seqüenciamento de nova geração do *Carolina Center for Genome Sciences* da Universidade da Carolina do Norte. Para cada biblioteca 10ug de RNA total foi usado para o seqüenciamento de acordo com o protocolo da Illumina. Controle e qualidade das amostras foram verificadas com o *Bioanalyzer Chip DNA 1000 series II* (Agilent). Para cada biblioteca foram seqüenciados fragmentos *single-end* de 36 ou 50 pares de base com *Illumina Genome Analyzer IIx*. Um total de 8 bibliotecas foram geradas, três réplicas biológicas do grupo controle (CT); duas réplicas biológicas do grupo suplementado cinco meses com narigenin (NAR); e uma biblioteca de cada grupo restante: cinco meses suplementação narigenin-chalcona (CH); um mês suplementação de narigenin (NARSTOP) e um mês suplementação narigenin-chalcona (CHSTOP). Cada biblioteca corresponde ao agrupamento de três amostras individuais.

3.3.3. Alinhamento

Os *reads* Illumina foram filtrados para excluir material ribossomal (utilizando a base de dados SILVA (Pruesse et al. 2007) e de baixa qualidade (Phred > = 20). Os restantes foram alinhados contra as variantes mais elevadas de transcritos do genoma de *Eucalyptus grandis* Phytozome 7,0 (44.974 seqüências) utilizando o alinhador SOAP2 (Li et al. 2009). Para preparar os dados para análise Genebrowser, leituras foram alinhadas contra o genoma de *E. grandis* usando TopHat (Trapnell et al. 2009) para permitir a alinhamentos *spliced*. Ambos os programas foram configurados para permitir até dois *mismatches* (SNPs podem gerar desequilíbrios no alinhamento, especialmente nestes casos, porque as seqüências são de espécies diferentes), descartar seqüências com ambigüidades (Ns) e retornar apenas alinhamentos únicos.

3.3.4 Anotação

O programa Autofact (Koski et al. 2005) foi usado para realizar a anotação automática de todos os contigs EUCANEXT. A principal característica do Autofact é a sua capacidade para retomar a anotação com base em pesquisas de similaridade de seqüência em vários bancos de dados. BLASTx (Altschul et al. 1997) (valor de corte de 1e-5) foi utilizado para alinhar os contigs contra as seguintes bases de dados públicas: a base de dados não redundante (NR), do NCBI; Uniref90 e Uniref100 a base de dados de vias metabólicas KEGG (Kanehisa 2000), e TAIR (versão 10 do

banco de dados de proteínas Arabidopsis. Anotação funcional (GO) foi realizada utilizando BLAST2GO (Conesa et al. 2005), e Mapman software (Thimm et al. 2004) com os parâmetros padrão.

3.3.5 Determinação dos valores de expressão gênica

Os valores de expressão gênica foram medidos utilizando o *FPKM* (*fragments per kilobase of exon per million fragments mapped*), utilizando apenas alinhamentos únicos como descrito por (Mortazavi et al. 2008). Todos os transcritos e seus valores de expressão podem ser acessados no banco de dados EUCANEXT – www.lge.ibi.unicamp.br/eucalyptusdb

3.4 Hidrólise enzimática

3.4.1 Preparação do substrato

O material vegetal de xilema de Eucalipto foi congelado em nitrogênio líquido e liofilizado usando o FreeZone6 (Labconco) a - 51°C e 25 Pa durante 48 horas. Após a liofilização o material foi moído em um moinho de bola (MARCA) em um ciclo de 5 minutos.

3.4.2. Produção do Secretoma

A linhagem de *Neurospora crassa* St.L 74A (Missouri University, Kansas City, <http://www.fgsc.net/>) foi usada para a produção do secretoma. A preparação dos conídios foi realizada inoculando o fungo em 100ml de meio mínimo com sais de Vogel suplementados com 143uL de biotina (5mg, etanol 50% (v/v), 143uL de solução (5g de ácido cítrico, 5g ZnSO₄. 7H₂O, 1g FE(NH₄) .6H₂O, 0,25g de CuSO₄.5H₂O, 0,05g MnSO₄.H₂O, 0,05g H₃BO₃, 0,05g Na₂MoO₄.2H₂O, para 1000 mL), 1,5% de Agar, 2% de sacarose e crescido durante três dias a 30°C no escuro e depois por sete dias com presença de luz a 25°C. A suspensão de conídios foi inoculada em 100mL do mesmo meio de crescimento contendo 2% do material de eucalipto moído na contendo *E. grandis*, *E. urograndis* and *E. urophylla* na proporção de 3:3:1 como única fonte de carbono (Phillips et al. 2011).

3.4.3 Hidrólise

A hidrólise foi realizada de acordo com (Bragatto et al. 2012) com as seguintes modificações: utilizou-se tubos eppendorf de 2ml agitados a 1000 rpm a 30°C no *Thermomixer* (Eppendorf) durante 48 horas. Cerca de 10 mg do substrato foi diluído em 400 ul de tampão 50 mM acetato de sódio pH 5.5 e adicionado 100 uL do secretoma de *N. crassa* na concentração de 0,4 ug/uL determinada usando o *Bradford Kit assay* (BioRad) com BSA padrão. Todas as reações foram realizadas em triplicata. Para a determinação da concentração de açúcar redutor e produção de glicose, os

tubos de 2 mL foram centrifugados a 20.000g durante 10 minutos a 4°C, o sobrenadante foi recuperado e utilizou-se 100 uL para cada reação. Para a determinação do açúcar redutor foi adicionado ao sobrenadante 100 uL de *dinitrosalicylic acid assay* - DNS (Miller 1959), a reação foi aquecida a 99°C por 10 minutos e transferido 100 uL para uma placa ELISA e medida a absorbância a 540 nm (Tecan Infinite M200 instrument). Para o cálculo do conteúdo de glicose, 20 uL do sobrenadante foi adicionado a 100 uL de solução de trabalho do Kit *Glucose Oxidase* (Laborlab) em placas de ELISA e incubados a 37°C por 10 minutos e medida a absorbância a 505 nm.

4. Apresentação dos trabalhos:

Este trabalho objetivou a avaliação da influência da suplementação pelos flavonóides narigenin-chalcona e narigenin, na formação da madeira do híbrido comercial *E. grandis* x *E. urophylla* extensamente plantado no Brasil. O trabalho foi organizado em dois capítulos respectivamente: :

Capítulo 1. Caracterização química da parede celular de mudas de eucalipto suplementadas com flavonóides. Para avaliar a influência da suplementação por flavonóides na composição química da parede celular, as mudas de eucalipto foram analisadas através métodos químicos tradicionais e da pirólise analítica, objetivando quantificar extrativos, pentoses, lignina e composição monomérica da mesma. Este trabalho foi realizado no estágio de doutorado sanduíche do autor, realizado no Instituto de Investigação Científico tropical da Universidade Técnica de Lisboa, sobre a supervisão do Dr. José Carlos Rodrigues.

Manuscrito: “**Flavonoid supplementation reduces the extractive content and increases the syringyl/guaiacyl ratio in *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid trees**” Lepikson-Neto, Jorge ; Alves, A. ; Simoes, R. ; Deckmann, A. C. ; Camargo, E.L.O. ; Salazar, Marcela Mendes ; Rio,MCS ; Nascimento, LC ; Pereira, G. A. G. ; Rodrigues, J. C. Bioresources (Raleigh, N.C), v. 8, p. 1747-1757, 2013.

Data de submissão: 10/2012

Data de aceite: 03/2013

Capítulo 2. Análise da expressão gênica e hidrólise enzimática: Para verificar o impacto da suplementação por flavonóides na expressão gênica e obtenção de açúcares das mudas previamente caracterizadas quimicamente, foi seqüenciado transcriptoma e realizada a hidrólise enzimática do material, no intuito de verificar os efeitos moleculares do processo e associar a sua expressão gênica e potencial obtenção de açúcares fermentáveis as alterações da parede celular verificadas no primeiro capítulo do trabalho, de modo que estas informações completamente inéditas na literatura possam contribuir para futuros trabalhos de melhoramento genético da espécie.

Manuscrito: “**Flavonoid supplementation affects the gene expression of cell wall formation and lignification metabolisms and increases sugar content and saccharification levels in the fast growing Eucalyptus hybrid *E. urophylla* x *E. grandis***” Lepikson-Neto, J.¹, Nascimento L.C.¹, Salazar M.M.¹, Camargo E.L.O.¹, Cairo, J.P.F.²; Teixeira P.J.¹, Marques, W.L.¹; Squina F.M.²; Mieczkowski, P.³; Deckmann, A.C.¹ and Pereira, G.A.G.¹

Data prevista para submissão; 06/2013

Devido aos resultados positivos obtidos, a patente da composição química utilizada para a suplementação dos eucaliptos foi depositada junto ao **INPI – BR1020120269813**, pelo autor do trabalho e seu orientador, juntamente com a parceria da International Paper do Brasil. A folha de rosto da mesma encontra-se no **anexo 1**.

Capitulo 1: Caracterização química da parede celular de mudas de eucalipto suplementadas com flavonóides.

Flavonoid Supplementation Reduces the Extractive Content and Increases the Syringyl/Guaiacyl Ratio in *Eucalyptus grandis* x *Eucalyptus urophylla* Hybrid Trees

Jorge Lepikson-Neto,^a Ana Alves,^b Rita Simões,^b Ana Carolina Deckmann,^a
Eduardo Leal Oliveira Camargo,^a Marcela Mendes Salazar,^a
Maria Carolina Scatollin Rio,^a Leandro Costa do Nascimento,^a
Gonçalo Amarante Guimarães Pereira,^{a,*} and José Carlos Rodrigues^b

The *Eucalyptus* genus plays an important role in the worldwide forest industry, with highly productive plantations supplying high-quality raw material for pulp and paper, wood, and biomass that would otherwise come from native forests. Lignin and extractives are important components for wood structure and protection but they are disruptive elements with respect to some industrial processes involving paper, pulp, and biomass production. This work evaluated effects of supplementation of flavonoids on the wood composition of *Eucalyptus grandis* x *Eucalyptus urophylla* (*E. urograndis*), a commercial hybrid. The wood samples were analyzed for extractives and lignin contents by wet chemical analysis, and the composition of lignin monomers and the carbohydrate hexosan/pentosan ratio were determined by analytical pyrolysis. The results showed that supplementation with the flavonoids naringenin and naringenin-chalcone led to an overall reduction of the extractive content and altered the monomeric composition of lignins towards a higher syringyl content. Thus, the treatment of *Eucalyptus* with flavonoids results in the improvement of wood quality for technological purposes.

Keywords: *Eucalyptus*; Extractives; Lignin; Flavonoids

Contact information a: Laboratório de Genômica e Expressão, Departamento de Genética Evolução e Bioagentes, Instituto de Biologia, Universidade Estadual de Campinas, CEP: 13083-970, Campinas, São Paulo, Brazil; b: Tropical Research Institute of Portugal (IICT), Forestry and Forest Products Group, Tapada da Ajuda, Lisboa, Portugal; *Corresponding author: goncalo@unicamp.br

INTRODUCTION

Wood is a natural resource that forms the basis of a global industry producing fiber, timber, and energy, and it is the fifth most important world trade product (Foucart *et al.* 2009). Wood represents one of the most important sources of energy and biomass on Earth and constitutes an environmentally friendly and renewable alternative to fossil resources. Moreover, wood is an important natural sink for carbon dioxide, one of the major causes of global warming due to the greenhouse effect (Demura and Fukuda 2007).

The genus *Eucalyptus* is the most widely planted hardwood crop in the tropical and subtropical world due to its superior growth, broad adaptability, and multipurpose wood properties (Hu *et al.* 1999). Plantation-grown *Eucalyptus* supplies high-quality woody biomass for several industrial applications while reducing the pressure on tropical forests and their associated biodiversity (Hu *et al.* 1999).

The chemical composition of wood plays an important role in wood applications, especially in chemical conversion. Wood extractives, the non-cell wall components that can be removed by solvents, are necessary for protecting the living tree and derived wood products against disease; however, extractives can be detrimental to pulp and paper, paint, and varnish films and adhesives (Alves *et al.* 2012). Pulpwood from tropically grown eucalyptus trees, including those grown in Brazil, contains a higher content of extractives than those grown in Europe, accounting for a decrease of up to 4% in the pulping yield (Gomide *et al.* 2005). The extractive content is lower at a young harvesting age, and the content increases rapidly with aging (Gomide *et al.* 2005). Lignin composition is also a very important parameter. Eucalyptus lignins are composed of syringyl (S) and guaiacyl (G) units in varying proportions. High syringyl/guaiacyl (S/G) ratios are advantageous for pulp production due to higher delignification rates, reduced chemical consumption, and higher pulp yields (Rodrigues *et al.* 1999). Because of high tree-to-tree variation (Rodrigues *et al.* 1999) and low variation within a tree, lignin composition is a wood trait under strong genetic control (Stackpole *et al.* 2010) even when tension wood is present (Rodrigues *et al.* 2001). Therefore, extensive research efforts are focused on comprehending the lignification process in an effort to design trees through genetic engineering that either have reduced lignin content, produce lignins that are more susceptible to chemical degradation (Weng *et al.* 2008; Mansfield *et al.* 2009), or have altered lignin content (Valerio *et al.* 2003).

Lignin synthesis is a relatively well-established process that starts with the assemblage of radicals produced during the single-electron oxidation of monolignols (Baucher *et al.* 2003). The units resulting from the monolignols, when incorporated into the lignin polymer, are called guaiacyl (G), syringyl (S), and p-hydroxyphenyl (H) (Vanholme *et al.* 2010).

The monolignols are synthesized from the amino acid phenylalanine through the phenylpropanoid and monolignol-specific pathways. Phenylalanine is derived from the shikimate biosynthetic pathway in the plastid (Boudet *et al.* 2003; Rippert *et al.* 2009), which is responsible for the synthesis of a wide variety of secondary metabolic compounds, including lignin and phenolic extractives (Dixon *et al.* 1996; Vogt *et al.* 2010).

The catalytic step performed by the enzyme 4-coumaroyl:CoA-ligase (4CL) likely represents the most important branch point within the central phenylpropanoid biosynthesis pathway in plants (Campbell and Sederoff 1996; Voo *et al.* 1995). Through 4CL activity, cells can produce the precursors for either flavonoid biosynthesis or guaiacyl (G) and syringyl (S) lignin units (Vogt, 2010). The product of 4CL, p-coumaroyl-CoA, is the substrate of the enzyme chalcone synthase (CHS) (Besseau *et al.* 2007), the committing step in flavonoid biosynthesis. This pathway is reviewed in detail elsewhere (Besseau *et al.* 2007; Vanholme *et al.* 2010).

The flavonoids naringenin-chalcone and naringenin, synthesized by the enzymes chalcone synthase (CHS) and chalcone isomerase (CHI), respectively, constitute the primary C15 intermediates in flavonoid biosynthesis (Moustafa 1967; Chen *et al.* 2011). Naringenin was reported to inhibit the activity of 4CL (Voo *et al.* 1995), and this inhibition is positively associated with naringenin sensitivity in several plant species (Denget *et al.* 2004; Yun *et al.* 2009).

The objective of the present work was to evaluate the effects of naringenin-chalcone and naringenin supplementation on the wood composition of *E. urophylla* x *E. grandis*, a commercial hybrid referred to hereafter as *E. urograndis*.

EXPERIMENTAL

Materials

Plantlets of a 6-month-old commercial clone of *Eucalyptus urograndis* were provided by International Paper (Mogi-Guaçu, Brazil) and grown in a greenhouse. The plantlets were divided into 5 groups, according to supplementation conditions (apart from the standard nutritional solution supplied to all groups) as follows: control group (CT); experimental group 1 (CH), supplemented with 0.1 mmol of naringenin-chalcone for 5 months; experimental group 2 (NAR), supplemented with 0.1 mmol of naringenin for 5 months; experimental group 3 (CHSTOP), supplemented with 0.1 mmol of naringenin-chalcone for only the first month; and experimental group 4 (NARSTOP), supplemented with 0.1 mmol naringenin for only the first month. All solutions were administered by root application at approximately 100 to 150 mL daily. The treatments lasted 5 months. The composition of the standard nutritional solution has been described previously (Sarruje 1975). At the end of the experiment, all 5 groups of plantlets were cut, and the stems were kept for analysis; no growth differences were observed between the control and the treatment groups. A total of 30 plantlets were individually analyzed; 9 were only used for histology and 21 for chemical analyses. The same extracted samples were used for pyrolysis. All samples were analyzed 5 months after the start of the experiment regardless of their supplementation. The main wood stems were debarked and ground in a Thomas–Wiley mill model ED-5 to pass a 1 mm sieve and screened in a vibratory sieving apparatus, and the 40 to 60 mesh wood meal fraction was retained for analysis.

Naringenin (4',5,7-trihydroxyflavanone, 95%) and naringenin-chalcone (1,3-diphenyl-2-propen-1-one, 97%) were purchased from Sigma-Aldrich Co. (Tokyo, Japan) and AcrosOrganics Co. (Tokyo, Japan), respectively.

Histology

After harvesting, three stem samples per group were fixed in FAA (formalin: acetic acid: 50% ethanol, 1:1:18 v/v) for at least 24 h (Johansen 1940). All materials were dehydrated using the tertiary butyl alcohol series (Johansen 1940), embedded in paraffin (Paraplast Plus® - Fischer), and sectioned into 12 to 14 µm thick sections with a rotary microtome (Model and brand). Deparaffinized sections were double stained with a 1% alcoholic solution of safranin-O and 1% aqueous astral blue (Gerlach 1969). The sections were observed using an Olympus BX51 microscope under white light, and the images were obtained using a DP-72 digital camera and Image Pro Plus 6.3 software. Due to group size limitation, only the prolonged treatment groups (CT, CH, and NAR) were analyzed.

Extractives Content

Between two and eight samples per group (0.3 to 1.5 g) were kept individually in Ankon filter bags (Ankon Technology, New York, USA) and sequentially extracted for 16 h with 95% ethanol followed by another 16 h in distilled water in a 125 mL Soxhlet apparatus. Afterwards, the extracted samples were allowed to cool and dry under room conditions overnight and then dried at 60°C overnight, followed by 2 hours at 102 ± 2°C. The extractive content was assessed according to weight loss after each step (Alves *et al.* 2012). Only one determination per sample was possible due to the low amount of sample available.

Lignin Content

For lignin analysis, a pool of five randomly selected individuals was prepared per group treatment, and Klason (acid-insoluble lignin) and total lignin content (acid-insoluble lignin plus acid-soluble lignin) were assessed using the averaged values based on the oven-dried, extractive-free weight of each pooled sample as determined using wet chemistry methods. Klason (acid-insoluble) lignin content was determined according to TAPPI T 222 om-02 following the modifications by Schwanninger and Hinterstoisser (2002), and the acid-soluble lignin content was determined according to TAPPI UM 250.

Analytical Pyrolysis

A 30 mg aliquot of each extracted sample was further milled in a vibratory ball mill (Mixer Mill MM, Retsch) for 5 min and kept in desiccators prior to analysis by analytical pyrolysis. Analytical pyrolysis was performed using a CDS Pyroprobe 1000 with a coil filament probe connected to a gas chromatography (GC) unit (Agilent 6890) with a flame ionization detector (FID) using a heated interface (270°C). The pyrolysis was carried out at 600°C for 5 s using 75 to 77 µg of the extractive-free milled samples. Capillary column: DB1701 (60 m x 0.25 mm, 0.25 µm film, J&W Scientific). GC conditions: injector, 270°C; detector, 270°C; temperature program, 45°C, 4 min isothermal, then heating rate 4°Cmin⁻¹ to 250°C and 6°C min⁻¹ to 270°C, hold for 8 min (Rodrigues *et al.* 1999, 2001; Alves *et al.* 2011).

From the pyrolysis product peaks, the following ratios were computed using Chemstation Software (Agilent Technologies, Palo Alto, USA). S/G and H/G ratios were calculated from the sum of the peak areas of the pyrolysis products assigned to syringyl (S), guaiacyl (G), and p-hydroxyphenyl (H) type phenols. The cP/cH ratio was calculated from the sum of the peak areas of the characteristic pyrolysis products of pentosans (cP) and hexosans (cH) and represents a rough estimation of the relative proportion of hemicelluloses and cellulose. Py-lignin was calculated as the ratio of the sum of the areas of the peaks from the lignin products divided by the sum of the area of all peaks used (lignin and polysaccharides, *ca.* 75% of the total area) multiplied by 100% (Alves *et al.* 2006a).

Statistical analysis

To verify significant differences between the controls and the flavonoid-supplemented groups, a one way ANOVA test was performed between the control and each supplemented group. The results were considered significant if $p < 0.05$ and were classified as follows: *, $p < 0.05$; **, $p < 0.01$; and ***, $p < 0.001$. The correlation between ethanol and water extractives was also assessed. All statistics were performed using Analysis ToolPak for Excel 2007.

RESULTS AND DISCUSSION

Histochemical Analysis

The photomicrographs of the stained (astral blue-safranin) transverse section cuttings show that the treatment groups had a more pronounced blue coloration than the control group, especially evident in the NAR group (Fig. 1). These results suggest a relative decrease in the lignin content with a consequent increase in the polysaccharide fraction of the flavonoid-treated wood plantlets because astral blue-safranin dye confers a distinct coloration for carbohydrates and colors cellulose in the absence of lignin blue and

lignin-rich regions red. As shown later using wet chemical analysis, no significant changes in lignin content were found, whereas a clear reduction in the content of ethanol extractives seemed to indicate that the difference in the coloration could result from different amounts of ethanol extractives. Another possibility is the mobilization of the stained compounds during the preparation stages of the material. In fact, mobilization alone could account for the differences between the stain results and the analytical results.




Fig. 1. Photomicrographs of the stained (astral blue-safranin) transverse section cuttings from the control (A-C) and the naringenin (D-F) and naringenin-chalcone (G-I) treatment groups

Extractives and Lignin Content

A summary of results for the extractive and lignin content of the samples is given in Table 1.

Across groups and plants, the ethanol extractives content varied between 4.4% and 13.7% of the dry weight of the wood meal, representing between 51% and 83% of the total extractives content. After ethanol extraction, the water extractive content varied between 2.0% and 6.6% and represented between 26% and 41% of the total extractives content. The ethanol and water extractives contents were positively correlated ($r = 0.56$), and the correlation was highly significant ($p < 0.004$).

Table 1. Summary of the Results of the Extractive and Lignin (Klason and total) Contents

	Extractives (%)			Lignin (%)	
	Ethanol	Water	Total	Klason	Total
CT	10.3(0.34)	3.8(1.09)	14.2(1.25)	22.6(0.03)	26.7(0.01)
CH	6.1(1.01) ***	2.9(0.47)	9.0(1.82) ***	21.5(0.03)	25.5(0.01)
CHSTOP	7.1(0.23) ***	3.4(0.44)	10.5(1.31) **	22.6(0.14)	26.6(0.03)
NAR	7.6(1.57) ***	4.2(0.14)	11.8(0.90) *	22.3(0.01)	26.3(0.17)
NARSTOP	6.9(0.99) **	2.5(0.28)	9.4(0.19) **	22.5(0.02)	26.2(0.40)

Extractive data refer to the mean values of the individual samples from each group. Lignin data refer to the replicate analysis of a 5-sample pool per treatment. CT – control group; CH – 5 months naringenin-chalcone; CHSTOP – 1 month naringenin-chalcone; NAR – 5 months naringenin; NARSTOP- 1 month naringenin. Values in brackets refer to variance.

A decrease in the ethanol extractives content was observed for the flavonoid-treated plants. The average ethanol extractives content decreased from 10.3% (control group) to between 6.1% (CH) and 7.6% (NAR), representing a decrease of between 41% (CH) and 26% (NAR) compared with the control group. The water extractive content was also reduced with flavonoid treatment from 14% (CH) to 34% (NARSTOP), with the exception of the NAR-treated group, which showed an 11% increase compared with the control group. The total extractive content followed the same pattern as the ethanol extractives.

The extractive content of these plants was higher than the extractive content of *Eucalyptus urograndis* at the age of commercial pulpwood. The ethanol extractive content alone was above or close to the total extractive content of the heartwood (7.6%) and at least twice that of the total extractive content of the sapwood (3.7%) of 5.6-year-old trees from an *E. urograndis* clone in a commercial pulpwood plantation in Brazil (Gominho *et al.* 2001). These differences could be explained, at least in part, by the fact that living cells (developing xylem) filled with easily extractable metabolites (Paiva *et al.* 2008) represent a larger percentage of the stem at this age than at older ages.

The reduced extractive content due to flavonoid treatment will increase the productivity of pulping because a higher extractives content accounts for up to 4% of the losses to the kraft pulping yield of clonal eucalyptus in Brazil (Gomide *et al.* 2005).

The Klason and total lignin contents of these plantlets were lower than the reported values for *E. urograndis* at commercial age in Brazil (6 to 7 years old), ranging from 24.2% to 27.1% (Klason) and 27.5% to 30.6% (total) lignin content (Gomide *et al.* 2005). However, these results were obtained on a limited number of samples (7 clones, each a composite sample of three individuals).

The lignin content (Klason and total) was not affected by the flavonoid treatment (Table 1), which was unexpected in light of the histological results (Fig. 1). This result seems to indicate that, in addition to changing the extractive content, the flavonoid treatment could also have changed the extractive composition, accounting for the differences in coloration between the treated and control samples.

Alternatively, the flavonoid treatment could have altered the lignin composition results by making cellulose and other carbohydrates more accessible to the dye astral blue (Fig. 1).

Analytical Pyrolysis

The summary of the analytical pyrolysis results is shown in Table 2. According to these results, flavonoid-treated plants showed a statistically significant increase in the S/G ratio (between 8 and 10% compared with the control) and a decrease in the H/G ratio (between 11 and 16%). Both changes have a favorable impact on pulping because of their effects on delignification rates, chemical consumption, and pulp yields (Rodrigues *et al.* 1999).

Table 2. Summary of the Analytical Pyrolysis Results

		cP/cH	H/G	S/G	Py-lignin%
CT	5	26.8(1.6)	0.092(0.008)	1.41(0.05)	23.2(0.6)
CH	8	28.8(3.3)	0.078(0.007)**	1.55(0.03)**	23.7(0.8)
CHSTOP	3	29.5(2.5)	0.078(0.006)*	1.55(0.03)**	24.2(0.3)
NAR	3	28.2(1.2)	0.077(0.004)*	1.52(0.02)*	24.1(0.7)
NARSTOP	2	29.6(4.2)	0.082(0.009)	1.52(0.01)*	23.0(0.1)

The mean values and standard deviation (brackets) results of the analytical pyrolysis. The first column refers to the number of individual samples analyzed.

The syringyl/guaiacyl (S/G) ratio ranged from 1.41 (CT) to 1.52 (CH and NAR), representing an approximate 7.8% increase in syringyl lignins in the flavonoid treatment. The same result was observed in the short-term flavonoid supplementation groups: the S/G ratio increased to 1.52 and 1.53 in the CHSTOP and NARSTOP groups, respectively, which represents an approximately 7.8% (CHSTOP) and 8.5% (NARSTOP) increase. These values are lower than the values reported for this species (S/G ratio above 2) using analytical pyrolysis and nitrobenzene oxidation (Barbosa *et al.* 2008; Lima *et al.* 2008). However, the analytical pyrolysis results cannot be directly compared because differences in the pyrolysis instruments and columns will inevitably lead to different results.

The H/G ratio decreased from 0.092 (CT) to 0.078 for both CH and NAR groups and to 0.078 and 0.081 in the CHSTOP and NARSTOP groups, respectively, representing an approximate 15% decrease in the H/G ratio in the prolonged treatments (CH and NAR) and 15% (CHSTOP) and 12% (NARSTOP) in the short-term treatments. Interestingly, these H/G values are higher than the average H/G ratios for *Pinus pinaster* (0.64) (Alves *et al.* 2006b), *Pinus caribaea* (0.50) (Godoy *et al.* 2007), and *Picea abies* (0.50), all determined by analytical pyrolysis using the same methodology (Alves *et al.* 2009).

Because no changes in the total lignin content were observed and the S/G ratio increased while the H/G ratio decreased, a direct substitution of p-hydroxyphenyl and guaiacyl units by syringyl units in the composition of lignin is likely. This substitution is one of the key desired aspects of *Eucalyptus* plants (Baucher *et al.* 2003; Grattapaglia and Kirst, 2008; Jung *et al.* 2011), increasing lignin solubility and cellulose accessibility by promoting better delignification (Baucher *et al.* 2003; Huntley *et al.* 2003; Stewart *et al.* 2006). To explain this result, a change in the phenylpropanoid pathway is necessary, with a shift towards S lignin synthesis; expression analyses of supplemented eucalyptus might help elucidate this change.

Both the cP/cH ratio and the Py-lignin content showed no statistically significant differences between the control and treatment groups, although the cP/cH ratio was, on average, higher for all treated groups compared with the control.

In conclusion, it was demonstrated that flavonoid supplementation of *Eucalyptus urograndis* plantlets has a strong influence on its wood composition; the present results are preliminary but have great potential for the improvement of this species towards increased productivity. The addition of flavonoids to a common nutrient medium is inexpensive and easy to accomplish in a eucalyptus nursery. The fact that short-term supplementation (only 1 month) was enough to produce changes in wood composition 4 months later was encouraging, and longer experiments should be performed to determine the impact of supplementation on full-grown Eucalyptus.

CONCLUSIONS

1. Root-applied flavonoid supplementation promotes a significant reduction in extractive content and increases the lignin monomeric composition (S/G ratio) of young *E. urograndis* trees.
2. These preliminary results indicate that flavonoid supplementation can potentially be used as a nutritional complement being a new, viable, and interesting method for improving *Eucalyptus* wood quality for its utilization by the paper and pulp industry and for biomass exploitation.

ACKNOWLEDGMENTS

This project was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível superior (CAPES) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ). The authors would especially like to thank International Paper (Brazil) for its assistance and for providing plant material. Chemical analysis was supported by Fundação para a Ciência e Tecnologia (Portugal), research projects P-KBBE/AGR-GPL/0001/2010 and PTDC/AGR-GPL/098179/2008.

REFERENCES CITED

- Alves, A., Gierlinger, N., Schwanninger, M., and Rodrigues, J. (2009). "Analytical pyrolysis as a direct method to determine the lignin content in wood. Part 3. Evaluation of species-specific and tissue-specific differences in softwood lignin composition using principal component analysis," *J. Anal. Appl. Pyrol.* 85(1-2), 30-37.
- Alves, A., Schwanninger, M., Pereira, H., and Rodrigues, J. (2006a). "Analytical pyrolysis as a direct method to determine the lignin content in wood. Part 1: Comparison of pyrolysis lignin with Klason lignin," *J. Anal. Appl. Pyrol.* 76(1-2), 209-213.

- Alves, A., Schwanninger, M., Pereira, H., and Rodrigues, J. (2006b). "Calibration of NIR to assess lignin composition (H/G ratio) in maritime pine wood using analytical pyrolysis as the reference method," *Holzforschung* 60(1), 29-31.
- Alves, A., Simoes, R., Stackpole, D. J., Vaillancourt, R. E., Potts, B. M., Schwanninger, M., and Rodrigues, J. (2011). "Determination of the syringyl/guaiacyl ratio of *Eucalyptus globulus* wood lignin by near infrared-based partial least squares regression models using analytical pyrolysis as the reference method," *J. Near Infrared Spectrosc.* 19(5), 343-348.
- Alves, A. M. M., Simões, R. F. S., Santos, C. A., and Potts, B. M. (2012). "Determination of *Eucalyptus globulus* wood extractives content by near infrared-based partial least squares regression models: Comparison between extraction procedures," *J. Near Infrared Spectrosc.* __, 1-11.
- Barbosa, L. C. A., Maltha, C. R. A., Silva, V. L., and Colodette, J. L. (2008). "Determination of the syringyl/guaiacyl ratio in eucalyptus wood by pyrolysis-gas chromatography/mass spectrometry (Py-GC/MS)," *Quimica Nova* 31(8), 2035-2041.
- Baucher, M., Halpin, C., and Petit-Conil, M. (2003). "Lignin: Genetic engineering and impact on pulping," *Crit. Rev. Biochem. Mol.* 305-350.
- Besseau, S., Hoffmann, L., Geoffroy, P., et al. (2007). "Flavonoid accumulation in *Arabidopsis* repressed in lignin synthesis affects auxin transport and plant growth," *Plant Cell* 19, 148-162.
- Boudet, A. M., Kajita, S., Grima-Pettenati, J., and Goffner, D. (2003). "Lignins and lignocellulosics: A better control of synthesis for new and improved uses," *Trends Plant Sci.* 8, 576-581.
- Chen, W.-J., Yun, M.-S., Deng, F., and Yogo, Y. (2011). "Chalcone suppresses lignin biosynthesis in illuminated soybean cells," *Wedd. Biol. Manag.* 11, 49-56.
- Campbell, M. M., and Sederoff, R. R. (1996). "Variation in lignin content and composition (mechanisms of control and implications for the genetic improvement of plants)," *Plant Physiol* 110, 3-13.
- Demura, T., and Fukuda, H. (2007). "Transcriptional regulation in wood formation," *Trends Plant Sci.* 12, 64-70.
- Deng, F., Aoki, M., and Yogo, Y. (2004). "Effect of naringenin on the growth and lignin biosynthesis of gramineous plants," *Wedd. Biol. Manag.* 4, 49-55.
- Dixon, R. A., Lamb, C. J., Masoud, S., Sewalt, V. J., and Paiva, N. L. (1996). "Metabolic engineering: Prospects for crop improvement through the genetic manipulation of phenylpropanoid biosynthesis and defense responses — A review," *Gene* 179, 61-71.
- Foucart, C., Jauneau, A., Gion, J.-M., Amelot, N., Martinez, Y., Panegos, P., Grimá Pettenati, J., and Sivadon, P. (2009). "Overexpression of EgROP1, a *Eucalyptus* vascular-expressed Rac-like small GTPase, affects secondary xylem formation in *Arabidopsis thaliana*," *New Phytol.* 183, 1014-1029.
- Gerlach, D. (1969). *Botanische Mikrotechnik: Eine Einführung*, Georg Thieme, Stuttgart
- Godoy, E. A., Rodrigues, J., Alves, A. M. M., and Lazo, D. A. (2007). "Content and quality study of the lignin by analytical pyrolysis in *Pinus caribaea*," *Maderas-Ciencia Y Tecnologia* 9(2), 179-188.
- Gomide, J. L., Colodette, J. L., Oliveira, R. C., and Silva, C. M. (2005). "Caracterização o tecnológica, para produção de celulose, da nova geração de clones de *Eucalyptus* do Brasil," *Rev Arvore* 29, 129-137.

- Gominho, J., Figueira, J., Rodrigues, J., and Pereira, H. (2001). "Within-tree variation of heartwood, extractives and wood density in the eucalypt hybrid urograndis (*Eucalyptus grandis* x *E-urophylla*)," *Wood Fiber Sci.* 33(1), 3-8.
- Grattapaglia, D., and Kirst, M. (2008). "Eucalyptus applied genomics: From gene sequences to breeding tools," *New Phytol.* 179, 911-929.
- Hu, W. J., Harding, S., Lung, J., *et al.* (1999). "Repression of lignin biosynthesis promotes cellulose accumulation and growth in transgenic trees," *Nat. Biotechnol.* 17, 808-812.
- Huntley, S. K., Ellis, D., Gilbert, M., *et al.* (2003). "Significant increases in pulping efficiency in C4H-F5H-transformed poplars: Improved chemical savings and reduced environmental toxins," *J. Agr. Food Chem.* 51, 6178-6183.
- Johansen, D. A. (1940). *Plant Microtechnique*, McGraw-Hill, New York
- Jung, H.-J. G., Samac, D. A., and Sarath, G. (2011). "Modifying crops to increase cell wall digestibility," *Plant Sci.* 185-186, 65-77.
- Lima, C. F., Barbosa, L. C. A., Marcelo, C. R., Silverio, F. O., and Colodette, J. L. (2008). "Comparison between analytical pyrolysis and nitrobenzene oxidation for determination of syringyl/guaiacyl ratio in *Eucalyptus* spp. lignin." *BioResources* 3(3), 701-712.
- Mansfield, S. D. (2009). "Solutions for dissolution – Engineering cell walls for deconstruction," *Curr. Opin. Biotech.* 20, 286-294.
- Moustafa, E. (1967). "Purification and properties of chalcone-flavanone isomerase from soya bean seed," *Phytochemistry* 6, 625-632.
- Paiva, J. A. P., Garnier-Gere, P. H., Rodrigues, J. C., Alves, A., Santos, S., Graca, J., LeProvost, G., Chaumeil, P., Da Silva-Perez, D., Bosc, A., Fevereiro, P., and Plomion, C. (2008). "Plasticity of maritime pine (*Pinus pinaster*) wood-forming tissues during a growing season," *New Phytologist* 179(4), 1080-1094.
- Rippert, P., Puyaubert, J., Grisolle, D., *et al.* (2009). "Tyrosine and phenylalanine are synthesized within the plastids in *Arabidopsis*," *Plant Physiol.* 149, 1251-1260.
- Rodrigues, J., Meier, D., Faix, O., and Pereira, H. (1999). "Determination of tree to tree variation in syringyl/guaiacyl ratio of *Eucalyptus globulus* wood lignin by analytical pyrolysis," *J. Anal. Appl. Pyrol.* 48(2), 121-128.
- Rodrigues, J., Graca, J., and Pereira, H. (2001). "Influence of tree eccentric growth on syringyl/guaiacyl ratio in *Eucalyptus globulus* wood lignin assessed by analytical pyrolysis," *J. Anal. Appl. Pyrol.* 58, 481-489.
- Sarruje, J. (1975). "Soluções nutritivas," *Summa Phytopathol.* 1, 231-233.
- Schwanninger, M., and Hinterstoisser, B. (2002). "Klason lignin: Modifications to improve the precision of the standardized determination," *Holzforschung* 56, 161-166.
- Stackpole, D. J., Vaillancourt, R. E., Downes, G. M., Harwood, C. E., and Potts, B. M. (2010). "Genetic control of kraft pulp yield in *Eucalyptus globules*," *Can. J. Forest Res.* 40, 917-927.
- Stewart, J. J., Kadla, J. F., and Mansfield, S. D. (2006). "The influence of lignin chemistry and ultrastructure on the pulping efficiency of clonal aspen (*Populus tremuloides* Michx.)," *Holzforschung* 60, 111-122.
- TAPPI Test Methods T222om-85: "Acid-insoluble lignin in wood and pulp," In: *TAPPI Test Methods 1994-1995*, TAPPI Press, Atlanta, GA, USA.
- TAPPI T 222 om-02, "Acid-insoluble lignin in wood and pulp," In: *2002-2003 TAPPI Test Methods*, 2002, TAPPI Press, Atlanta, GA, USA.

- TAPPI UM 250, "Acid-soluble lignin in wood and pulp," in: *1991 TAPPI Useful Methods*, TAPPI Press, Atlanta, GA, USA.
- Valério, L., Carter, D., Rodrigues, J., Tournier, V., Gominho, J., Marque, C., Boudet, A. M., Maunders, M., Pereira, H., and Teulieres, C. (2003). "Down regulation of cinnamyl alcohol dehydrogenase, a lignification enzyme, in *Eucalyptus camaldulensis*," *Mol. Breeding* 12(2), 157-167.
- Vanholme, R., Demedts, B., Morreel, K., *et al.* (2010). "Lignin biosynthesis and structure," *Plant Physiol* 153, 895-905.
- Vogt, T. (2010). "Phenylpropanoid biosynthesis," *Mol. Plant* 3, 2-20.
- Voo, K. S., Whetten, R. W., O'Malley, D. M., and Sederoff, R. R. (1995). "4-coumarate:coenzyme A ligase from loblolly pine xylem. Isolation, characterization, and complementary DNA cloning," *Plant Physiol.* 108, 85-97.
- Weng, J.-K., Li, X., Bonawitz, N. D., and Chapple, C. (2008). "Emerging strategies of lignin engineering and degradation for cellulosic biofuel production," *Curr. Opin. Biotech.* 19, 166-172.
- Yun, M.-S., Chen, W., Deng, F., and Yogo, Y. (2009). "Selective growth suppression of five annual plant species by chalcone and naringenin correlates with the total amount of 4-coumarate: Coenzyme A ligase," *Wedd. Biol. Manag.* 9, 27-37.

Article submitted: October 31, 2012;

Capitulo 2: Análise da expressão gênica e hidrólise enzimática

Flavonoid supplementation affects the expression of genes involved in cell wall formation and lignification metabolism and increases sugar content and saccharification in the fast-growing *Eucalyptus* hybrid *E. urophylla* x *E. grandis*

Lepikson-Neto, J.¹, Nascimento L.C.¹, Salazar M.M.¹, Camargo E.L.O.¹, Cairo, J.P.F.²; Teixeira P.J.¹, Marques, W.L.¹; Squina F.M.²; Mieczkowski, P.³; Deckmann, A.C.¹ and Pereira, G.A.G.¹

1 - Laboratório de Genômica e Expressão, Departamento de Genética e Evolução, Instituto de Biologia, Universidade Estadual de Campinas.

2 - Laboratório Nacional de Ciência e Tecnologia do Bioetanol, CTBE, Campinas, São Paulo, Brazil

3 - Department of Genetics, School of Medicine, University of North Carolina at Chapel Hill (UNC), Chapel Hill, NC, USA

ABSTRACT

Eucalyptus species are the most widely planted hardwood species in the world and are renowned for their rapid growth and adaptability. These species supply the timber industry with high-quality material for producing pulp, paper, wood and biomass while preserving natural forests. In Brazil, one of the most widely grown *Eucalyptus* cultivars is the fast-growing *Eucalyptus urophylla* x *Eucalyptus grandis* hybrid. In a previous study, we described a chemical characterization of these hybrids when subjected to flavonoid supplementation on 2 distinct timetables, and our results revealed marked differences between the wood composition of the treated and untreated trees. In this work, we report the transcriptional responses occurring in these trees that may be related to the observed chemical differences. Gene expression was analyzed through mRNA sequencing, and the results showed that the supplemented trees exhibit a distinct expression profile in comparison with the control group. Notably, compared to control trees, the treated trees display differential down-regulation of cell wall formation pathways such as phenylpropanoid metabolism as well as differential expression of genes involved in sucrose, starch and minor CHO metabolism and genes that play a role in several stress and environmental responses. Because phenylpropanoid metabolism is involved in lignin and flavonoid biosynthesis, these results may provide clues regarding the molecular events that determine wood quality in *Eucalyptus* trees. We also performed enzymatic hydrolysis of wood samples from the different treatments, and the results indicated higher sugar contents and glucose yields in the flavonoid-treated plants, further illustrating the potential use of flavonoids as a nutritional complement for modifying *Eucalyptus* wood.

Keywords: *Eucalyptus*; lignin, phenylpropanoid metabolism, syringyl/guaiacyl ratio, gene expression, hydrolysis.

Contact information: Laboratório de Genômica e Expressão, Departamento de Genética Evolução e Bioagentes, Instituto de Biologia, Universidade Estadual de Campinas, CEP: 13083-970, Campinas, São Paulo, Brazil; *Corresponding author: goncalo@unicamp.br

Introduction

Trees constitute the majority of the lignocellulosic biomass on Earth and are expected to play a significant role in the future as a renewable and environmentally cost-effective alternative feedstock for biofuel production, a source of fibers and solid wood products and a major sink for excess atmospheric CO₂ (Boudet et al. 2003; Plomion et al. 2001; Rengel et al. 2009). In Brazil, the pulp and paper industries have been efficiently fed by *Eucalyptus* forests due to their rapid growth, adaptability and wood quality, but with the dramatic increase in industrial demands and the interest in second-generation biofuels and renewable chemicals, the quality and quantity of wood produced must also increase (Paiva et al. 2011; Grattapaglia et al. 2012).

Wood is a highly variable material that differs among trees and is composed of the secondary xylem, a specialized type of conductive and structural support tissue produced through the lateral growth and differentiation of the meristematic vascular cambium (Fukuda 1996). Most of the genes expressed during the formation of the secondary xylem (xylogenesis) are involved in determining the physical and chemical properties of wood (Plomion et al. 2001; Paux et al. 2005). Despite the progress that has been made in defining the molecular and cellular events involved in xylogenesis, the mechanisms regulating the rate of this process and the variation in wood properties remain largely unknown (Kirst et al. 2004; Foucart et al. 2009; Vanholme et al. 2010).

The secondary xylem cell wall of *Eucalyptus* trees is mostly composed by cellulose (β -1,4-glucan), lignin (a phenolic polymer) and hemicelluloses (heterogeneous polysaccharides), in an approximate ratio of 2:1:1 (Hu et al. 1999). During tree growth, cellulose microfibrils give the cell walls tensile strength, and the lignin encasing the cellulose microfibrils imparts rigidity to the cell walls. Despite its importance during growth, lignin becomes problematic during postharvest, cellulose-based wood processing because it must be extracted during industrial handling through a complicated process, resulting in an enormous expenditure of energy and chemicals and strain on the environment (Hu et al. 1999; Fu et al. 2010). Thus, it is of major interest to investigate the molecular basis of lignification to further increase our overall comprehension of this metabolic process for better adaptation of industrial processes.

Lignin synthesis is a relatively well-understood process that begins with the assembly of radicals produced during the single-electron oxidation of monolignols (Baucher et al. 2003; Li et al. 2010; Vanholme et al. 2010). The industrial exploitation of wood to obtain cellulose depends mostly on the composition of lignins because lignins determine the rigidity of the wood and the feasibility of cellulose extraction, which are of major concern in the paper and pulp industries. In angiosperms, lignin is composed of 2 major units: the guaiacyl (G) and syringyl (S) units, which are derived from corresponding monolignol precursors, the coniferyl and sinapyl alcohols, respectively (Boudet 2000; Boudet et al. 2003). The S/G ratio dictates the degree and nature of polymeric cross-linking; an increased G content leads to highly cross-linked lignin (more rigid

wood), whereas S subunits are typically linked through more labile ether bonds at the 4-hydroxyl position (Dixon et al. 1996; Guo et al. 2001; Ferrer et al. 2008;). Thus, S-rich lignins are much easier to dissociate from cellulosic content, resulting in a much cleaner and cheaper process (Ferrer et al. 2008). The S/G ratio is variable among species and is commonly used to evaluate the quality of wood in commercial tree plantations (Rodrigues et al. 1999; Foelkel 2009;).

The formation of lignin monomers begins with the catalytic step performed by the 4-coumaroyl:CoA-ligase (4CL) enzyme, which likely represents the most important branch point in the central phenylpropanoid biosynthesis pathway in plants (Voo et al. 1995; Campbell and Sederoff 1996). Through 4CL activity, cells can produce the precursors for either flavonoids or the G and S lignin precursors (Vogt 2010). The product of 4CL, p-coumaroyl-CoA, is the substrate of the enzyme chalcone synthase (CHS) (Besseau et al. 2007), which carries out the committing step in flavonoid biosynthesis. This pathway is reviewed in detail elsewhere (Besseau et al. 2007; Vanholme et al. 2010).

The flavonoids naringenin-chalcone and naringenin, which are synthesized by the enzymes chalcone synthase (CHS) and chalcone isomerase (CHI), respectively, are the primary C15 intermediates in flavonoid biosynthesis (Moustafa 1967; Chen et al. 2011). This metabolic pathway appears to be a promising target for improving wood quality in *Eucalyptus* trees, as shown by our previous work (Lepikson-Neto et al. 2013) demonstrating that flavonoid supplementation of the fast-growing *Eucalyptus urophylla* x *Eucalyptus grandis* hybrid, hereafter referred to as *E. urograndis*, changes its wood composition, reduces its extractive contents and alters its syringyl monomer composition.

In this context, the objective of the present work was to perform further studies on the effects of flavonoid supplementation on *E. urograndis* trees by analyzing gene expression in xylem tissue from treated and non-treated trees and by measuring the effect on sugar accessibility through enzymatic hydrolysis. We analyzed the obtained data with special emphasis on results that might be correlated with the previously observed changes in wood composition (Lepikson-Neto et al. 2013).

Materials and Methods

Plant materials and tissue harvesting

Plantlets of a 6-month-old commercial clone of *Eucalyptus urograndis* were provided by International Paper (Mogi-Guaçu, Brazil) and grown in a greenhouse. The plantlets were divided into 5 groups according to supplementation conditions (apart from the standard nutritional solution supplied to all groups), as follows: control group (CT); experimental group 1 (CH), supplemented with 0.1 mmol of naringenin-chalcone for 5 months; experimental group 2 (NAR), supplemented with 0.1 mmol of naringenin for 5 months; experimental group 3 (CHSTOP), supplemented with 0.1 mmol of naringenin-chalcone for only the first month and then given the standard nutrition solution for the next 5 months; and experimental group 4 (NARSTOP), supplemented with 0.1 mmol of naringenin for only the first month and then given the standard nutrition solution for the next 5 months. Approximately 100-150 mL of each solution was administered via root application daily. The treatments lasted 5 months. The composition of the standard nutritional solution has been described previously (Sarruje, 1975). At the end of the experiment, all 5 groups of plantlets were cut, and their stems were debarked, immediately frozen in liquid nitrogen and kept at -80°C for analysis; no growth differences were observed between the control and treatment groups. All samples were analyzed 5 months after the beginning of the experiment, regardless of the applied supplementation.

Naringenin (4',5-,7-trihydroxyflavanone, 95%) and naringenin-chalcone (1,3-diphenyl-2-propen-1-one, 97%) were purchased from Sigma-Aldrich Co. (Tokyo, Japan) and AcrosOrganics Co. (Tokyo, Japan), respectively.

Total RNA extraction

Total RNA was extracted according to the protocol described by Zeng and Yang (2002), with the modifications proposed by Le Provost et al. (2003). The obtained RNA concentration and quality were verified using a Nanodrop 2000 spectrophotometer (Thermo Scientific).

mRNA sequencing

mRNA sequencing was performed at the High-Throughput Sequencing Facility of the Carolina Center for Genome Sciences (University of North Carolina, USA). From each xylem sample, 10 µg of total RNA was used to prepare an mRNAseq library according to the protocol provided by Illumina. The gel extraction step was modified by dissolving excised gel slices at room temperature to avoid underrepresentation of AT-rich sequences (Quail et al. 2008). Quality control and quantification of the libraries were performed using a DNA 1000 series II Bioanalyzer Chip (Agilent). For each library, single-end sequences of 36 or 50 bp were generated in a single lane using an Illumina Genome Analyzer IIx. A total of 8 libraries were generated: 3 biological replicates of the control group (CT); 2 biological replicates of the 5-month naringenin-supplemented groups (NAR); and 1 library for each remaining group (subjected to 1 month of supplementation with naringenin (NARSTOP), 5 months of supplementation with naringenin-chalcone (CH) or 1 month of supplementation with naringenin-chalcone (CHSTOP). Each library was constructed from a sample pooled from 3 individual trees.

Read alignment

The obtained Illumina reads were filtered to exclude ribosomal sequences (using the SILVA database (Pruesse et al. 2007) and low quality reads (phred ≥ 20). The remaining reads were aligned against the greater splice variants of *E. grandis* transcripts from Phytozome 7.0 (44,974 sequences) using the SOAP2 alignment software package (Li et al. 2009). To prepare the data for Genebrowser analysis, the reads were aligned to the *E. grandis* genome using the TopHat aligner (Trapnell et al. 2009) to allow for spliced alignments. Both programs were configured to allow up to two mismatches (because SNPs can generate mismatches in the alignments, especially in cases such as the present analysis, where the sequences come from different

species), discard sequences with ambiguities (Ns) and return only reads with unique alignments.

Gene annotation

The Autofact program (Koski et al. 2005) was used to perform an automatic annotation of all *E. grandis* transcripts. The main feature of Autofact is its ability to perform gene annotation based on sequence similarity searches of several databases. BLASTx (Altschul et al. 1997) (e-value cutoff of 1e-5) was used to align the obtained contigs against the following public databases: the NCBI non-redundant (NR) database; the Uniref90 and Uniref100 databases, which contain clustered sets of proteins from Uniprot (Supek et al. 2007); the KEGG database of metabolic pathways (Kanehisa 2000); and TAIR (version 10), an *Arabidopsis* proteins database. Functional annotation (GO) was performed using BLAST2GO (Conesa et al. 2005) and MaPMAN with the default parameters.

Determination of gene expression levels

Gene expression was measured via the FPKM (fragments per kilobase of exon per million fragments mapped) method using only one read alignment for each transcript, as described by Mortazavi et al. (2008). The FPKM values for all transcripts are available for searching in the EUCANEXT database (www.lge.ibi.unicamp.br/eucalyptusdb).

Enzymatic Hydrolysis

Substrate preparation

Samples from each Eucalyptus treatment were frozen in liquid nitrogen and then dried in FreeZone6 (Labconco) at - 51°C and 25 Pa for 48 hours. Subsequently, the lignocellulosic material was reduced through 1 cycle of 5 minutes in a ball-milling reactor. The milled material was used as a substrate for fungal growth and hydrolysis assays.

Secretome production for enzymatic hydrolysis

The *Neurospora crassa* wide strain St.L. 74A (Missouri University, Kansas City, <http://www.fgsc.net/>) was used for secretome production. Conidia preparation was performed by inoculating the fungus in 100 mL of minimal medium plus Vogel's salts supplemented with 143 µL of biotin (biotin 5 mg, ethanol 50% (v/v), 143 µL of a trace element solution (5 g monohydrate citric acid, 5 g of ZnSO₄·7H₂O, 1 g of Fe(NH₄)₂·6H₂O, 0.25 g of CuSO₄·5H₂O, 0.05 g MnSO₄·H₂O, 0.05 g H₃BO₃, 0.05 g Na₂MoO₄·2H₂O, qsp 1,000 mL), 1.5% agar and 2% sucrose. The prepared samples were grown for 3 days at 30°C in the dark and then for 7 days in the light at 25°C. A conidial suspension was then inoculated in 100 mL of the same medium described above without agar (Phillips et al. 2011) containing as the only carbon source 2% of a substrate blend of 3 *Eucalyptus* species: *E. grandis*, *E. urograndis* and *E. urophylla*, in a ratio of 3:3:1, prepared as described above.

Eucalyptus hydrolysis

Hydrolysis was performed as described by Bragatto et al. (2012) with the following modifications: enzymatic hydrolysis was performed in 2 mL tubes shaken at 1,000 rpm at 30°C in a Thermomixer (Eppendorf) for 48 hours. Approximately 10 mg of substrate from each substrate preparation was diluted in 400 µL of 50 mM sodium acetate buffer pH 5.5, and 100 µL of the *N. crassa* secretome was then added. The protein concentration of the secretome was 0.4 µg/µL, as determined in a Bradford Kit assay (BioRad) with BSA as a standard. This temperature and pH were optimal for the hydrolysis of carboxymethyl cellulose, xylan and β-

glucan, as determined by testing the temperatures of 25-40 °C and pH levels of 4.0–9.0. All hydrolysis reactions were performed in triplicate. For determination of the reducing sugar concentration and glucose production in supernatants derived from *Eucalyptus* hydrolysis, 2 mL tubes were centrifuged at 20,000 x g for 10 minutes at 4°C. The supernatant was then recovered, and 100 µL of each reaction was used to determine the content of reducing sugars by adding 100 µL of the dinitrosalicylic acid (DNS) assay reagent (Miller 1959) heated to 99 °C. A 100 µL aliquot of the sample was next transferred to an ELISA plate, and its absorbance was measured at 540 nm using a Tecan Infinite M200 microplate reader, referring to calibration curves generated from glucose solutions. To calculate glucose contents, 20 µL of the supernatant was added to 100 µL of a working solution from a Glucose Oxidase Kit (Laborlab) in ELISA plates. The reaction was subsequently incubated at 37°C for 10 minutes, and its absorbance was measured at 505 nm (in a Tecan Infinite M200 microplate reader). Glucose concentrations were calculated with a factor referring to a standard solution of glucose at 1 mg/mL. A blank reaction containing only buffer and substrate was subtracted from the measurements obtained for each assay.

Results

RNA sequencing and differential gene expression

A total of over 335 million reads were generated from 8 samples: 3 samples from the control group (CT); 3 from the naringenin-supplemented groups (2 NAR and 1 NARSTOP); and 2 from the naringenin-chalcone supplemented groups (1 CH and 1 CHSTOP). The number of reads per sample ranged from 32 to 54 million (total) and 30 to 48 million (after filtering). The reads were mapped against the greater splice variants (44,974 sequences) of the *E.grandis* gene predictions from Phytozome 7.0 (54,935 transcripts) using the SOAP2 alignment software package (Li et al., 2009) (Supplementary file 1).

Heat map clustering of all transcripts was performed using Expander software (Shamir et al. 2005), resulting in 2 major groups: 1 formed by the 3 control sample replicates and the other by the flavonoid-supplemented samples (Figure 1).

The read counts from each sample were used to test the differential expression of the genes between the control (CT) and supplemented (CH, NAR, CHSTOP and NARSTOP) treatments using the baySeq package (Hardcastle and Kelly 2010). A total of 1,573 genes were considered to be differentially expressed ($FDR < 0.01$), which were distributed among the treatments (917 CH; 1,289 NAR; 268 CHSTOP; 47 NARSTOP) (Supplementary file 2).

The gene expression patterns observed for the supplemented and control groups were distinct, while similar profiles were observed within treatments, indicating similarities among the different types of flavonoid supplementation studied here. Most of the differences were observed in the long-term supplementation treatments, which comprised almost all of the genes that were differentially expressed in the short-term treatments as well. The NAR-supplemented plants displayed the greatest number of genes that were differentially expressed, while the NARSTOP-supplemented plants had fewer, which may indicate that naringenin supplementation has a strong, but short-lasting impact on gene expression, whereas naringenin-chalcone has a smaller but more durable impact.

Functional analyses

To determine the biological functions of the genes responding to flavonoid supplementation, functional analyses were performed using the web-based tools Blast2GO and Mapman. The genes considered differentially expressed in each treatment were mapped to their corresponding metabolic pathways, and the treatments were tested for enrichment of particular metabolic responses.

Only 36 genes were differentially expressed in all four treatments, including genes encoding several heat-shock proteins, sequences with no hits and unknown proteins (Table 1).

Each supplemented group was analyzed individually. Common categories between different treatments are shown in Figure 2, and all affected GO categories are listed in Supplementary file 3.

Many of the down-regulated categories that were common to all treatments are involved in cell wall formation and development. On the other hand, the common up-regulated categories are all related to stress and environmental responses. Interestingly, NARSTOP, which resulted in fewer differentially expressed genes, only led to enriched GO categories among up-regulated genes.

Mapman analyses of all of the differentially expressed genes also indicated down-regulation of cell wall-related genes and phenylpropanoid pathways, whereas flavonoid, minor CHO and starch and sucrose metabolism and stress response were associated with the most genes up-regulated (Figure3).

The phenylpropanoid genes

To further analyze the impact of flavonoid supplementation on lignification, a broader analysis was performed on the genes from the phenylpropanoid pathway, especially those related to lignin biosynthesis.

Several phenylpropanoid genes were differentially expressed between the treated samples and controls (Table 2), including the following genes that are directly related to lignin synthesis: 4CL, HCT, 2 OMT-methyltransferases, CCR and 2 CAD genes; 4CL, HCT and CCR were down-regulated, while the 2 methyltransferases and CAD genes were up-regulated. Additionally, several laccases were down-regulated among the treatments. These results are highly significant in terms of explaining the higher S/G ratio found in supplemented plants.

Interestingly, no gene related to the phenylpropanoid pathway was differentially expressed as a result of NARSTOP treatment.

Secondary cell wall genes

In addition to genes from the phenylpropanoid pathway, many genes related to secondary cell wall formation were differentially expressed in response to flavonoid supplementation (Table 3). Among these genes, we observed sucrose synthases, cellulose synthases and many glucosylases and transferases, most of which were down-regulated following the prolonged supplementation treatments. However, we also observed the up-regulation of several genes related to secondary cell wall formation after both prolonged and short-term flavonoid supplementation, including galactinol synthase, stachyose synthase, raffinose synthase and starch synthase.

Stress-related genes

Some of the most differentially expressed genes belonged to stress-related gene categories, which were up-regulated in all of the supplemented groups. These genes included several encoding heat-shock proteins and UDP-glycosyl transferases (Table 4).

Enzymatic Hydrolysis

To verify the effects of flavonoid supplementation on sugar yields and saccharification in *Eucalyptus* wood, enzymatic hydrolysis was performed. The hydrolysates were analyzed for total sugar contents ('reduced sugars'), which included most of the pentoses and hexoses from the hemicellulose fraction, and glucose content ('glucose'), allowing an estimate of the percent of saccharification to be obtained.

Flavonoid-supplemented plantlets showed increased sugar and glucose values compared to the control groups. The reduced sugar content was increased from 50% (CH) to 250% (NARSTOP), and the glucose content was increased from 43% (CH) to 253% (NARSTOP). With the exception of the naringenin-chalcone prolonged supplementation treatment (CH), all of the treatment values were considered statistically significant (Table 5).

Discussion

The metabolism of phenylpropanoids follows 2 main pathways: the lignin branch and the flavonoid branch. The two pathways share common substrates and enzymes, and these shared components lead to a high level of interdependence between the pathways. Considering the economic interest in *Eucalyptus* trees for paper and pulp production, and given that flavonoids are known to have a direct influence on lignification and wood formation in several species (Chen et al. 2004; Deng et al. 2004), including *Eucalyptus* species, as previously demonstrated by our group (Lepikson-Neto et al 2013), it is of high interest to verify the effects of flavonoid supplementation on gene expression, especially concerning genes related to wood formation. Additionally, there is a pressing interest in expanding the industrial uses of *Eucalyptus* because *Eucalyptus* forest cultures are well-established in Brazil and may affect other strategic sectors, such as second-generation biochemicals. In this case, *Eucalyptus* wood could be employed as lignocellulosic biomass for biological fermentation (Mizrachi et al. 2011; Mansfield et al. 2012).

With this objective, we designed the present work to investigate the molecular basis of the differences in wood observed in flavonoid-supplemented *E. urograndis* trees. Additionally, in light of our previous findings, we paid special attention to the expression of genes involved with lignin and secondary cell wall formation and to the possible association between gene expression and the chemical composition of wood in *Eucalyptus*.

We analyzed the whole genome (44,974 genes) of *Eucalyptus* plants following supplementation with different flavonoids. A total of 1,573 (3,5%) differentially expressed genes were identified, which were distributed among the supplementation groups: 963 genes were down-regulated and 610 genes were up-regulated. Most of the differentially expressed genes were associated with the prolonged supplementation groups (1,289 for NAR and 917 for CH), while the short-term supplementation groups displayed fewer differentially expressed genes (268 for CHSTOP and 47 for NARSTOP). Most of the differentially expressed genes in the CHSTOP and NARSTOP groups were also differentially expressed in the NAR and CH groups. Thus, naringenin supplementation appears to have had a stronger but less durable effect, while naringenin-chalcone supplementation has a longer-lasting effect on gene expression.

GO enrichment analyses demonstrated that there were several categories involved in cell wall formation that were down-regulated in all of the supplemented groups, including the phenylpropanoid pathway in the NAR-supplemented samples. The up-regulated gene categories included many responses to stress and the environment as well as genes related to sugar alcohols, through being involved in polyol, hexitol and alditol metabolism (minor CHOs), in the CH group. This pattern could also be observed in the mapping analysis of differentially expressed genes performed using MapMan software, in which several pathways, most notably those associated with the cell wall and phenylpropanoids, were down-regulated, while the metabolic pathways associated with minor CHOs, flavonoids, sucrose and starch displayed up-regulated genes. Furthermore, there was strong evidence that stress may play a major role, as several stress-related gene categories were found to be enriched via GO analysis, even in the groups subjected to short-term supplementation.

It was therefore clear that lignification and the phenylpropanoid pathway are affected by a great number of factors, and we believe that our work can help to clarify some of these factors. The interdependence of the phenylpropanoid, flavonoid and lignin branches has been explored in other studies. For example, it has been reported that 4CL activity is inhibited by some flavonoids, such as naringenin-chalcone and naringenin, which are the products of the chalcone synthase (CHS) and chalcone isomerase (CHI) enzymes, respectively (Chen et al. 2004; Chen et al. 2005). The same study demonstrated that the administration of flavonoids suppressed the growth of 20 plant species, although the sensitivities of the plants to flavonoids were different.

In addition, the activation of the lignin precursor cinnamic acid (catalyzed by C4H) and p-coumaroyl-CoA (catalyzed by 4CL) is, to some extent, regulated by the activity of the CHS enzyme, which is involved in the first step of flavonoid biosynthesis (Chen et al. 2005). It has also been reported that CHS is associated with growth suppression via the regulation of 4CL. This association has major importance in lignin biosynthesis in a great number of species (Deng et al. 2004; Chen et al. 2005).

As demonstrated by our results, several genes involved in the phenylpropanoid pathway were differentially expressed in plants subjected to supplementation with flavonoids (Table 2; Figure3). Our most noteworthy findings revealed the differential expression of genes directly related to lignin synthesis. The NAR-supplemented group presented down-regulation of both the 4CL and CCR genes, whereas the ATOMT1 and 2 CAD genes were up-regulated. The CH-supplemented group exhibited HCT down-regulation and 1 CAD gene that was up-regulated. In the CHSTOP-supplemented group, 1 methyltransferase was up-regulated. No genes from the phenylpropanoid pathway were differentially expressed following supplementation with NARSTOP.

Surprisingly, the gene encoding F5H, which is one of the key enzymes involved in the synthesis of the monolignol sinapyl alcohol and, ultimately, the S lignin moiety, was not found to be differentially expressed on our analyses. This result is particularly interesting in light of our finding that the S/G ratios in all of the flavonoid-supplemented groups were higher than that of the control group. Thus, we expected a change in the expression of F5H following flavonoid treatment. Because phenylpropanoid metabolism is complex, it is likely that the differential regulation of other enzymatic steps, such as those encoded by the 4CL, HCT, CCR, ATOM1 and CAD genes, may underlie this response.

Some findings reported in the literature support this possibility. For example, 4CL plays a major role in phenylpropanoid metabolism, as its product, p-coumaroyl-CoA, is a substrate that is common to the flavonoid and lignin synthesis pathway. HCT silencing in *Arabidopsis* represses lignin synthesis and plant growth, and the metabolic flux is redirected toward flavonoids by chalcone synthase activity (Besseau et al. 2007). CCR catalyzes the reduction of hydroxycinnamoyl-CoA thioesters to the corresponding aldehydes; this reaction is considered to be a potential control point that regulates the overall carbon flux in favor of lignin (Lacombe et al. 1997). *Arabidopsis* ATOMT1 knock-out mutants lack S units (Goujon et al. 2003), and CAD catalyzes the reduction of cinnamaldehydes to cinnamyl alcohols, which is the last step in the biosynthesis of the monolignols, thus playing a pivotal role in determining the lignin monomer composition and increasing S contents (Baucher et al. 2003).

There are also several laccases that have been demonstrated to be involved in lignification (Berthet et al. 2011), and many laccases were found to be down-regulated in the NAR-, CH- and CHSTOP-supplemented samples.

Our results further corroborate those of Li et al. (1993), who suggested that *Arabidopsis* responds to the accumulation of 1 or more intermediates from the flavonoid pathway by down-regulating either the whole phenylpropanoid pathway or the specific branch leading to monocyclic phenolic compounds. According to our results, it is possible that the accumulation of naringenin-chalcone and naringenin, the products of CHS and CHI, respectively, due to exogenous supplementation, results in the down-regulation of genes from the phenylpropanoid

pathway, with the exception of 2 genes involved in the final steps of sinapilic acid synthesis (ATOMT1 and CAD). This down-regulation may at least partially explain the higher S/G ratios observed in the supplemented samples.

While the NARSTOP-supplemented plants did not show differential expression of any genes that are related to lignin synthesis according to our statistical analyses, they exhibited FPKM values that were similar to those of the NAR-, CH- and CHSTOP-supplemented groups, but closer to the control values than the other groups. This indicates that an early impact on gene expression may be sufficient to promote the phenotypic differences observed in this group.

Another possibility is that factors other than the genes from the lignification pathway *per se* influence the lignin monomer composition. Cook and collaborators (2012) reported that the levels of cellulose, xylan and lignin are not completely dependent on the transcription of the genes involved in these metabolic pathways. Thus, the regulation of cell wall biosynthesis occurs at different levels, not only at the transcriptional level (Cook et al. 2012).

Additionally, other genes that have not yet been discovered may be causing the observed differences, as many no hits and unknown proteins were found among the most differentially expressed genes following flavonoid treatment. The stress and environmental response pathways were significantly enriched and associated with lignification; thus, these pathways may play major roles in the alterations of lignin composition after flavonoid supplementation.

Stress and lignification are closely related. Many of the products of the phenylpropanoid pathway are induced by biotic and abiotic stress (Dixon and Paiva 1995). Both flavonoids and sinapate esters, which are used for lignin synthesis, are important for UV protection (Li et al. 1993). *Arabidopsis* mutants with reduced levels of CHS and CHI activity show up to 60% higher levels of sinapate esters (Li et al. 1993; Dixon et al 95).

Moreover, a large number of phenylpropanoids are induced by stress, such as those derived from the C15 flavonoid skeleton that are synthesized via the chalcone synthase (CHS)-mediated condensation of p-coumaroyl-coenzyme A (CoA) and three molecules of malonyl-CoA (Harbone 1988). In most plant families, the initial product of CHS is a tetrahydroxychalcone, which is further converted to other flavonoid classes, such as flavones, flavanones, flavanols, anthocyanins and 3-deoxyanthocyanidins, all of which are compounds that are important in the response to stress (Holton and Cornish 1995).

The observation that several genes related to stress responses are differentially expressed in flavonoid-supplemented trees confirms the importance of stress responses in defining *Eucalyptus* wood properties as has been previously shown by our group (Salazar et al 2013). In this work, a comparison between three *Eucalyptus* species revealed differential expression of stress-related genes in *E. urophylla*, which could explain the higher plasticity and

adaptability of this species compared to *E. grandis* and *E. globulus*, the two other studied species.

In the present study, the stress-related genes that were differentially expressed following flavonoid treatments included a noteworthy group composed of several UDP-glucosyltransferases (UGTs), which were up-regulated in all of our treatments. In plants, UGTs utilize UDP-glucose, UDP-galactose, and UDP-rhamnose as sugar donors and are involved in the modulation of plant architecture and the water stress response in *Arabidopsis* (Tognetti et al. 2010). The glucosylation of coniferyl aldehyde and sinapyl aldehyde may regulate both lignin biosynthesis and the metabolism of other phenylpropanoids, such as ferulic acid, 5-hydroxyferulic acid, sinapic acid and their derivatives (Lim et al. 2005). Thus, the presence of up-regulated UGTs in all of the groups is another interesting result that might help elucidate the chemical differences present in the flavonoid-supplemented trees.

Another notable finding regarding cell wall formation was the differential expression of genes involved in the metabolism of sucrose, starch, CHOs and minor sugars. Despite the down-regulation of sucrose synthase (Sus) and cellulose synthase (CesA), there were several other enzymes involved in this metabolic pathway that were up-regulated in the prolonged flavonoid treatments, even in the short-term treatments. Starting with galactinol synthase (GolS2), which was one of the most differentially expressed genes, all of the downstream genes in this pathway were differentially expressed (up-regulated), most of which were up-regulated after both the prolonged and short-term treatments.

The most notable of these genes was stachyose synthase (STS), which converts raafinose (a trisaccharide of galactose, fructose and glucose) into stachyose (a tetrasaccharide) by transferring a galactosyl moiety from galactinol, (Peterbauer et al. 2002). Raafinose synthase (RafS) was also differentially up-regulated. Additionally, sucrose phosphate synthase (SPS1F), which catalyses the conversion of UDP-glucose and D-fructose 6-phosphate into UDP and sucrose 6-phosphate (Chua et al. 2008), was also differentially up-regulated (Figure 4).

Starch synthase (SS) was up-regulated as well, as were enzymes involved in the degradation of starch into maltose (beta-amylase; BAM) and glucose (glycoside hydrolase; GH and phosphoglucan water dikinases; PWD). These results suggest a shift from sucrose and cellulose production to the synthesis of starch and minor sugars (galactinol, raffinose and stachyose).

Thus, to verify the possible effects of these transcriptional responses on cell wall formation and sugar accessibility, enzymatic hydrolysis was performed in wood samples from all of the experimental and control plants. The results revealed an increase in sugar contents (up to 250% in the NARSTOP group) and glucose yields of all of the flavonoid-supplemented *Eucalyptus* plants.

This may be the result of the plants producing more sugar or a result of the increased digestibility of lignin due to modifications of lignin structure and other cell wall components. The total values should increase exponentially with the use of pre-treatments and the extension of milling times (Chen and Dixon 2007; Inoue et al. 2008).

Galactinol, raffinose and stachyose have been described as being involved in freezing and stress tolerance (Saravitz et al. 1987; Taji et al. 2002). The levels of galactinol and raffinose are increased in the leaves of *Arabidopsis* plants over-expressing HsfA2, a heat shock factor, suggesting a possible role for these compounds in protection from oxidative damage (Nishizawa et al. 2008). Moreover, this factor may also constitute another link between the stress response and cell wall formation in flavonoid-supplemented *Eucalyptus* plants because all of our treatment groups showed more than 1 up-regulated HsfA2 gene.

Our results are also in agreement with those of Unda et al. (2012), who verified that in poplar trees, the over-expression of GoS and its product, galactinol, may serve as a molecular signal that initiates metabolic changes associated with combating stress, culminating in the formation of tension wood and increased glucose contents. Additionally, over-expression of raffinose synthase was found to result in increased biomass and total cellulose contents, while the total contents of lignin and xylose moieties were slightly reduced. Furthermore, the total amount of glucose was commensurately increased in the transgenic trees, by from ~1 to 4% (Unda et al 2012). Moreover, repression of the flavonoid pathway in *Arabidopsis* increases starch levels (Rubin et al. 2009), and a chalcone isomerase-deficient *Arabidopsis* mutant exhibits increased levels of starch and soluble sugars in its leaves.

Based on our results, flavonoid supplementation causes a stress response in *E. urograndis*, greatly affecting cell wall development, modifying lignification by affecting the expression of genes involved in the phenylpropanoid pathway and altering sugar metabolism in favor of starch and minor sugar synthesis, resulting in increased sugar accessibility and saccharification.

Conclusion

The effects of flavonoid supplementation on cell wall development in *Eucalyptus* plants are most likely due to a combination of transcriptional changes in several distinct pathways. The down-regulation of the phenylpropanoid pathway, combined with the up-regulation of ATOMT1 and CAD, results in a higher S/G ratio, which in turn, increases lignin solubility and facilitates access to cellulose and hemicellulose. Subsequently, as a result of the stress response, sugar metabolism is shifted towards starch and minor sugars, culminating in the increased sugar and saccharification levels identified due to hydrolysis.

Given the importance of *Eucalyptus* in several industrial sectors, there is great interest in expanding the use of these species as a resource for cellulose, paper and pulp production and as an alternative source of biomass for second-generation biochemicals. Our results contribute not only to our understanding of the molecular responses involved in wood formation but will also have a significant impact on the use of *Eucalyptus* as biomass. Finally, we expect our findings to guide future genetic manipulation and nutritional supplementation analyses of *Eucalyptus* trees aimed at achieving significant improvements in their productivity yields.

Acknowledgments: This project was funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ). We would especially like to thank International Paper of Brasil for their assistance.

References

- Altschul SF, Madden TL, Schäffer AA, et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic acids research* 25:3389–402.
- Baucher M, Halpin C, Petit-conil M (2003) Lignin : Genetic Engineering and Impact on Pulping. *Critical Reviews in Biochemistry* 305–350. doi: 10.1080/10409230390242443
- Berthet S, Demont-Caulet N, Pollet B, et al. (2011) Disruption of LACCASE4 and 17 results in tissue-specific alterations to lignification of *Arabidopsis thaliana* stems. *The Plant cell* 23:1124–37. doi: 10.1105/tpc.110.082792
- Besseau S, Hoffmann L, Geoffroy P, et al. (2007) Flavonoid accumulation in *Arabidopsis* repressed in lignin synthesis affects auxin transport and plant growth. *The Plant cell* 19:148–62. doi: 10.1105/tpc.106.044495
- Boudet A-M (2000) Lignins and lignification: Selected issues. *Plant Physiology and Biochemistry* 38:81–96. doi: 10.1016/S0981-9428(00)00166-2
- Boudet AM, Kajita S, Grima-Pettenati J, Goffner D (2003) Lignins and lignocellulosics: a better control of synthesis for new and improved uses. *Trends in plant science* 8:576–81.

- Bragatto J, Segato F, Cota J, et al. (2012) Insights on how the activity of an endoglucanase is affected by physical properties of insoluble celluloses. *The journal of physical chemistry B* 116:6128–36. doi: 10.1021/jp3021744
- Campbell MM, Sederoff RR (1996) Variation in Lignin Content and Composition (Mechanisms of Control and Implications for the Genetic Improvement of Plants). *Plant physiology* 110:3–13.
- Chen F, Dixon RA (2007) Lignin modification improves fermentable sugar yields for biofuel production. *Nature biotechnology* 25:759–61. doi: 10.1038/nbt1316
- Chen W, Yun M-S, Deng F, Yogo Y (2005) The rates of maize growth and lignin biosynthesis change after root-applied chalcone. *Weed Biology and Management* 5:118–122. doi: 10.1111/j.1445-6664.2005.00165.x
- Chen W-J, Yun M-S, Deng F, Yogo Y (2011) Chalcone suppresses lignin biosynthesis in illuminated soybean cells. *Weed Biology and Management* 11:49–56. doi: 10.1111/j.1445-6664.2011.00404.x
- Chen WEIJUN, Yun MINSOO, Deng FAN, et al. (2004) Effects of root-applied naringenin and chalcone on the growth of annual plants. *Weed Biology and Management* 238:235–238.
- Chua TK, Bujnicki JM, Tan T-C, et al. (2008) The structure of sucrose phosphate synthase from *Halothermothrix orenii* reveals its mechanism of action and binding mode. *The Plant cell* 20:1059–72. doi: 10.1105/tpc.107.051193
- Conesa A, Götz S, García-Gómez JM, et al. (2005) Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics (Oxford, England)* 21:3674–6. doi: 10.1093/bioinformatics/bti610
- Cook CM, Daudi A, Millar DJ, et al. (2012) Transcriptional changes related to secondary wall formation in xylem of transgenic lines of tobacco altered for lignin or xylan content which show improved saccharification. *Phytochemistry* 74:79–89. doi: 10.1016/j.phytochem.2011.10.009
- Deng F, Aoki M, Yogo Y (2004) Effect of naringenin on the growth and lignin biosynthesis of gramineous plants. *Weed Biology and Management* 4:49–55. doi: 10.1111/j.1445-6664.2003.00119.x
- Dixon RA, Lamb CJ, Masoud S, et al. (1996) Metabolic engineering: prospects for crop improvement through the genetic manipulation of phenylpropanoid biosynthesis and defense responses — a review. *Gene* 179:61–71. doi: 10.1016/S0378-1119(96)00327-7
- Dixon RA, Paiva NL (1995) Stress-Induced Phenylpropanoid Metabolism. *The Plant cell* 7:1085–1097. doi: 10.1105/tpc.7.7.1085
- Ferrer J-L, Austin MB, Stewart C, Noel JP (2008) Structure and function of enzymes involved in the biosynthesis of phenylpropanoids. *Plant physiology and biochemistry : PPB / Société française de physiologie végétale* 46:356–70. doi: 10.1016/j.plaphy.2007.12.009
- Foelkel C (2009) Papermaking Properties of Eucalyptus Trees , Woods , and Pulp Fibers Celso Foelkel. In: *Eucalyptus Online Book & newsletter*. <http://www.eucalyptus.com.br/eucaliptos/ENG14.pdf>.

- Foucart C, Jauneau A, Gion J-M, et al. (2009) Overexpression of EgROP1, a Eucalyptus vascular-expressed Rac-like small GTPase, affects secondary xylem formation in *Arabidopsis thaliana*. *The New phytologist* 183:1014–29. doi: 10.1111/j.1469-8137.2009.02910.x
- Fu D, Mazza G, Tamaki Y (2010) Lignin extraction from straw by ionic liquids and enzymatic hydrolysis of the cellulosic residues. *Journal of agricultural and food chemistry* 58:2915–22. doi: 10.1021/jf903616y
- Fukuda H (1996) XYLOGENESIS: INITIATION, PROGRESSION, AND CELL DEATH. *Annual review of plant physiology and plant molecular biology* 47:299–325. doi: 10.1146/annurev.arplant.47.1.299
- Goujon T, Sibout R, Pollet B, et al. (2003) A new *Arabidopsis thaliana* mutant deficient in the expression of O-methyltransferase impacts lignins and sinapoyl esters. *Plant molecular biology* 51:973–89.
- Grattapaglia D, Vaillancourt RE, Shepherd M, et al. (2012) Progress in Myrtaceae genetics and genomics: Eucalyptus as the pivotal genus. *Tree Genetics & Genomes* 8:463–508. doi: 10.1007/s11295-012-0491-x
- Guo D, Chen F, Inoue K, et al. (2001) Downregulation of caffeic acid 3-O-methyltransferase and caffeoyl CoA 3-O-methyltransferase in transgenic alfalfa. impacts on lignin structure and implications for the biosynthesis of G and S lignin. *The Plant cell* 13:73–88.
- Harbone J (1988) *The Flavonoids: Advances in Research Since 1980*. 624.
- Hardcastle TJ, Kelly KA (2010) baySeq: empirical Bayesian methods for identifying differential expression in sequence count data. *BMC bioinformatics* 11:422. doi: 10.1186/1471-2105-11-422
- Holton TA, Cornish EC (1995) Genetics and Biochemistry of Anthocyanin Biosynthesis. *The Plant cell* 7:1071–1083. doi: 10.1105/tpc.7.7.1071
- Hu WJ, Harding S a, Lung J, et al. (1999) Repression of lignin biosynthesis promotes cellulose accumulation and growth in transgenic trees. *Nature biotechnology* 17:808–12. doi: 10.1038/11758
- Inoue H, Yano S, Endo T, et al. (2008) Combining hot-compressed water and ball milling pretreatments to improve the efficiency of the enzymatic hydrolysis of eucalyptus. *Biotechnology for biofuels* 1:2. doi: 10.1186/1754-6834-1-2
- Kanehisa M (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research* 28:27–30. doi: 10.1093/nar/28.1.27
- Kirst ME, Scott J, Kirst M, et al. (2004) Coordinated Genetic Regulation of Growth and Lignin Revealed by Quantitative Trait Locus Analysis of cDNA Microarray Data in an Interspecific Backcross of Eucalyptus 1. *Society* 135:2368–2378. doi: 10.1104/pp.103.037960.At
- Koski LB, Gray MW, Lang BF, Burger G (2005) AutoFACT: an automatic functional annotation and classification tool. *BMC bioinformatics* 6:151. doi: 10.1186/1471-2105-6-151

- Lacombe E, Hawkins S, Van Doorselaere J, et al. (1997) Cinnamoyl CoA reductase, the first committed enzyme of the lignin branch biosynthetic pathway: cloning, expression and phylogenetic relationships. *The Plant journal : for cell and molecular biology* 11:429–41.
- Lepikson-Neto J, Alves A, Deckmann AC, et al. (2013) Flavonoid supplementation reduces extractives content and increases S/G ratio on *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid trees. *Bioresources* 1–11.
- Li J, Ou-Lee TM, Raba R, et al. (1993) Arabidopsis Flavonoid Mutants Are Hypersensitive to UV-B Irradiation. *The Plant cell* 5:171–179. doi: 10.1105/tpc.5.2.171
- Li R, Yu C, Li Y, et al. (2009) SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics (Oxford, England)* 25:1966–7. doi: 10.1093/bioinformatics/btp336
- Li X, Ximenes E, Kim Y, et al. (2010) Lignin monomer composition affects Arabidopsis cell-wall degradability after liquid hot water pretreatment. *Biotechnology for biofuels* 3:27. doi: 10.1186/1754-6834-3-27
- Lim E-K, Jackson RG, Bowles DJ (2005) Identification and characterisation of Arabidopsis glycosyltransferases capable of glucosylating coniferyl aldehyde and sinapyl aldehyde. *FEBS letters* 579:2802–6. doi: 10.1016/j.febslet.2005.04.016
- Mansfield SD, Kang K-Y, Chapple C (2012) Designed for deconstruction—poplar trees altered in cell wall lignification improve the efficacy of bioethanol production. *The New phytologist* 194:91–101. doi: 10.1111/j.1469-8137.2011.04031.x
- Miller G. (1959) Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. *Analytical Chemistry*. *Analytical chemistry* 31:426–428.
- Mizrachi E, Mansfield SD, Myburg A a. (2011) Cellulose factories: advancing bioenergy production from forest trees. *New Phytologist* no–no. doi: 10.1111/j.1469-8137.2011.03971.x
- Mortazavi A, Williams BA, McCue K, et al. (2008) Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nature methods* 5:621–8. doi: 10.1038/nmeth.1226
- Moustafa E (1967) Purification and properties of chalcone-flavanone isomerase from soya bean seed. *Phytochemistry* 6:625–632. doi: 10.1016/S0031-9422(00)86001-X
- Nishizawa A, Yabuta Y, Shigeoka S (2008) Galactinol and raffinose constitute a novel function to protect plants from oxidative damage. *Plant physiology* 147:1251–63. doi: 10.1104/pp.108.122465
- Paiva JAP, Prat E, Vautrin S, et al. (2011) Advancing Eucalyptus genomics: identification and sequencing of lignin biosynthesis genes from deep-coverage BAC libraries. *BMC genomics* 12:137. doi: 10.1186/1471-2164-12-137
- Paux E, Carocha V, Marques C, et al. (2005) Transcript profiling of Eucalyptus xylem genes during tension wood formation. *The New phytologist* 167:89–100. doi: 10.1111/j.1469-8137.2005.01396.x
- Peterbauer T, Mucha J, Mach L, Richter A (2002) Chain Elongation of raffinose in pea seeds. Isolation, characterization, and molecular cloning of multifunctional enzyme catalyzing the

- synthesis of stachyose and verbascose. *The Journal of biological chemistry* 277:194–200. doi: 10.1074/jbc.M109734200
- Phillips CM, Iavarone AT, Marletta MA (2011) Quantitative proteomic approach for cellulose degradation by *Neurospora crassa*. *Journal of proteome research* 10:4177–85. doi: 10.1021/pr200329b
- Plomion C, Leprovost G, Stokes A (2001) Wood Formation in Trees. *PLANT PHYSIOLOGY* 127:1513–1523. doi: 10.1104/pp.010816
- Le Provost G, Paiva J, Pot D, et al. (2003) Seasonal variation in transcript accumulation in wood-forming tissues of maritime pine (*Pinus pinaster* Ait.) with emphasis on a cell wall glycine-rich protein. *Planta* 217:820–30. doi: 10.1007/s00425-003-1051-2
- Pruesse E, Quast C, Knittel K, et al. (2007) SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic acids research* 35:7188–96. doi: 10.1093/nar/gkm864
- Quail MA, Kozarewa I, Smith F, et al. (2008) A large genome center's improvements to the Illumina sequencing system. *Nature methods* 5:1005–10. doi: 10.1038/nmeth.1270
- Rengel D, San Clemente H, Servant F, et al. (2009) A new genomic resource dedicated to wood formation in *Eucalyptus*. *BMC plant biology* 9:36. doi: 10.1186/1471-2229-9-36
- Rodrigues J, Meier D, Faix O, Pereira H (1999) Determination of tree to tree variation in syringyl/guaiacyl ratio of *Eucalyptus globulus* wood lignin by analytical pyrolysis. *Journal of Analytical and Applied Pyrolysis* 48:121–128. doi: 10.1016/S0165-2370(98)00134-X
- Rubin G, Tohge T, Matsuda F, et al. (2009) Members of the LBD family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in *Arabidopsis*. *The Plant cell* 21:3567–84. doi: 10.1105/tpc.109.067041
- Saravitz DM, Pharr DM, Carter TE (1987) Galactinol synthase activity and soluble sugars in developing seeds of four soybean genotypes. *Plant physiology* 83:185–9.
- Shamir R, Maron-Katz A, Tanay A, et al. (2005) EXPANDER--an integrative program suite for microarray data analysis. *BMC bioinformatics* 6:232. doi: 10.1186/1471-2105-6-232
- Suzek BE, Huang H, McGarvey P, et al. (2007) UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics (Oxford, England)* 23:1282–8. doi: 10.1093/bioinformatics/btm098
- Taji T, Ohsumi C, Iuchi S, et al. (2002) Important roles of drought- and cold-inducible genes for galactinol synthase in stress tolerance in *Arabidopsis thaliana*. *The Plant journal : for cell and molecular biology* 29:417–26.
- Tognetti VB, Van Aken O, Morreel K, et al. (2010) Perturbation of indole-3-butyric acid homeostasis by the UDP-glucosyltransferase UGT74E2 modulates *Arabidopsis* architecture and water stress tolerance. *The Plant cell* 22:2660–79. doi: 10.1105/tpc.109.071316
- Trapnell C, Pachter L, Salzberg SL (2009) TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics (Oxford, England)* 25:1105–11. doi: 10.1093/bioinformatics/btp120

- Unda F (2012) EVALUATING THE ROLE OF THE RAFFINOSE FAMILY OF IN HYBRID POPLAR (*Populus alba* × *grandidentata*). The University of British Columbia
- Unda F, Canam T, Preston L, Mansfield SD (2012) Isolation and characterization of galactinol synthases from hybrid poplar. *Journal of experimental botany* 63:2059–69. doi: 10.1093/jxb/err411
- Vanholme R, Demedts B, Morreel K, et al. (2010) Lignin biosynthesis and structure. *Plant physiology* 153:895–905. doi: 10.1104/pp.110.155119
- Vogt T (2010) Phenylpropanoid biosynthesis. *Molecular plant* 3:2–20. doi: 10.1093/mp/ssp106
- Voo KS, Whetten RW, O'Malley DM, Sederoff RR (1995) 4-coumarate:coenzyme a ligase from loblolly pine xylem. Isolation, characterization, and complementary DNA cloning. *Plant physiology* 108:85–97.
- Zeng Y, Yang T (2002) RNA isolation from highly viscous samples rich in polyphenols and polysaccharides. *Plant Molecular Biology Reporter* 20:417–417. doi: 10.1007/BF02772130

FIGURES AND TABLES

Table 1. Gene ID, FPKM values and annotation of the 36 genes that found to bedifferentially expressed in all tested conditions.

Gene ID	Annotation	FPKM				
		CT	CH	NAR	CHSTOP	NARSTOP
Eucgr.F04479.1	HSP20	0.12	35.55	40.44	23.84	57.92
Eucgr.K02389.1	unknown	0.04	13.69	10.18	9.89	26.99
Eucgr.K02399.1	unknown	0.08	18.09	18.67	19.48	53.82
Eucgr.G01188.2	EGY3	2.78	45.78	41.56	33.40	76.35
Eucgr.J01979.1	HSP18.2	0.34	17.61	19.94	14.63	27.21
Eucgr.K02410.1	unknown	0.13	14.27	10.66	12.52	29.61
Eucgr.J01980.1	HSP18.2	0.02	12.20	11.59	9.15	26.39
Eucgr.L02233.1	no hit	1.29	41.47	38.15	29.24	118.40
Eucgr.F02898.1	HSP20	2.76	525.99	343.17	280.68	342.59
Eucgr.J01985.1	HSP18.2	0.31	22.25	16.62	13.21	33.33
Eucgr.K03553.1	STS	0.04	4.11	4.59	2.83	6.87
Eucgr.L03261.1	HSP18.2	1.44	47.41	27.81	19.98	75.67
Eucgr.C03449.1	HSFA2	0.29	14.55	12.98	14.36	18.58
Eucgr.C00684.1	HSP17.6II	2.00	341.30	299.30	243.65	272.90
Eucgr.K02384.1	unknown	0.07	14.56	10.59	10.60	27.23
Eucgr.J01969.1	HSP20	4.89	192.23	134.53	103.09	310.44
Eucgr.K03472.1	ARATH	0.07	109.57	84.15	62.04	20.94
Eucgr.H04513.1	HSP70	0.23	15.72	18.79	11.56	21.62
Eucgr.A00595.1	PEBP	0.10	81.17	76.66	57.33	98.83
Eucgr.E02421.1	unknown	0.19	260.04	220.12	119.63	51.05
Eucgr.H04692.1	HSP21	2.97	83.31	59.57	43.11	313.22
Eucgr.G02440.1	UGT73B2	0.00	5.46	5.80	3.33	4.06
Eucgr.G02259.1	UGT73B3	0.00	2.73	2.14	1.18	3.10
Eucgr.J01959.1	HSP18.2	3.19	142.90	89.76	58.22	148.21
Eucgr.K00295.1	HSP90.1	2.11	46.13	38.25	35.52	62.69
Eucgr.A01833.1	AAC3	0.13	32.00	24.43	15.08	10.47
Eucgr.C03071.1	HSP17.6II	3.64	517.08	509.51	451.66	324.54
Eucgr.B03843.1	no hit	1.45	93.17	63.39	67.61	20.77
Eucgr.C03320.1	DUF1677	0.38	24.39	18.22	14.01	9.78
Eucgr.B00176.2	PIMT2	3.86	153.25	109.01	82.97	57.39
Eucgr.J02588.1	no hit	3.20	225.90	182.76	194.01	130.04
Eucgr.C00690.1	HSP17.6II	2.48	563.68	498.40	514.67	458.94
Eucgr.K00237.1	PEBP	0.04	115.83	64.41	61.47	11.49
Eucgr.F03196.1	GSTU25	1.43	292.73	240.63	168.71	38.29
Eucgr.I02136.1	HSP20	1.68	226.73	147.43	90.02	259.35
Eucgr.H04427.1	MEE32	49.92	0.52	0.85	1.32	13.93

A total of 36 genes were differentially expressed in all four conditions. FPKM -fragments per kilobase of exon per million fragments mapped. All results are controlled for the FDR – false discovery rate ≤ 0.01 . CT – control; CH – prolonged naringenin-chalcone supp; NAR – prolonged naringenin supp; CHSTOP- short-term naringenin-chalcone supp; NARSTOP – short-term naringenin supp. *HSP20*– *HSP20-like chaperone superfamily protein*; *unknown*– *unknown protein*; *EGY3*–*ethylene-dependent gravitropism-deficient and yellow-green-like 3*; *HSP18.2*–*heat shock protein 18.2*; *HSP20*- *HSP20-like chaperones superfamily protein*; *STS*- *stachyose synthase*; *HSFA2*- *heat shock transcription factor A2*; *HSP17.6II* –*17.6 kDa class II heat shock protein*; *ARATH* –*Adenine nucleotide alpha hydrolases-like superfamily protein*; *HSP70* –*BIP1heat shock protein 70 family protein*; *PEBP* – *phosphatidylethanolamine-binding protein family protein*; *HSP21* –*heat shock protein 21*; *UGT73B2* –*UDP-glucosyltransferase 73B2*; *UGT73B3UDP* – *glucosyl transferase 73B3*; *HSP90.1* –*heat shock protein 90.1*; *AAC3* – *ADP/ATP carrier 3*; *DUF1677* –*protein of unknown function*; *PIMT2* –*protein-l-isoaspartate methyltransferase 2*;

Table 2. Differentially expressed phenylpropanoid-related genes

Gene ID	Annotation	FPKM				
		CT	CH	NAR	CHSTOP	NARSTOP
Eucgr.C00859.1	U91A1	0.00	1.60*	0.83	0.28	0.00
Eucgr.K00903.1	AAT	0.38	3.91	4.61*	3.45	1.26
Eucgr.K00901.1	AAT	0.83	0.03*	0.98	1.18	1.51
Eucgr.E01250.1	PRR1	38.77	3.72*	4.68*	7.66	21.15
Eucgr.B03781.1	AA	24.04	0.04*	0.86	1.39	9.77
Eucgr.D02454.1	DFR	0.05	3.10*	2.73*	1.32	0.48
Eucgr.G02325.1	DFR	2.51	28.24*	20.16*	21.06	9.55
Eucgr.F04163.1	LAC14	6.22	0.15	0.32*	0.57	1.82
Eucgr.F02646.1	LAC14	1.99	0.00*	0.02*	0.06*	0.44
Eucgr.F04162.1	LAC14	1.95	0.09	0.04*	0.06	0.72
Eucgr.H04937.1	LAC14	13.67	0.02*	0.15*	0.21	4.66
Eucgr.F04160.1	LAC14	17.17	0.04*	0.25*	0.26*	3.08
Eucgr.F02674.1	LAC14	7.13	0.28	0.27*	0.42	2.90
Eucgr.H04936.1	LAC14	8.51	0.04*	0.03*	0.08	2.50
Eucgr.B02796.1	LAC4	12.04	0.28*	1.95	3.31	5.69
Eucgr.K00957.1	ATOMT1	1.36	17.54	17.19*	10.55	15.16
Eucgr.A01877.1	OMT-like	0.00	0.31	0.79	1.75*	0.22
Eucgr.J00363.1	HCT	88.68	3.76*	16.43	29.57	53.35
Eucgr.B00137.1	4CL	12.31	1.83	2.50*	4.21	6.24
Eucgr.E00270.1	CCR	30.29	3.21	3.29*	4.03	8.00
Eucgr.G01350.2	CAD5	23.73	146.29*	126.30*	144.55	62.96
Eucgr.E01110.2	CAD1	4.34	59.21	49.17*	51.45	27.12

U91A1 –UDP-Glycosyltransferase superfamily protein; *AAT* –HXXXD-type acyl-transferase family protein; *DFR* –Dihydroflavonol-4-reductase; *LAC14* –laccase 14; *LAC4* –laccase 4; *ATOMT1* –O-methyltransferase 1; *OMT-like* –O-methyltransferase family protein; *HCT* –hydroxycinnamoyl-CoA shikimate transferase; *4CL* –4 coumarate CoA ligase; *CCR*–cinnamoyl-CoA reductase; *CAD*–cinnamyl alcohol dehydrogenase.*Denotes differential expression

Table 3. Differentially expressed secondary cell wall genes

Gene ID	Annotation	FPKM				
		CT	CH	NAR	CHSTOP	NARSTOP
Eucgr.C03199.1	SUS4	1,532.75	66.91*	100.38*	132.04	213.15
Eucgr.C01715.1	SPS1F	3.32	63.28*	54.72	35.80*	28.00
Eucgr.F00464.1	SUT4	28.18	85.70	85.04	82.49*	36.89
Eucgr.D01765.2	CSLG3	0.07	3.26	6.28*	7.07*	1.24
Eucgr.F04010.1	CSLC05	7.51	0.11*	0.35	0.47*	3.06
Eucgr.J00420.1	CSLA2	41.08	2.63*	5.07*	5.87*	25.59
Eucgr.E00226.1	CSLD3	10.13	0.53	0.78	0.68	2.11
Eucgr.E00821.1	CSLG2	3.07	0.38*	0.35	0.78*	3.03
Eucgr.J02497.1	AMR1	1.00	5.25	6.14	6.11*	2.68
Eucgr.J02407.1	MUR1	74.28	18.89	17.82	19.95*	38.28
Eucgr.B03204.1	MUR2	13.62	55.51	54.92	47.91*	32.70
Eucgr.J01663.1	XTH5	97.41	1.21*	1.87*	3.31*	75.71
Eucgr.B03348.1	XTH33	26.89	0.21	0.15*	0.07*	9.47
Eucgr.K00883.2	XTH9	607.60	16.06*	29.74	37.61*	288.85
Eucgr.C00184.1	XTH23	45.26	0.45	0.72*	0.22*	37.20
Eucgr.H02634.1	XTH16	386.73	21.78*	47.14	72.69	396.09
Eucgr.D01294.1	XTH8	10.82	1.01	1.54	1.66*	7.36
Eucgr.J00827.1	GSL12	0.04	0.90*	1.56*	0.98*	0.76*
Eucgr.A02002.1	GSL7	0.20	1.24	2.13	1.38*	0.79
Eucgr.A02008.1	GSL7	0.16	1.07	1.89	1.66*	0.83
Eucgr.K02988.2	GH	16.20	90.51*	69.09*	53.94	47.27*
Eucgr.H00494.1	PWD	6.13	20.24	26.24*	27.22	7.21
Eucgr.H03767.1	BAM9	39.25	225.21	235.09*	217.12	115.07
Eucgr.E00460.1	TPS	0.08	5.75*	6.21*	4.26	0.34
Eucgr.K00387.1	SS	9.88	66.76*	54.46*	39.19	19.93
Eucgr.C04266.1	RafS	26.09	1,317.97*	1007.17	544.49	193.24
Eucgr.K03553.1	STS	0.04	4.11*	4.59*	2.83*	6.87*
Eucgr.H00997.1	STS	0.81	37.45*	29.83*	17.09*	7.06*
Eucgr.K03563.1	GoSL1	0.23	4.48	11.12*	12.00	8.48*
Eucgr.L00249.1	GoSL2	0.34	280.16*	135.36*	52.37*	1.25
Eucgr.L00243.1	GoSL2	0.02	29.83*	19.77*	9.75*	1.37*
Eucgr.L00251.1	GoSL2	0.21	325.56*	149.21*	61.46*	3.38
Eucgr.L03245.1	GoSL2	0.07	190.81*	124.71*	44.86*	2.79*
Eucgr.L00240.1	GoSL2	0.02	34.50*	22.77*	10.27*	0.66
Eucgr.L00248.1	GoSL2	0.17	162.07*	86.93*	32.28*	0.69
Eucgr.L03244.1	GoSL2	0.12	279.83*	137.40*	63.66*	2.99*
Eucgr.L00235.1	GosL2	0.04	73.19*	38.88*	15.60*	0.03
Eucgr.L00245.1	GosL2	1.83	287.87*	164.66	80.81*	5.28
Eucgr.F01661.1	Invertase	0.15	2.82*	2.16	2.39*	1.33
Eucgr.J00457.2	Invertase	5.69	47.20*	42.83	33.08*	18.78
Eucgr.G01751.1	Invertase	4.83	0.23*	0.42	1.17*	2.23
Eucgr.A02888.1	Invertase	7.36	0.04*	0.08*	0.08*	1.55

Sus4 –sucrose synthase 4; *SPS1F* –sucrose phosphate synthase 1F; *SUT4* –sucrose transporter 4; *CSLG3* –cellulose synthase like G3; *CSLD3* –cellulose synthase-like D3; *CSLC05* –Cellulose-synthase-like C5; *CSLA2* –cellulose synthase-like A02; *CSLG2* –cellulose synthase like G2; *CSLG3* –cellulose synthase like G3; *CESA3* –cellulose synthase family protein; *AMR1* –ascorbic acidmannose pathway regulator  *GDP-mannose 4,6 dehydratase 2*; *MUR2* fucosyltransferase 1; *XTH5* xyloglucan endotransglucosylase/hydrolase 5; *XTH33* xyloglucosyl transferase 33; *XTH9* xyloglucan endotransglucosylase/hydrolase 9; *XTH23* xyloglucan endotransglucosylase 6; *XTH16* xyloglucan endotransglucosylase/hydrolase 16; *XTH8* xyloglucan endotransglucosylase/hydrolase 8; *GSL12* glucan synthase-like 12; *GSL7* glucan synthase-like 7; *GH* glycoside hydrolase; *PWD* phosphoglucan water dikinases; *BAM9* beta-amylase 9 *TPS* trehalose-6-phosphate synthase; *SS* starch synthase; *Rafs* rafiinose synthase; *STS* stachyose synthase; *GoSL1* galactinol synthase 1; *GoSL2* galactinol synthase 2.

*Denotes differential expression

Table 4. Differentially expressed stress-related genes

Gene ID	Annotation	FPKM				
		CT	CH	NAR	CHSTOP	NARSTOP
Eucgr.H05081.4	ALDH3I1	4.84	0.93*	1.11*	1.60	5.34
Eucgr.C00112.1	CIA2	0.70	12.29*	14.05*	17.42*	3.57
Eucgr.K01387.2	COL9	3.33	26.78*	22.63*	18.02*	2.56
Eucgr.C03449.1	HSFA2	0.29	14.55*	12.98*	14.36*	18.58*
Eucgr.C03456.1	HSFA2	0.09	2.67*	3.64*	2.25*	1.45
Eucgr.C00873.1	HSFA2	1.92	24.38*	18.82*	14.75*	7.94
Eucgr.C03434.1	HSFA2	0.31	6.75*	6.22*	6.19*	2.03
Eucgr.C01043.1	HSFC1	2.92	122.26*	116.87*	96.98*	13.96
Eucgr.J01981.1	HSP18.2	2.00	34.43	50.34	69.86*	105.48*
Eucgr.J01980.1	HSP18.2	0.02	12.20*	11.59*	9.15*	26.39*
Eucgr.J01959.1	HSP18.2	3.19	142.90*	89.76*	58.22*	148.21*
Eucgr.J01958.1	HSP18.2	2.68	115.32*	90.34*	65.56	100.03
Eucgr.J01979.1	HSP18.2	0.34	17.61*	19.94*	14.63*	27.21*
Eucgr.J01958.1	HSP18.2	2.68	115.32*	90.34*	65.56	100.03
Eucgr.J01979.1	HSP18.2	0.34	17.61*	19.94*	14.63*	27.21*
Eucgr.J01977.1	HSP18.2	0.37	9.43*	8.59*	7.08	19.26*
Eucgr.J01964.1	HSP18.2	13.65	145.21*	119.05*	111.04	251.88*
Eucgr.J01985.1	HSP18.2	0.31	22.25*	16.62*	13.21*	33.33*
Eucgr.J01982.1	HSP18.2	0.21	8.65*	6.06*	4.24	13.03*
Eucgr.F04479.1	HSP20-like	0.12	35.55*	40.44*	23.84*	57.92*
Eucgr.I02136.1	HSP20-like	1.68	226.73*	147.43*	90.02*	259.35*
Eucgr.J01969.1	HSP20-like	4.89	192.23*	134.53*	103.09*	310.44*
Eucgr.G00061.1	HSP20-like	9.45	834.94*	858.21*	726.74*	814.87
Eucgr.E00433.1	HSP20-like	3.35	295.94	205.55*	185.03	165.89
Eucgr.F02898.1	HSP20-like	2.76	525.99*	343.17*	280.68*	342.59*
Eucgr.A01416.1	HSP21	0.08	12.40*	7.60*	4.34*	0.29
Eucgr.H04692.1	HSP21	2.97	83.31*	59.57*	43.11*	313.22*
Eucgr.J03127.1	Hsp70b	8.27	1576.75	1347.55*	881.92	925.42
Eucgr.H03518.1	HSP70T-2	6.06	282.04*	218.12*	152.91	249.47
Eucgr.K00295.1	HSP90-1	2.11	46.13*	38.25*	35.52*	62.69*
Eucgr.F03704.1	MSL6	1.45	27.74*	21.67*	11.53*	10.03
Eucgr.H02896.1	MYB305	0.07	6.57*	8.20*	6.34*	1.14
Eucgr.C00618.1	Oleosin	0.50	38.75*	24.66*	18.36*	2.85
Eucgr.F01003.1	PAT1	2.30	43.54*	47.40*	43.89*	6.02
Eucgr.K00237.1	PEBP	0.04	115.83*	64.41*	61.47*	11.49*
Eucgr.B00176.2	PIMT2	3.86	153.25*	109.01*	82.97*	57.39*
Eucgr.G01510.1	RLK	1.71	10.40*	11.17*	11.20*	6.80
Eucgr.F01854.1	TRX1	4.16	598.03*	295.93*	191.61*	25.01
Eucgr.G02440.1	UGT73B2	0.00	5.46*	5.80*	3.33*	4.06*
Eucgr.L03261.1	UGT73B3	1.44	47.41*	27.81*	19.98*	75.67*
Eucgr.G02259.1	UGT73B3	0.00	2.73*	2.14*	1.18*	3.10*
Eucgr.I00409.1	UGT73B3	0.06	3.27*	4.18*	2.91*	0.60
Eucgr.B02291.1	UGT76E11	10.86	52.79*	42.69*	39.45	45.69

Eucgr.K01389.2	XERICO	19.81	1,348.64*	796.35*	569.70*	120.14
----------------	--------	-------	-----------	---------	---------	--------

ALDH3I1 –aldehyde dehydrogenase 3I1; CIA2 –chloroplast import apparatus 2; COL9 –CONSTANS-like 9; HSFA2 –heat shock transcription factor A2; HSFC1 –heat shock transcription factor C1; HSP18.2 –heat shock protein 18.2; HSP20 like chaperones superfamily protein; HSP21 –heat shock protein 21; Hsp70b –heat shock protein 70B; HSP70T-2 –heat-shock protein 70T-2; HSP90.1 –heat shock protein 90.1; MSL6 – mechanosensitive channel of small conductance-like 6; MYB305 –myb domain protein 305; Oleosin family protein; PAT1 –GRAS family transcription factor; PEBP –phosphatidylethanolamine-binding protein family protein; RLK –receptor lectin kinase; TRX1 –thioredoxin H-type 1; UGT73B2 –UDP-glucosyltransferase 73B2; UGT73B3 –UDP-glucosyl transferase 73B3; UGT76E11 –UDP-glucosyl transferase 76E11; XERICO –RING/U-box superfamily protein

Table 5. Total sugar and glucose values

	n	Reduced sugars (mg/ml)	Reduced sugar Yield %	Glucose (mg/ml)	Saccharification %
CT	7	1.17 (0.67)	5.69 (3.23)	0.39 (0.23)	1.76 (1.13)
CH	4	1.8 (0.33)	9.06 (1.83)	0.56 (0.07)	2.59 (0.37)
NAR	3	2.54 (0.005)**	12.72 (0.66) **	0.87 (0.10) **	4.22 (0.5) **
CHSTOP	3	2.32 (0.24)*	11.81 (1.08)*	0.85 (0.29)*	4.1 (1.46)*
NARSTOP	3	3 (0.85) **	14.59 (4.34) **	0.99 (0.09) **	4.83 (0.46) **

Mean values and standard deviations (parentheses) for total sugar and glucose levels. n –number of biological replicates; CT – control; CH – prolonged naringenin-chalcone supp; N – prolonged naringenin supp; CHSTOP –short-termnaringenin-chalcone supp; NSTOP – short-termnaringenin supp. * p-value < 0.05; ** p-value < 0.01




Figure 1. A) Heat map clustering of differentially expressed transcripts and comparison of the estimated log2 fold change correlations between each group subjected to differential expression analyses. B) Venn diagram of differentially expressed genes. CT – control; CH – prolonged naringenin-chalcone supp; NAR – prolonged naringenin supp; CHSTOP- short-term naringenin-chalcone supp; NARSTOP – short-termnaringenin supp

Enriched GO Categories





Figure 2. Common GO categories that were enriched (p -values ≤ 0.05) between treatments. CH – prolonged naringenin-chalcone supp; NAR – prolonged naringenin supp; CHSTOP- short term naringenin-chalcone supp; NARSTOP – short-term naringenin supplementation.




Figure 4. The effects of flavonoid supplementation on the expression of secondary cell wall-related genes. *Sus* – sucrose synthase; *SPS* –sucrose phosphate synthase; *CesA* –cellulose synthase;; *GH* –glycoside hydrolase; *PWD* –phosphoglucan water dikinases; *BAM* –beta-amylase; *TPS* –trehalose-6-phosphate synthase; *SS* –starch synthase; *RafS* –raafinose synthase; *STS* –stachyose synthase; *GoSL* –galactinol synthase

5. Discussão

Para avaliar a influência da suplementação por flavonóides na formação da madeira em Eucalipto, mudas de um híbrido comercial de *E. urograndis*, foram suplementadas com os flavonóides narigenin e narigenin-chalcona, durante um ou cinco meses e após a suplementação foram analisadas quanto a composição química da sua parede celular, acessibilidade de enzimas hidrolíticas a seus açúcares e expressão gênica. Todos os grupos tratados por flavonóides apresentaram diferenças fenotípicas em relação aos controles (não tratados), e os fenótipos dos grupos foram bastante semelhantes entre si, independentemente do tipo de flavonóide utilizado ou tempo de suplementação.

Após as análises químicas e de expressão gênica, fica claro que a suplementação por flavonóides exerce grande influência na formação da madeira em Eucalipto. Quimicamente, verifica-se uma redução bastante significativa na quantidade de extrativos, o que constitui um importante resultado uma vez que afeta a cor da pasta de papel. Apesar dos números totais do teor de lignina permanecerem semelhantes entre os grupos suplementados e os grupos controle, sua composição monomérica é significativamente alterada após suplementação pelos flavonóides, com favorecimento da via de produção de lignina siringílica (S) em detrimento da guaiacilica (G) e cumarilica (H). Este resultado também é bastante relevante, uma vez que estes monômeros, por fazerem menos ligações intermôméricas, são mais solúveis e, portanto, mais fáceis de serem removidos.

Ambos os resultados são de grande importância tanto para a indústria de papel e celulose, como para a utilização do Eucalipto como fonte de material lignocelulósico para obtenção de açúcares fermentáveis para a produção de etanol de segunda geração. No caso de seu uso para produção de papel e celulose, o aumento da relação S/G pode significar um menor consumo de materiais químicos para a remoção da lignina da celulose usada para a fabricação de papel, sem comprometer o balanço energético da fábrica, uma vez que a lignina também é utilizada como fonte de energia para estas. Já para a produção de etanol de segunda geração, o fato da lignina estar mais “frouxa” significa maior acessibilidade das enzimas hidrolíticas à celulose e sua conseqüente quebra à açúcares fermentáveis. Fato que se confirmou nos resultados da hidrólise enzimática do material, com aumento tanto dos açúcares redutores totais quanto da quantidade de glicose encontrada nas mudas suplementadas.

Para melhor entender o que estaria causando estas modificações, analisamos a expressão gênica das mudas suplementadas através do seqüenciamento do seu transcriptoma, e verificamos a expressão diferencial de diversos genes e vias

metabólicas chaves para o desenvolvimento vegetal. Dentre os resultados mais relevantes, destacam-se a grande quantidade de genes relacionados a vias metabólicas de resposta ao estresse e às condições ambientais, todas induzidas nos grupos de suplementação por flavonóides em relação aos controles. Também foram diferencialmente expressos diversos genes relacionados à formação da parede celular e da via de fenilpropanóides, a qual inclui as rotas de síntese de lignina.

Olhando especificamente a via de formação de lignina, varios genes estão diferencialmente expressos entre os tratamentos. 4CL, HCT, diversas lacases e CCR estão reprimidos, enquanto duas metiltransferases e dois genes da família CAD, estão induzidos nos grupos de suplementação por flavonóides. Este resultado é muito interessante por fornecer uma explicação, ainda que parcial, para o aumento da razão S/G nas plantas suplementadas. Consideramos que este resultado constitui uma explicação parcial porque, apesar de observamos valores de expressão (FPKM) semelhantes entre os grupos de suplementação, estes resultados só foram considerados estatisticamente significativos nos grupos que tiveram a suplementação prolongada (NAR e CH). Dessa forma, hipotetizamos que deve existir um controle mais complexo relacionado à síntese dos monômeros de lignina que poderia ser explicado pela presença de muitos genes sem identidade e função definida (*no hits; unknown function*) como diferencialmente expressos. Outra possibilidade é a contribuição de outras vias metabólicas.

Entre as vias que podem estar afetando a composição da madeira das plantas suplementadas em relação ao grupos controle, nos chamou a atenção as vias relacionadas com resposta a estresse. Diversos genes relacionados a estresse estão induzidos em todos os grupos de suplementação, independentemente do tipo ou duração do tratamento. Estresse e lignificação estão intimamente ligados, sendo esta uma resposta natural dos organismos vegetais. Os resultados deste trabalho indicam que existe uma ligação entre suplementação por flavonóides, estresse e consequentemente, modificação da composição monomérica da lignina e da parede celular.

Analisando os genes diferencialmente expressos relacionados com a formação da parede celular, verificamos que existe um desvio na síntese de açúcares e polissacarídeos. Genes relacionados com a síntese da celulose e sacarose estão reprimidos, enquanto que estão induzidos, genes envolvidos com a síntese de amido (e de sua degradação em maltose e glicose) e açúcares menores, estaquiose e rafinose. Genes desta via participam da resposta a estresse abióticos, seca e temperaturas baixas, mas também estão envolvidos com a formação de madeira de tensão, aumento de biomassa e do conteúdo celulósico. Além disso em plantas de

arabidopsis super-expressando o gene HsfA2 (Heat-shock factor A2), que também está induzido em todos os tratamentos, foi verificado um aumento dos níveis de rafinose.

Desta forma, os altos valores de FPKM encontrados em todos os grupos suplementados com flavonóides para os genes desta via, ligam à resposta ao estresse, que pode estar sendo causado pela simples adição destes flavonóides, aos resultados histológicos que demonstram uma aparente maior presença de madeira de tensão, aos resultados de expressão gênica e da hidrólise enzimática.

As diferenças fenotípicas encontradas são, provavelmente, resultado não de um desses fatores, mas sim de uma combinação dos mesmos. A suplementação por flavonóides seria o agente causador de estresse no organismo, e a resposta do vegetal envolveria a expressão de diversos genes e vias metabólicas, resultando numa redução de extrativos, modificação da composição monomérica da lignina, em uma possível maior produção e desvio da síntese de açúcares e polissacarídeos na direção da formação de amido, estaquiose e rafinose, e a maior acessibilidade a estes.

Assim, a presente tese de doutorado abre novas perspectivas de pesquisa na área florestal e no melhoramento vegetal do Eucalipto. Testes mais amplos em campo são necessários para determinar a extensão do efeito da suplementação nas árvores adultas e para determinar a otimização dos flavonóides como composto nutricional, como por exemplo a concentração mais adequada, o tempo necessário de aplicação, idade correta das plantas para início e fim da aplicação, dentre outros. Além de serem necessários ensaios fermentativos e uma maior caracterização dos açúcares obtidos na hidrólise para verificar o potencial da suplementação por flavonóides na produção de etanol.

Entretanto, a simples perspectiva de alcançar mudanças fenotípicas benéficas e significativas pela adição de um composto barato ao meio nutricional costumeiramente utilizado é muito favorável, especialmente em se tratando do melhoramento de um clone comercial, o qual já passou por inúmeras etapas de melhoramento clássico e tem sido amplamente selecionado devido a suas características ligadas a produtividade.

6. Conclusões

A suplementação de mudas de Eucalipto com flavonóides:

- Reduz a quantidade de extrativos em até 41%;
- Não altera as quantidades totais de lignina;
- Altera a composição monomérica da lignina, aumentando em até 8% a razão S/G e diminuindo a razão H/G em até 15%;
- Promove alterações transcricionais, modificando a expressão de genes presentes em vias de formação da parede celular e desenvolvimento vegetal, desviando a via para a formação e degradação de amido, além de carboidratos e açúcares menores e de resposta ao estresse e ao meio ambiente;
- Melhora a acessibilidade de enzimas hidrolíticas aos açúcares da parede celular;
- Aumenta a quantidade total de açúcares e de glicose obtidos através da hidrólise enzimática;
- Os flavonóides narigenin-chalcona e narigenin tem potencial para serem utilizados pela indústria de papel, celulose e energia para aumentar o rendimento de suas plantações.

7. Referências

- Andersen, Øyvind M. II. Markham, Kenneth R. (2006) Flavonoids : chemistry, biochemistry, and applications. Taylor & Francis.
- Altschul SF, Madden TL, Schäffer AA, et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic acids research* 25:3389–402.
- Alves A, Schwanninger M, Pereira H, Rodrigues J (2006) Analytical pyrolysis as a direct method to determine the lignin content in wood. *Journal of Analytical and Applied Pyrolysis* 76:209–213. doi: 10.1016/j.jaap.2005.11.004
- Alves A, Simões R, Stackpole D, et al. (2011) Determination of the syringyl/guaiacyl (S/G) ratio of Eucalyptus globulus wood lignin by NIR-based PLS-R models using analytical pyrolysis as the reference method. *Journal of Near Infrared Spectroscopy* 19:343. doi: 10.1255/jnirs.946
- Alves AMM, Simões RFS, Santos CA, Potts BM (2012) Determination of Eucalyptus globulus wood extractives content by near infrared-based partial least squares regression models : comparison between extraction procedures. *Journal of Near Infrared Spectroscopy* 1–11. doi: 10.1255/jnirs.xxx
- Baucher M, Halpin C, Petit-conil M (2003) Lignin : Genetic Engineering and Impact on Pulp. *Critical Reviews in Biochemistry* 305–350. doi: 10.1080/10409230390242443
- Berthet S, Demont-Caulet N, Pollet B, et al. (2011) Disruption of LACCASE4 and 17 results in tissue-specific alterations to lignification of Arabidopsis thaliana stems. *The Plant cell* 23:1124–37. doi: 10.1105/tpc.110.082792
- Besseau S, Hoffmann L, Geoffroy P, et al. (2007) Flavonoid accumulation in Arabidopsis repressed in lignin synthesis affects auxin transport and plant growth. *The Plant cell* 19:148–62. doi: 10.1105/tpc.106.044495
- Boerjan W, Ralph J, Baucher M (2003) Lignin biosynthesis. *Annual review of plant biology* 54:519–46. doi: 10.1146/annurev.arplant.54.031902.134938
- Boland, Brooker MIH, Chippendale GM, et al. (2006) *Forest Trees of Australia*. 768.
- Bolwell GP, Mavandad M, Millar DJ, et al. (1988) Inhibition of mRNA levels and activities by trans-cinnamic acid in elicitor-induced bean cells. *Phytochemistry* 27:2109–2117. doi: 10.1016/0031-9422(88)80106-7
- Boudet A-M (2000) Lignins and lignification: Selected issues. *Plant Physiology and Biochemistry* 38:81–96. doi: 10.1016/S0981-9428(00)00166-2
- Boudet AM, Kajita S, Grima-Pettenati J, Goffner D (2003) Lignins and lignocellulosics: a better control of synthesis for new and improved uses. *Trends in plant science* 8:576–81.

- Bragatto J, Segato F, Cota J, et al. (2012) Insights on how the activity of an endoglucanase is affected by physical properties of insoluble celluloses. *The journal of physical chemistry B* 116:6128–36. doi: 10.1021/jp3021744
- Campbell MM, Sederoff RR (1996) Variation in Lignin Content and Composition (Mechanisms of Control and Implications for the Genetic Improvement of Plants). *Plant physiology* 110:3–13.
- Carroll A, Somerville C (2009) Cellulosic biofuels. *Annual review of plant biology* 60:165–82. doi: 10.1146/annurev.arplant.043008.092125
- Caño-Delgado A, Penfield S, Smith C, et al. (2003) Reduced cellulose synthesis invokes lignification and defense responses in *Arabidopsis thaliana*. *The Plant journal : for cell and molecular biology* 34:351–62.
- Chaffey N, Cholewa E, Regan S, Sundberg B (2002) Secondary xylem development in *Arabidopsis*: a model for wood formation. *Physiologia plantarum* 114:594–600.
- Chen F, Dixon RA (2007) Lignin modification improves fermentable sugar yields for biofuel production. *Nature biotechnology* 25:759–61. doi: 10.1038/nbt1316
- Chen W, Yun M-S, Deng F, Yogo Y (2005) The rates of maize growth and lignin biosynthesis change after root-applied chalcone. *Weed Biology and Management* 5:118–122. doi: 10.1111/j.1445-6664.2005.00165.x
- Chen W-J, Yun M-S, Deng F, Yogo Y (2011) Chalcone suppresses lignin biosynthesis in illuminated soybean cells. *Weed Biology and Management* 11:49–56. doi: 10.1111/j.1445-6664.2011.00404.x
- Chen WEIJUN, Yun MINSOO, Deng FAN, et al. (2004) Effects of root-applied naringenin and chalcone on the growth of annual plants. *Weed Biology and Management* 238:235–238.
- Chisti Y (2007) Biodiesel from microalgae. *Biotechnology advances* 25:294–306. doi: 10.1016/j.biotechadv.2007.02.001
- Chua TK, Bujnicki JM, Tan T-C, et al. (2008) The structure of sucrose phosphate synthase from *Halotheothrix orenii* reveals its mechanism of action and binding mode. *The Plant cell* 20:1059–72. doi: 10.1105/tpc.107.051193
- Conesa A, Götz S, García-Gómez JM, et al. (2005) Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics (Oxford, England)* 21:3674–6. doi: 10.1093/bioinformatics/bti610
- Darley CP, Forrester AM, McQueen-Mason SJ (2001) The molecular basis of plant cell wall extension. *Plant molecular biology* 47:179–95.
- Demura T, Fukuda H (2007) Transcriptional regulation in wood formation. *Trends in plant science* 12:64–70. doi: 10.1016/j.tplants.2006.12.006
- Deng F, Aoki M, Yogo Y (2004) Effect of naringenin on the growth and lignin biosynthesis of gramineous plants. *Weed Biology and Management* 4:49–55. doi: 10.1111/j.1445-6664.2003.00119.x

- Dixon RA, Harrison MJ (1990) Activation, structure, and organization of genes involved in microbial defense in plants. *Advances in genetics* 28:165–234.
- Dixon RA, Lamb CJ, Masoud S, et al. (1996) Metabolic engineering: prospects for crop improvement through the genetic manipulation of phenylpropanoid biosynthesis and defense responses — a review. *Gene* 179:61–71. doi: 10.1016/S0378-1119(96)00327-7
- Dixon RA, Paiva NL (1995) Stress-Induced Phenylpropanoid Metabolism. *The Plant cell* 7:1085–1097. doi: 10.1105/tpc.7.7.1085
- Ferrer J-L, Austin MB, Stewart C, Noel JP (2008) Structure and function of enzymes involved in the biosynthesis of phenylpropanoids. *Plant physiology and biochemistry : PPB / Société française de physiologie végétale* 46:356–70. doi: 10.1016/j.plaphy.2007.12.009
- Foelkel C (2009) Papermaking Properties of Eucalyptus Trees , Woods , and Pulp Fibers Celso Foelkel. In: *Eucalyptus Online Book & newsletter*. <http://www.eucalyptus.com.br/eucalptos/ENG14.pdf>.
- Foucart C, Jauneau A, Gion J-M, et al. (2009) Overexpression of EgROP1, a Eucalyptus vascular-expressed Rac-like small GTPase, affects secondary xylem formation in *Arabidopsis thaliana*. *The New phytologist* 183:1014–29. doi: 10.1111/j.1469-8137.2009.02910.x
- Fu D, Mazza G, Tamaki Y (2010) Lignin extraction from straw by ionic liquids and enzymatic hydrolysis of the cellulosic residues. *Journal of agricultural and food chemistry* 58:2915–22. doi: 10.1021/jf903616y
- Fukuda H (1996) XYLOGENESIS: INITIATION, PROGRESSION, AND CELL DEATH. *Annual review of plant physiology and plant molecular biology* 47:299–325. doi: 10.1146/annurev.arplant.47.1.299
- Gerlach D (1969) *Botanische Mikrotechnik: Eine Einführung*. Georg Thieme, Stuttgart
- Goicoechea M, Lacombe E, Legay S, et al. (2005) EgMYB2, a new transcriptional activator from Eucalyptus xylem, regulates secondary cell wall formation and lignin biosynthesis. *The Plant journal : for cell and molecular biology* 43:553–67. doi: 10.1111/j.1365-313X.2005.02480.x
- Gomes FJB, Gouvea A de FG, Colodette JL, et al. (2008) Influence of content and S/G relation of the wood lignin on kraft pulping performance *Influencia do teor da O PAPEL* 69:95–105.
- Goujon T, Sibout R, Pollet B, et al. (2003) A new *Arabidopsis thaliana* mutant deficient in the expression of O-methyltransferase impacts lignins and sinapoyl esters. *Plant molecular biology* 51:973–89.
- Grattapaglia D, Vaillancourt RE, Shepherd M, et al. (2012) Progress in Myrtaceae genetics and genomics: Eucalyptus as the pivotal genus. *Tree Genetics & Genomes* 8:463–508. doi: 10.1007/s11295-012-0491-x

- Grima-pettenati J, Soler M, Leal O, Camargo E, Wang H (2012) Transcriptional Regulation of the Lignin Biosynthetic Pathway Revisited : New Players and Insights. *Advances in Botanical Research*. Elsevier, pp 173–218
- Guo D, Chen F, Inoue K, et al. (2001) Downregulation of caffeic acid 3-O-methyltransferase and caffeoyl CoA 3-O-methyltransferase in transgenic alfalfa. impacts on lignin structure and implications for the biosynthesis of G and S lignin. *The Plant cell* 13:73–88.
- Harbone J (1988) *The Flavonoids: Advances in Research Since 1980*. 624.
- Hardcastle TJ, Kelly KA (2010) baySeq: empirical Bayesian methods for identifying differential expression in sequence count data. *BMC bioinformatics* 11:422. doi: 10.1186/1471-2105-11-422
- Hinchee M, Rottmann W, Mullinax L, et al. (2009) Short-rotation woody crops for bioenergy and biofuels applications. *In vitro cellular & developmental biology Plant : journal of the Tissue Culture Association* 45:619–629. doi: 10.1007/s11627-009-9235-5
- Holton TA, Cornish EC (1995) Genetics and Biochemistry of Anthocyanin Biosynthesis. *The Plant cell* 7:1071–1083. doi: 10.1105/tpc.7.7.1071
- Hu WJ, Harding S a, Lung J, et al. (1999) Repression of lignin biosynthesis promotes cellulose accumulation and growth in transgenic trees. *Nature biotechnology* 17:808–12. doi: 10.1038/11758
- Inoue H, Yano S, Endo T, et al. (2008) Combining hot-compressed water and ball milling pretreatments to improve the efficiency of the enzymatic hydrolysis of eucalyptus. *Biotechnology for biofuels* 1:2. doi: 10.1186/1754-6834-1-2
- Johansen DA (1940) *Plant microtechnique*. McGraw-Hill, New York
- Kanehisa M (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research* 28:27–30. doi: 10.1093/nar/28.1.27
- Kirst ME, Scott J, Kirst M, et al. (2004) Coordinated Genetic Regulation of Growth and Lignin Revealed by Quantitative Trait Locus Analysis of cDNA Microarray Data in an Interspecific Backcross of Eucalyptus 1. *Society* 135:2368–2378. doi: 10.1104/pp.103.037960.At
- Koski LB, Gray MW, Lang BF, Burger G (2005) AutoFACT: an automatic functional annotation and classification tool. *BMC bioinformatics* 6:151. doi: 10.1186/1471-2105-6-151
- Lacombe E, Hawkins S, Van Doorselaere J, et al. (1997) Cinnamoyl CoA reductase, the first committed enzyme of the lignin branch biosynthetic pathway: cloning, expression and phylogenetic relationships. *The Plant journal : for cell and molecular biology* 11:429–41.
- Ladiges P, Udovicic F, Nelson G (2003) Journal Article - Australian biogeographical connections and the phylogeny of large genera in the plant family Myrtaceae - *Encyclopedia of Australian Science. Journal of Biogeography* 30:989–998.

- Lee D, Meyer K, Chapple C, Douglas CJ (1997) Antisense suppression of 4-coumarate:coenzyme A ligase activity in *Arabidopsis* leads to altered lignin subunit composition. *The Plant cell* 9:1985–98. doi: 10.1105/tpc.9.11.1985
- Legay S, Sivadon P, Blervacq A-S, et al. (2010) EgMYB1, an R2R3 MYB transcription factor from eucalyptus negatively regulates secondary cell wall formation in *Arabidopsis* and poplar. *The New phytologist* 188:774–86. doi: 10.1111/j.1469-8137.2010.03432.x
- Lepikson-Neto J, Alves A, Deckmann AC, et al. (2013) Flavonoid supplementation reduces extractives content and increases S/G ratio on *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid trees. *Bioresources* 1–11.
- Li J, Ou-Lee TM, Raba R, et al. (1993) *Arabidopsis* Flavonoid Mutants Are Hypersensitive to UV-B Irradiation. *The Plant cell* 5:171–179. doi: 10.1105/tpc.5.2.171
- Li R, Yu C, Li Y, et al. (2009) SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics (Oxford, England)* 25:1966–7. doi: 10.1093/bioinformatics/btp336
- Li X, Ximenes E, Kim Y, et al. (2010) Lignin monomer composition affects *Arabidopsis* cell-wall degradability after liquid hot water pretreatment. *Biotechnology for biofuels* 3:27. doi: 10.1186/1754-6834-3-27
- Lim E-K, Jackson RG, Bowles DJ (2005) Identification and characterisation of *Arabidopsis* glycosyltransferases capable of glucosylating coniferyl aldehyde and sinapyl aldehyde. *FEBS letters* 579:2802–6. doi: 10.1016/j.febslet.2005.04.016
- Mansfield SD (2009) Solutions for dissolution--engineering cell walls for deconstruction. *Current opinion in biotechnology* 20:286–94. doi: 10.1016/j.copbio.2009.05.001
- Mansfield SD, Kang K-Y, Chapple C (2012) Designed for deconstruction--poplar trees altered in cell wall lignification improve the efficacy of bioethanol production. *The New phytologist* 194:91–101. doi: 10.1111/j.1469-8137.2011.04031.x
- Mavandad M, Edwards R, Liang X, et al. (1990) Effects of trans-Cinnamic Acid on Expression of the Bean Phenylalanine Ammonia-Lyase Gene Family. *Plant physiology* 94:671–80.
- Miller G. (1959) Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. *Analytical Chemistry*. *Analytical chemistry* 31:426–428.
- Mizrachi E, Mansfield SD, Myburg A a. (2011) Cellulose factories: advancing bioenergy production from forest trees. *New Phytologist* no–no. doi: 10.1111/j.1469-8137.2011.03971.x
- Mortazavi A, Williams BA, McCue K, et al. (2008) Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nature methods* 5:621–8. doi: 10.1038/nmeth.1226
- Moustafa E (1967) Purification and properties of chalcone-flavanone isomerase from soya bean seed. *Phytochemistry* 6:625–632. doi: 10.1016/S0031-9422(00)86001-X

- Nishizawa A, Yabuta Y, Shigeoka S (2008) Galactinol and raffinose constitute a novel function to protect plants from oxidative damage. *Plant physiology* 147:1251–63. doi: 10.1104/pp.108.122465
- Paiva JAP, Prat E, Vautrin S, et al. (2011) Advancing Eucalyptus genomics: identification and sequencing of lignin biosynthesis genes from deep-coverage BAC libraries. *BMC genomics* 12:137. doi: 10.1186/1471-2164-12-137
- Paux E, Carocha V, Marques C, et al. (2005) Transcript profiling of Eucalyptus xylem genes during tension wood formation. *The New phytologist* 167:89–100. doi: 10.1111/j.1469-8137.2005.01396.x
- Peterbauer T, Mucha J, Mach L, Richter A (2002) Chain Elongation of raffinose in pea seeds. Isolation, characterization, and molecular cloning of multifunctional enzyme catalyzing the synthesis of stachyose and verbascose. *The Journal of biological chemistry* 277:194–200. doi: 10.1074/jbc.M109734200
- Phillips CM, Iavarone AT, Marletta MA (2011) Quantitative proteomic approach for cellulose degradation by *Neurospora crassa*. *Journal of proteome research* 10:4177–85. doi: 10.1021/pr200329b
- Plomion C, Leprovost G, Stokes A (2001) Wood Formation in Trees. *PLANT PHYSIOLOGY* 127:1513–1523. doi: 10.1104/pp.010816
- Le Provost G, Paiva J, Pot D, et al. (2003) Seasonal variation in transcript accumulation in wood-forming tissues of maritime pine (*Pinus pinaster* Ait.) with emphasis on a cell wall glycine-rich protein. *Planta* 217:820–30. doi: 10.1007/s00425-003-1051-2
- Pruesse E, Quast C, Knittel K, et al. (2007) SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic acids research* 35:7188–96. doi: 10.1093/nar/gkm864
- Pu Y, Zhang D, Singh PM, Ragauskas AJ (2008) The new forestry biofuels sector. *Biofuels, Bioproducts and Biorefining* 2:58–73. doi: 10.1002/bbb.48
- Quail MA, Kozarewa I, Smith F, et al. (2008) A large genome center's improvements to the Illumina sequencing system. *Nature methods* 5:1005–10. doi: 10.1038/nmeth.1270
- Ragland KW, Aerts DJ, Baker AJ (1991) Properties of Wood for Combustion Analysis. 37:161–168.
- Ralph J, Lundquist K, Brunow G, et al. (2004) Lignins: Natural polymers from oxidative coupling of 4-hydroxyphenyl- propanoids. *Phytochemistry Reviews* 3:29–60. doi: 10.1023/B:PHYT.0000047809.65444.a4
- Rengel D, San Clemente H, Servant F, et al. (2009) A new genomic resource dedicated to wood formation in Eucalyptus. *BMC plant biology* 9:36. doi: 10.1186/1471-2229-9-36
- Richard TL (2010) Challenges in scaling up biofuels infrastructure. *Science (New York, NY)* 329:793–6. doi: 10.1126/science.1189139

- Rodrigues J, Graça J, Pereira H (2001) Influence of tree eccentric growth on syringyl/guaiacyl ratio in *Eucalyptus globulus* wood lignin assessed by analytical pyrolysis. *Journal of Analytical and Applied Pyrolysis* 58-59:481–489. doi: 10.1016/S0165-2370(00)00121-2
- Rodrigues J, Meier D, Faix O, Pereira H (1999) Determination of tree to tree variation in syringyl/guaiacyl ratio of *Eucalyptus globulus* wood lignin by analytical pyrolysis. *Journal of Analytical and Applied Pyrolysis* 48:121–128. doi: 10.1016/S0165-2370(98)00134-X
- Rubin G, Tohge T, Matsuda F, et al. (2009) Members of the LBD family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in *Arabidopsis*. *The Plant cell* 21:3567–84. doi: 10.1105/tpc.109.067041
- Ruegger M, Meyer K, Cusumano JC, Chapple C (1999) Regulation of Ferulate-5-Hydroxylase Expression in *Arabidopsis* in the Context of Sinapate Ester Biosynthesis. *PLANT PHYSIOLOGY* 119:101–110. doi: 10.1104/pp.119.1.101
- Saravitz DM, Pharr DM, Carter TE (1987) Galactinol synthase activity and soluble sugars in developing seeds of four soybean genotypes. *Plant physiology* 83:185–9.
- Sarruje J (1975) Soluções nutritivas. *Summa Phytopathologica* v.1:p.231–233,.
- Schwanninger, Manfred / Hinterstoisser B (2002) Klason Lignin: Modifications to Improve the Precision of the Standardized Determination : *Holzforschung*. 56:161–166.
- Shamir R, Maron-Katz A, Tanay A, et al. (2005) EXPANDER--an integrative program suite for microarray data analysis. *BMC bioinformatics* 6:232. doi: 10.1186/1471-2105-6-232
- Somerville C, Youngs H, Taylor C, et al. (2010) Feedstocks for lignocellulosic biofuels. *Science (New York, NY)* 329:790–2. doi: 10.1126/science.1189268
- Sticklen MB (2008) Plant genetic engineering for biofuel production: towards affordable cellulosic ethanol. *Nature reviews Genetics* 9:433–43. doi: 10.1038/nrg2336
- Suzek BE, Huang H, McGarvey P, et al. (2007) UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics (Oxford, England)* 23:1282–8. doi: 10.1093/bioinformatics/btm098
- Séguin A (2011) How could forest trees play an important role as feedstock for bioenergy production? *Current Opinion in Environmental Sustainability* 3:90–94. doi: 10.1016/j.cosust.2010.12.006
- Taji T, Ohsumi C, Iuchi S, et al. (2002) Important roles of drought- and cold-inducible genes for galactinol synthase in stress tolerance in *Arabidopsis thaliana*. *The Plant journal : for cell and molecular biology* 29:417–26.
- Tamagnone L, Merida A, Parr A, et al. (1998) The AmMYB308 and AmMYB330 transcription factors from *Antirrhinum* regulate phenylpropanoid and lignin biosynthesis in transgenic tobacco. *The Plant cell* 10:135–54.

- Thimm O, Bläsing O, Gibon Y, et al. (2004) MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *The Plant journal : for cell and molecular biology* 37:914–39.
- Tognetti VB, Van Aken O, Morreel K, et al. (2010) Perturbation of indole-3-butyric acid homeostasis by the UDP-glucosyltransferase UGT74E2 modulates Arabidopsis architecture and water stress tolerance. *The Plant cell* 22:2660–79. doi: 10.1105/tpc.109.071316
- Trapnell C, Pachter L, Salzberg SL (2009) TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics (Oxford, England)* 25:1105–11. doi: 10.1093/bioinformatics/btp120
- Tronchet M, Balagué C, Kroj T, et al. (2010) Cinnamyl alcohol dehydrogenases-C and D, key enzymes in lignin biosynthesis, play an essential role in disease resistance in Arabidopsis. *Molecular plant pathology* 11:83–92. doi: 10.1111/j.1364-3703.2009.00578.x
- Unda F (2012) EVALUATING THE ROLE OF THE RAFFINOSE FAMILY OF IN HYBRID POPLAR (*Populus alba* × *grandidentata*). The University of British Columbia
- Unda F, Canam T, Preston L, Mansfield SD (2012) Isolation and characterization of galactinol synthases from hybrid poplar. *Journal of experimental botany* 63:2059–69. doi: 10.1093/jxb/err411
- UNICA- União da indústria de cana de açúcar; <http://www.unica.com.br/>
- Vanholme R, Demedts B, Morreel K, et al. (2010) Lignin biosynthesis and structure. *Plant physiology* 153:895–905. doi: 10.1104/pp.110.155119
- Vogt T (2010) Phenylpropanoid biosynthesis. *Molecular plant* 3:2–20. doi: 10.1093/mp/ssp106
- Voo KS, Whetten RW, O'Malley DM, Sederoff RR (1995) 4-coumarate:coenzyme a ligase from loblolly pine xylem. Isolation, characterization, and complementary DNA cloning. *Plant physiology* 108:85–97.
- Wagner A, Donaldson L, Kim H, et al. (2009) Suppression of 4-Coumarate-CoA Ligase in the Coniferous Gymnosperm *Pinus radiata* 1 [W]. *Society* 149:370–383. doi: 10.1104/pp.108.125765
- Wang W, Vinocur B, Shoseyov O, Altman A (2004) Role of plant heat-shock proteins and molecular chaperones in the abiotic stress response. *Trends in plant science* 9:244–52. doi: 10.1016/j.tplants.2004.03.006
- Weng J-K, Akiyama T, Bonawitz ND, et al. (2010) Convergent evolution of syringyl lignin biosynthesis via distinct pathways in the lycophyte *Selaginella* and flowering plants. *The Plant cell* 22:1033–45. doi: 10.1105/tpc.109.073528
- Weng J-K, Li X, Bonawitz ND, Chapple C (2008) Emerging strategies of lignin engineering and degradation for cellulosic biofuel production. *Current opinion in biotechnology* 19:166–72. doi: 10.1016/j.copbio.2008.02.014

- Yáñez-S M, Rojas J, Castro J, et al. (2012) Fuel ethanol production from Eucalyptus globulus wood by autocatalized organosolv pretreatment ethanol-water and SSF. Journal of Chemical Technology & Biotechnology n/a–n/a.
- Zeng Y, Yang T (2002) RNA isolation from highly viscous samples rich in polyphenols and polysaccharides. Plant Molecular Biology Reporter 20:417–417. doi: 10.1007/BF02772130
- Zhang J, Elo A, Helariutta Y (2011) Arabidopsis as a model for wood formation. Current opinion in biotechnology 22:293–9. doi: 10.1016/j.copbio.2010.11.008
- Zhao Q, Dixon RA (2011) Transcriptional networks for lignin biosynthesis: more complex than we thought? Trends in plant science 16:227–33. doi: 10.1016/j.tplants.2010.12.005
- Zhong R, Lee C, Ye Z-H (2010) Evolutionary conservation of the transcriptional network regulating secondary cell wall biosynthesis. Trends in plant science 15:625–32. doi: 10.1016/j.tplants.2010.08.007
- Zhong R, Lee C, Zhou J, et al. (2008) A battery of transcription factors involved in the regulation of secondary cell wall biosynthesis in Arabidopsis. The Plant cell 20:2763–82. doi: 10.1105/tpc.108.061325
- Zhong R, Richardson EA, Ye Z-H (2007) The MYB46 transcription factor is a direct target of SND1 and regulates secondary wall biosynthesis in Arabidopsis. The Plant cell 19:2776–92. doi: 10.1105/tpc.107.053678
- Zhong R, Ye Z-H (2007) Regulation of cell wall biosynthesis. Current opinion in plant biology 10:564–72. doi: 10.1016/j.pbi.2007.09.001
- Zhu JY, Pan XJ (2010) Woody biomass pretreatment for cellulosic ethanol production: Technology and energy consumption evaluation. Bioresource technology 101:4992–5002.

Folha de rosto do pedido de patente depositado junto ao INPI relativo à formulação de uma composição para a modificação química da madeira de eucalipto utilizando a suplementação por flavonóides.

Formulário 1.01 – Depósito de Pedido de Patente ou de Certificado de Adição (folha 1/2)

7. Declaração na forma do item 3.2 do Ato Normativo nº 127/97:

☐ 7.1 Declaro que os dados fornecidos no presente formulário são idênticos ao da certidão de depósito ou documento equivalente do pedido cuja prioridade está sendo reivindicada.

☐ em anexo

8. Declaração de divulgação anterior não prejudicial: (Período de Graça):
(art. 12 da LPI e item 2 do AN nº 127/97)

☐ em anexo

9. Procurador (74)

9.1 Nome: FERNANDA LAVRAS COSTALLAT SILVADO

9.2 CNPJ/CPF: 295.166.068-57

9.3 API/OAB: 210.899

9.4 Endereço completo: PROCURADORIA GERAL DA UNICAMP, CAMPINAS - SP

9.5 CEP: 13083-970

9.6 Telefone: 19 3521-4771

9.7 Fax: 00 0000-0000

9.8 E-Mail: patentes@inova.unicamp.br

10. Listagem de sequências Biológicas (documentos anexados) (se houver):

☐ Listagem de sequências em arquivo eletrônico: nº de CDs ou DVDs (original e cópia).

☐ Código de controle alfanumérico no formato de código de barras: fl.

☐ Listagem de sequências em formato impresso: fls.

☐ Declaração de acordo com o artigo da Resolução INPI nº 228/09: fls.

11. Documentos anexados (assinale e indique também o número de folhas):

(Deverá ser indicado o nº total de somente uma das vias de cada documento)

☒	11.1 Guia de Recolhimento	1 fls.	☒	11.5 Relatório descritivo	2.1 fls.
☒	11.2 Procuração	2 fls.	☒	11.6 Reivindicações	4 fls.
☐	11.3 Documentos de Prioridade	fls.	☐	11.7 Desenhos	fls.
☐	11.4 Doc. de contrato de trabalho	fls.	☒	11.8 Resumo	1 fls.
☐	11.9 Outros que não aqueles definidos no campo 11 (especificar) ANEXOS				10 fls.

12. Total de folhas anexadas (referentes aos campos 10 e 11): 39 fls.

13. Declaro, sob penas da Lei, que todas as informações acima prestadas são completas e verdadeiras.

CAMPINAS, SP, DE OUTUBRO DE 2012

Local e Data



Assinatura e Carimbo

Fernanda Lavras Costallat Silvado,
Procuradora de Universidade Subchefe
Matricula nº 28.574-2
OAB/SP nº 210.899

Anexo 2. Identidade, valores de expressão e anotação dos 1573 genes diferencialmente expressos.

GENE ID	FPKM					Annotation	
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR	
Eucgr.F04479.1	0,12	35,55	40,44	23,84	57,92	AT1G53540.1	Symbols: HSP20-like chaperones superfamily protein chr1:19980510-19980983 FORWARD LENGTH=157
Eucgr.K02389.1	0,04	13,69	10,18	9,89	26,99	AT3G10020.2	Symbols: unknown protein
Eucgr.K02399.1	0,08	18,09	18,67	19,48	53,82	AT3G10020.1	Symbols: unknown protein
Eucgr.G01188.2	2,78	45,78	41,56	33,40	76,35	AT1G17870.1	Symbols: ATEGY3, EGY3 ethylene-dependent gravitropism-deficient and yellow-green-like 3
Eucgr.J01979.1	0,34	17,61	19,94	14,63	27,21	AT5G59720.1	Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.K02410.1	0,13	14,27	10,66	12,52	29,61	AT3G10020.1	Symbols: unknown protein
Eucgr.J01980.1	0,02	12,20	11,59	9,15	26,39	AT5G59720.1	Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.L02233.1	1,29	41,47	38,15	29,24	118,40	No hits found	
Eucgr.C03056.2	4,97	32,66	25,61	15,26	91,38	AT2G26150.1	Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2 chr2:11135856-11137217 FORWARD LENGTH=345
Eucgr.F02898.1	2,76	525,99	343,17	280,68	342,59	AT5G37670.1	Symbols: HSP20-like chaperones superfamily protein chr5:14969035-14969448 FORWARD LENGTH=137
Eucgr.J01985.1	0,31	22,25	16,62	13,21	33,33	AT5G59720.1	Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.K03553.1	0,04	4,11	4,59	2,83	6,87	AT4G01970.1	Symbols: AtSTS, STS stachyose synthase chr4:854073-856953 REVERSE LENGTH=876
Eucgr.L03261.1	1,44	47,41	27,81	19,98	75,67	AT5G59720.1	Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.C03449.1	0,29	14,55	12,98	14,36	18,58	AT2G26150.1	Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2 chr2:11135856-11137217 FORWARD LENGTH=345
Eucgr.C00684.1	2,00	341,30	299,30	243,65	272,90	AT5G12020.1	Symbols: HSP17.6II 17.6 kDa class II heat shock protein chr5:3882409-3882876 REVERSE LENGTH=155
Eucgr.J01972.1	5,54	85,59	56,26	32,51	108,67	AT5G59720.1	Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.K02384.1	0,07	14,56	10,59	10,60	27,23	AT3G10020.1	Symbols: unknown protein
Eucgr.J01969.1	4,89	192,23	134,53	103,09	310,44	AT2G29500.1	Symbols: HSP20-like chaperones superfamily protein chr2:12633279-12633740 REVERSE LENGTH=153
Eucgr.K03472.1	0,07	109,57	84,15	62,04	20,94	AT2G47710.1	Symbols: Adenine nucleotide alpha hydrolases-like superfamily protein
Eucgr.H04513.1	0,23	15,72	18,79	11,56	21,62	AT5G28540.1	Symbols: BIP1 heat shock protein 70 (Hsp 70) family protein
Eucgr.A00595.1	0,10	81,17	76,66	57,33	98,83	AT5G01300.1	Symbols: PEBP (phosphatidylethanolamine-binding protein) family protein
Eucgr.E02421.1	0,19	260,04	220,12	119,63	51,05	AT1G30190.1	Symbols: unknown protein
Eucgr.H04692.1	2,97	83,31	59,57	43,11	313,22	AT4G27670.1	Symbols: HSP21 heat shock protein 21 chr4:13819048-13819895 REVERSE LENGTH=227
Eucgr.G02440.1	0,00	5,46	5,80	3,33	4,06	AT4G34135.1	Symbols: UGT73B2 UDP-glucosyltransferase 73B2 chr4:16345476-16347016 REVERSE LENGTH=483
Eucgr.J01978.1	1,35	36,39	28,77	20,90	71,33	AT1G53540.1	Symbols: HSP20-like chaperones superfamily protein chr1:19980510-19980983 FORWARD LENGTH=157
Eucgr.J01977.1	0,37	9,43	8,59	7,08	19,26	AT5G59720.1	Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.G02259.1	0,00	2,73	2,14	1,18	3,10	AT4G34131.1	Symbols: UGT73B3 UDP-glucosyl transferase 73B3 chr4:16343268-16344713 REVERSE LENGTH=481

GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	
Eucgr.J01959.1	3,19	142,90	89,76	58,22	148,21	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.J01982.1	0,21	8,65	6,06	4,24	13,03	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.K00295.1	2,11	46,13	38,25	35,52	62,69	AT5G52640.1 Symbols: HSP81-1, ATHS83, HSP81.1, HSP83, ATHSP90.1, AtHsp90-1, HSP90.1 heat shock protein 90.1
Eucgr.A01833.1	0,13	32,00	24,43	15,08	10,47	AT4G28390.1 Symbols: AAC3, ATAAC3 ADP/ATP carrier 3 chr4:14041486-14042781 REVERSE LENGTH=379
Eucgr.J01964.1	13,65	145,21	119,05	111,04	251,88	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.C03071.1	3,64	517,08	509,51	451,66	324,54	AT5G12020.1 Symbols: HSP17.6II 17.6 kDa class II heat shock protein chr5:3882409-3882876 REVERSE LENGTH=155
Eucgr.B03843.1	1,45	93,17	63,39	67,61	20,77	No hits found
Eucgr.I01982.1	13,71	110,50	95,01	54,71	184,40	AT5G15450.1 Symbols: APG6, CLPB3, CLPB-P casein lytic proteinase B3 chr5:5014399-5018255 REVERSE LENGTH=968
Eucgr.C03320.1	0,38	24,39	18,22	14,01	9,78	AT5G20670.1 Symbols: Protein of unknown function (DUF1677) chr5:6993332-6993793 REVERSE LENGTH=153
Eucgr.B00176.2	3,86	153,25	109,01	82,97	57,39	AT5G50240.1 Symbols: PIMT2, AtPIMT2 protein-L-isoaspartate methyltransferase 2
Eucgr.J02588.1	3,20	225,90	182,76	194,01	130,04	No hits found
Eucgr.C00690.1	2,48	563,68	498,40	514,67	458,94	AT5G12020.1 Symbols: HSP17.6II 17.6 kDa class II heat shock protein chr5:3882409-3882876 REVERSE LENGTH=155
Eucgr.H01413.1	0,61	14,45	13,47	8,20	14,25	AT1G03070.2 Symbols: Bax inhibitor-1 family protein chr1:730148-731379 FORWARD LENGTH=247
Eucgr.H02887.1	4,80	37,80	28,32	19,46	30,12	AT4G22340.1 Symbols: CDS2 cytidinediphosphate diacylglycerol synthase 2 chr4:11800150-11802549 REVERSE LENGTH=423
Eucgr.G01414.1	0,04	1,91	1,28	1,10	2,80	AT4G34131.1 Symbols: UGT73B3 UDP-glucosyl transferase 73B3 chr4:16343268-16344713 REVERSE LENGTH=481
Eucgr.J01981.1	2,00	34,43	50,34	69,86	105,48	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.K00237.1	0,04	115,83	64,41	61,47	11,49	AT5G62040.1 Symbols: PEBP (phosphatidylethanolamine-binding protein) family protein
Eucgr.F03196.1	1,43	292,73	240,63	168,71	38,29	AT1G17180.1 Symbols: ATGSTU25, GSTU25 glutathione S-transferase TAU 25
Eucgr.I02136.1	1,68	226,73	147,43	90,02	259,35	AT1G54050.1 Symbols: HSP20-like chaperones superfamily protein chr1:20179558-20180122 REVERSE LENGTH=155
Eucgr.H04427.1	49,92	0,52	0,85	1,32	13,93	AT3G06350.1 Symbols: EMB3004, MEE32 dehydroquinase dehydratase, putative / shikimate dehydrogenase, putative
Eucgr.L02860.1	2,44	408,70	383,21	397,30	273,12	AT5G12020.1 Symbols: HSP17.6II 17.6 kDa class II heat shock protein chr5:3882409-3882876 REVERSE LENGTH=155
Eucgr.B03280.1	0,36	165,28	127,02	117,88	51,95	AT1G63840.1 Symbols: RING/U-box superfamily protein chr1:23689991-23690491 REVERSE LENGTH=166
Eucgr.C04378.1	1,60	7,63	9,88	6,52	8,45	AT1G50030.1 Symbols: TOR target of rapamycin chr1:18522626-18539619 REVERSE LENGTH=2481
Eucgr.L01770.1	0,17	15,89	16,32	10,41	19,09	AT5G28540.1 Symbols: BIP1 heat shock protein 70 (Hsp 70) family protein
Eucgr.L02697.1	0,02	1,00	1,30	0,92	1,27	AT5G53990.1 Symbols: UDP-Glycosyltransferase superfamily protein chr5:21915707-21917050 REVERSE LENGTH=447
Eucgr.H03046.1	1,04	8,76	4,89	4,14	11,58	AT1G65660.1 Symbols: SMP1 Pre-mRNA splicing Prp18-interacting factor
Eucgr.L02363.1	0,24	39,93	126,78	154,82	28,88	AT5G47550.1 Symbols: Cystatin/monellin superfamily protein chr5:19286596-19286964 REVERSE LENGTH=122
Eucgr.K01389.2	19,81	1348,64	796,35	569,70	120,14	AT2G04240.2 Symbols: XERICO RING/U-box superfamily protein chr2:1461816-1462304 REVERSE LENGTH=162
Eucgr.D02334.1	1,64	250,51	223,70	129,50	253,35	AT2G46240.1 Symbols: BAG6, ATBAG6 BCL-2-associated athanogene 6

Eucgr.H00118.1	27,73	1624,22	1433,16	1046,96	372,67	AT1G04560.1 Symbols: AWP19-like family protein chr1:1245070-1245888 FORWARD LENGTH=186
GENE ID	FPKM	Annotation				
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.B02743.1	0,06	5,44	6,05	6,01	10,26	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.J02137.1	2,47	27,99	28,19	18,78	15,91	AT5G15070.1 Symbols: Phosphoglycerate mutase-like family protein chr5:4876898-4885615 FORWARD LENGTH=1049
Eucgr.F03363.1	1,18	115,06	100,03	71,61	51,24	AT2G20900.2 Symbols: DGK5, ATDGK5 diacylglycerol kinase 5 chr2:8989794-8992798 REVERSE LENGTH=491
Eucgr.K02873.1	0,31	9,45	11,35	7,36	4,06	AT5G62890.3 Symbols: Xanthine/uracil permease family protein chr5:25243723-25247075 FORWARD LENGTH=532
Eucgr.J01958.1	2,68	115,32	90,34	65,56	100,03	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.K01402.1	0,68	489,73	335,43	243,43	159,69	No hits found
Eucgr.H01638.1	7,89	123,54	93,85	65,50	121,11	AT1G03070.2 Symbols: Bax inhibitor-1 family protein chr1:730148-731379 FORWARD LENGTH=247
Eucgr.H03044.1	0,96	11,74	5,01	4,78	12,51	AT1G65660.1 Symbols: SMP1 Pre-mRNA splicing Prp18-interacting factor chr1:24418295-24421231 REVERSE LENGTH=535
Eucgr.B02244.1	0,12	11,57	8,11	6,82	11,20	AT2G30000.1 Symbols: PHF5-like protein chr2:12804042-12804374 REVERSE LENGTH=110
Eucgr.D01874.3	7,75	39,28	36,64	29,74	27,09	AT1G12800.1 Symbols: Nucleic acid-binding, OB-fold-like protein chr1:4361778-4365189 REVERSE LENGTH=767
Eucgr.A01806.1	17,58	0,52	1,62	3,50	4,84	AT4G28250.1 Symbols: ATEXPB3, EXPB3, ATHEXP BETA 1.6 expansin B3 chr4:14000446-14001945 REVERSE LENGTH=264
Eucgr.J00827.1	0,04	0,90	1,56	0,98	0,76	AT5G13000.1 Symbols: ATGSL12, gsl12 glucan synthase-like 12 chr5:4110445-4121202 REVERSE LENGTH=1955
Eucgr.K03563.1	0,23	4,48	11,12	12,00	8,48	AT2G47180.1 Symbols: AtGolS1, GolS1 galactinol synthase 1 chr2:19369049-19370372 REVERSE LENGTH=344
Eucgr.B03963.1	0,01	0,78	1,33	1,08	0,51	AT3G62150.1 Symbols: PGP21 P-glycoprotein 21 chr3:23008755-23013579 REVERSE LENGTH=1296
Eucgr.B02291.1	10,86	52,79	42,69	39,45	45,69	AT3G46670.1 Symbols: UGT76E11 UDP-glucosyl transferase 76E11 chr3:17192795-17194227 REVERSE LENGTH=451
Eucgr.E02947.1	0,37	8,89	9,66	7,62	7,69	AT1G16770.1 Symbols: unknown protein
Eucgr.D00704.1	5,24	77,32	63,35	45,93	125,37	AT1G61260.1 Symbols: Protein of unknown function (DUF761) chr1:22593756-22594986 REVERSE LENGTH=344
Eucgr.D01410.1	25,73	0,43	0,86	1,34	6,87	AT3G07880.1 Symbols: SCN1 Immunoglobulin E-set superfamily protein chr3:2514175-2515544 FORWARD LENGTH=240
Eucgr.I02375.1	25,29	1,93	4,94	6,67	8,37	AT5G66330.1 Symbols: Leucine-rich repeat (LRR) family protein chr5:26500531-26501787 REVERSE LENGTH=418
Eucgr.G01464.1	0,16	19,83	11,39	8,24	8,14	No hits found
Eucgr.J01960.1	4,43	365,87	267,66	190,17	357,50	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.H03725.1	5,38	13,43	16,12	13,10	20,16	AT4G13550.1 Symbols: triglyceride lipases;triglyceride lipases chr4:7871251-7876160 REVERSE LENGTH=715
Eucgr.D01040.2	15,96	134,23	130,11	126,16	81,83	AT5G22480.1 Symbols: ZPR1 zinc-finger domain protein chr5:7451643-7456168 REVERSE LENGTH=493
Eucgr.A02381.1	4,49	172,85	136,95	91,17	101,55	AT5G12840.3 Symbols: HAP2A, ATHAP2A, NF-YA1 nuclear factor Y, subunit A1
Eucgr.I00312.1	2,82	14,42	15,28	12,03	15,91	AT2G21860.1 Symbols: violaxanthin de-epoxidase-related chr2:9318333-9319990 REVERSE LENGTH=522
Eucgr.H04079.1	9,38	74,45	65,19	42,14	40,33	AT3G17180.1 Symbols: scpl33 serine carboxypeptidase-like 33 chr3:5855861-5859157 REVERSE LENGTH=478
Eucgr.G02845.1	9,82	0,00	0,19	0,25	2,74	AT5G03250.1 Symbols: Phototropic-responsive NPH3 family protein chr5:774591-776855 FORWARD LENGTH=592
Eucgr.K02208.2	18,76	0,74	0,60	1,39	0,92	AT5G35525.1 Symbols: PLAC8 family protein chr5:13707084-13707818 FORWARD LENGTH=152

Eucgr.G02748.1	17,47	0,31	0,44	0,60	5,40	AT5G04310.1 Symbols: Pectin lyase-like superfamily protein chr5:1203356-1207352 REVERSE LENGTH=518
Eucgr.F03704.1	1,45	27,74	21,67	11,53	10,03	AT1G78610.1 Symbols: MSL6 mechanosensitive channel of small conductance-like 6 chr1:29569226-29572126 REVERSE LENGTH=856
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.I01368.1	4,42	135,89	110,06	83,93	39,49	AT4G38810.2 Symbols: Calcium-binding EF-hand family protein chr4:18115607-18118860 REVERSE LENGTH=375
Eucgr.D02073.1	24,48	469,78	380,75	330,52	1060,38	AT4G25200.1 Symbols: ATHSP23.6-MITO, HSP23.6-MITO
Eucgr.G02781.1	0,16	80,65	61,04	42,50	3,75	AT5G07330.1 Symbols: unknown protein
Eucgr.H03519.1	0,38	4,60	4,12	1,63	5,88	No hits found
Eucgr.H03564.1	2,05	0,42	0,08	0,12	0,10	AT5G19260.1 Symbols: Protein of unknown function (DUF3049) chr5:6479494-6480360 REVERSE LENGTH=288
Eucgr.G01238.2	2,34	33,68	37,22	26,67	20,98	AT3G18990.1 Symbols: VRN1, REM39 AP2/B3-like transcriptional factor family protein chr3:6549077-6551568 REVERSE LENGTH=341
Eucgr.H04153.1	7,60	33,07	24,65	21,90	31,09	AT3G17611.1 Symbols: ATRBL14, RBL14 RHOMBOID-like protein 14 chr3:6024946-6026173 FORWARD LENGTH=334
Eucgr.H00260.2	4,36	52,47	46,33	43,37	29,63	AT3G03080.1 Symbols: Zinc-binding dehydrogenase family protein chr3:698530-700278 REVERSE LENGTH=350
Eucgr.B03897.1	11,62	57,07	44,48	34,48	68,19	AT1G27650.1 Symbols: ATU2AF35A U2 snRNP auxiliary factor small subunit, putative chr1:9615152-9616042 FORWARD LENGTH=296
Eucgr.D00856.1	24,31	247,98	198,32	196,32	181,11	AT4G21320.1 Symbols: HSA32 Aldolase-type TIM barrel family protein chr4:11340492-11341732 FORWARD LENGTH=286
Eucgr.C01031.1	2,73	476,52	301,85	296,55	156,41	AT1G52240.2 Symbols: ATROPGEF11, ROPGEF11, PIRF1 RHO guanyl-nucleotide exchange factor 11 chr1:19458844-19459235 REVERSE LENGTH=94
Eucgr.H03386.1	0,25	64,14	50,54	45,19	12,54	No hits found
Eucgr.C00553.1	0,06	7,73	4,99	3,03	2,70	AT4G32900.2 Symbols: Peptidyl-tRNA hydrolase II (PTH2) family protein chr4:15878944-15880979 REVERSE LENGTH=193
Eucgr.E01440.1	15,55	311,66	257,44	211,20	244,86	AT2G45380.1 Symbols: unknown protein
Eucgr.C03325.1	13,51	0,33	0,56	0,53	2,23	AT2G05920.1 Symbols: Subtilase family protein chr2:2269831-2272207 REVERSE LENGTH=754
Eucgr.L00243.1	0,02	29,83	19,77	9,75	1,37	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.J03055.1	1,30	52,54	39,45	19,47	44,88	AT1G80160.1 Symbols: Lactoylglutathione lyase / glyoxalase I family protein chr1:30151101-30151930 FORWARD LENGTH=167
Eucgr.K00185.1	18,10	65,15	61,81	52,54	59,96	AT5G51820.1 Symbols: PGM, ATPGMP, PGM1, STF1 phosphoglucomutase
Eucgr.B02319.1	1,11	62,29	41,02	20,31	18,83	No hits found
Eucgr.J01986.1	15,59	1423,51	1288,64	1295,51	1725,77	No hits found
Eucgr.F03317.1	7,94	99,41	63,55	61,29	61,03	No hits found
Eucgr.H03518.1	6,06	282,04	218,12	152,91	249,47	AT2G32120.2 Symbols: HSP70T-2 heat-shock protein 70T-2 chr2:13651720-13653411 REVERSE LENGTH=563
Eucgr.H04961.1	0,09	83,28	46,51	26,22	2,80	AT4G33467.2 Symbols: unknown protein
Eucgr.B03883.1	4,47	146,11	114,74	104,60	310,58	AT4G21105.1 Symbols: cytochrome-c oxidases;electron carriers chr4:11266273-11266724 FORWARD LENGTH=68
Eucgr.L03244.1	0,12	279,83	137,40	63,66	2,99	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335

Eucgr.H03093.1	1,70	0,10	0,10	0,16	0,17	AT2G39510.1 Symbols: nodulin MtN21 /EamA-like transporter family protein
Eucgr.H03531.1	74,41	0,51	2,84	6,02	12,22	AT5G19530.1 Symbols: ACL5 S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
Eucgr.C03260.1	25,50	0,97	1,08	1,41	4,84	AT2G05920.1 Symbols: Subtilase family protein chr2:2269831-2272207 REVERSE LENGTH=754
Eucgr.G02190.1	0,54	97,19	71,42	60,16	12,44	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.I01937.1	0,63	234,53	148,33	111,94	21,63	No hits found
Eucgr.A01691.1	0,62	27,11	26,11	24,65	21,17	AT2G36970.1 Symbols: UDP-Glycosyltransferase superfamily protein chr2:15529050-15530712 FORWARD LENGTH=490
Eucgr.J02226.1	2,50	129,38	84,12	80,04	22,01	AT5G40382.1 Symbols: Cytochrome c oxidase subunit Vc family protein chr5:16155486-16155683 REVERSE LENGTH=65
Eucgr.H01945.1	6,14	62,68	72,80	52,36	31,27	AT3G50380.1 Symbols: Protein of unknown function (DUF1162) chr3:18686527-18700533 REVERSE LENGTH=3072
Eucgr.A02785.1	12,29	0,68	1,90	3,40	2,59	AT3G48970.1 Symbols: Heavy metal transport/detoxification superfamily protein
Eucgr.K03449.2	8,93	62,92	60,36	43,02	30,45	AT4G03200.1 Symbols: catalytics chr4:1408296-1412566 FORWARD LENGTH=818
Eucgr.E02934.1	0,08	4,87	7,41	6,39	3,64	AT1G16770.1 Symbols: unknown protein
Eucgr.L03214.1	24,68	888,01	641,74	501,21	1002,76	AT1G53540.1 Symbols: HSP20-like chaperones superfamily protein chr1:19980510-19980983 FORWARD LENGTH=157
Eucgr.F04333.1	2,83	0,05	0,29	0,41	0,38	AT1G77400.1 Symbols: unknown protein
Eucgr.H02320.1	0,63	3,53	4,39	3,01	4,07	AT5G65890.1 Symbols: ACR1 ACT domain repeat 1 chr5:26355701-26357721 FORWARD LENGTH=477
Eucgr.B00140.1	15,42	176,75	142,66	163,92	115,92	AT3G48510.1 Symbols: unknown protein
Eucgr.J01966.1	1,38	27,34	22,88	23,24	44,20	AT1G53540.1 Symbols: HSP20-like chaperones superfamily protein chr1:19980510-19980983 FORWARD LENGTH=157
Eucgr.K00957.1	1,36	17,54	17,19	10,55	15,16	AT5G54160.1 Symbols: ATOMT1, OMT1 O-methyltransferase 1 chr5:21982075-21984167 FORWARD LENGTH=363
Eucgr.I02689.1	21,71	218,24	199,47	162,07	136,90	AT1G80670.1 Symbols: Transducin/WD40 repeat-like superfamily protein
Eucgr.C00370.2	57,14	230,72	219,00	189,70	274,04	AT5G58070.1 Symbols: ATTIL, TIL temperature-induced lipocalin chr5:23500512-23501156 REVERSE LENGTH=186
Eucgr.G00061.1	9,45	834,94	858,21	726,74	814,87	AT4G10250.1 Symbols: ATHSP22.0 HSP20-like chaperones superfamily protein
Eucgr.H04349.1	0,10	0,48	1,13	0,87	0,99	AT3G63380.1 Symbols: ATPase E1-E2 type family protein / haloacid dehalogenase-like hydrolase family protein
Eucgr.I02057.1	0,00	1,43	0,99	0,73	1,01	AT3G30210.1 Symbols: ATMYB121, MYB121 myb domain protein 121 chr3:11838567-11840313 FORWARD LENGTH=276
Eucgr.C03457.2	0,17	2,38	3,72	2,39	2,36	AT2G26150.2 Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2
Eucgr.G01510.1	1,71	10,40	11,17	11,20	6,80	AT2G37710.1 Symbols: RLK receptor lectin kinase chr2:15814934-15816961 REVERSE LENGTH=675
Eucgr.C02666.1	11,79	71,43	66,15	46,70	55,43	AT2G25140.1 Symbols: HSP98.7, CLPB-M, CLPB4 casein lytic proteinase B4
Eucgr.L02232.1	0,02	0,81	0,48	0,27	0,82	AT5G59720.1 Symbols: HSP18.2 heat shock protein 18.2 chr5:24062632-24063117 FORWARD LENGTH=161
Eucgr.J02919.1	11,70	0,07	0,49	1,01	4,08	AT2G19190.1 Symbols: FRK1 FLG22-induced receptor-like kinase 1 chr2:8326067-8329893 REVERSE LENGTH=876
Eucgr.C03456.1	0,09	2,67	3,64	2,25	1,45	AT2G26150.1 Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2
Eucgr.H00781.1	4,55	36,43	23,07	20,24	37,78	AT3G07150.1 Symbols: unknown protein;

Eucgr.K02521.1	48,86	363,27	336,91	234,15	831,75	AT1G74310.1 Symbols: ATHSP101, HSP101, HOT1 heat shock protein 101
Eucgr.B02460.3	41,11	1,17	2,07	2,91	10,24	AT5G60210.2 Symbols: RIP5 ROP interactive partner 5 chr5:24243529-24245889 REVERSE LENGTH=564
Eucgr.H01544.1	3,03	72,01	54,80	52,16	36,82	AT4G02980.1 Symbols: ABP1, ABP endoplasmic reticulum auxin binding protein 1
Eucgr.H03768.1	3,91	117,64	84,62	55,77	59,05	No hits found
Eucgr.D00956.1	17,54	116,86	103,49	88,96	84,51	AT5G47810.1 Symbols: PFK2 phosphofructokinase 2 chr5:19356569-19357989 REVERSE LENGTH=444
Eucgr.H03959.1	5,98	0,07	0,05	0,07	0,61	AT3G06840.1 Symbols: unknown protein;
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.G03295.1	4,33	0,00	0,01	0,03	0,13	AT3G63470.1 Symbols: scpl40 serine carboxypeptidase-like 40 chr3:23438561-23440862 FORWARD LENGTH=502
Eucgr.F01711.1	18,73	226,64	194,05	169,82	184,15	AT5G20890.1 Symbols: TCP-1/cpn60 chaperonin family protein chr5:7087020-7089906 REVERSE LENGTH=527
Eucgr.F01656.1	0,00	8,31	3,72	1,43	1,06	AT1G75750.1 Symbols: GASA1 GAST1 protein homolog 1 chr1:28441813-28442284 REVERSE LENGTH=98
Eucgr.K02988.2	16,21	90,52	69,09	53,94	47,28	AT5G01260.1 Symbols: Carbohydrate-binding-like fold chr5:105367-106488 FORWARD LENGTH=306
Eucgr.C00382.1	11,83	39,61	36,35	36,96	30,51	AT5G58200.2 Symbols: Calcineurin-like metallo-phosphoesterase superfamily protein
Eucgr.F02405.1	0,08	1,19	2,13	1,16	0,98	AT4G31940.1 Symbols: CYP82C4 cytochrome P450, family 82, subfamily C, polypeptide 4
Eucgr.B03053.1	11,13	0,31	0,65	0,83	2,91	AT1G25570.1 Symbols: Di-glucose binding protein with Leucine-rich repeat domain
Eucgr.G02214.1	0,04	42,30	20,44	13,24	1,86	No hits found
Eucgr.G02673.1	0,51	13,39	10,33	7,14	3,98	AT4G03320.1 Symbols: tic20-IV translocon at the inner envelope membrane of chloroplasts 20-IV
Eucgr.A01369.1	5,88	19,37	15,11	12,88	16,92	AT1G04130.1 Symbols: TPR2, AtTPR2 Tetratricopeptide repeat (TPR)-like superfamily protein
Eucgr.B02930.1	0,16	1,00	1,20	1,11	1,20	AT3G47570.1 Symbols: Leucine-rich repeat protein kinase family protein
Eucgr.I00418.1	21,92	0,86	1,80	2,35	7,81	AT5G10150.1 Symbols: Domain of unknown function (DUF966) chr5:3182176-3183834 REVERSE LENGTH=414
Eucgr.G01240.1	5,22	51,15	62,63	47,18	27,12	AT3G18990.1 Symbols: VRN1, REM39 AP2/B3-like transcriptional factor family protein
Eucgr.F00083.2	5,75	30,30	23,10	18,30	15,70	AT1G27840.1 Symbols: ATCSA-1 Transducin/WD40 repeat-like superfamily protein
Eucgr.H02860.1	275,17	0,98	4,25	5,40	27,03	AT1G04680.1 Symbols: Pectin lyase-like superfamily protein chr1:1304052-1307780 REVERSE LENGTH=431
Eucgr.G02722.2	6,40	71,75	59,67	48,41	39,53	AT5G04520.1 Symbols: Protein of unknown function DUF455 chr5:1290012-1291063 REVERSE LENGTH=291
Eucgr.I00519.1	31,40	3,85	4,68	5,80	10,85	AT4G39980.1 Symbols: DHS1 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase 1
Eucgr.H00997.1	0,81	37,45	29,83	17,09	7,06	AT4G01970.1 Symbols: AtSTS, STS stachyose synthase chr4:854073-856953 REVERSE LENGTH=876
Eucgr.G02747.1	26,28	0,69	1,15	1,75	11,16	AT5G22930.1 Symbols: Protein of unknown function (DUF1635) chr5:7668229-7669315 REVERSE LENGTH=238
Eucgr.H04537.3	25,06	97,97	79,89	63,75	86,77	AT1G09140.2 Symbols: ATSRP30, SR30, At-SR30 SERINE-ARGININE PROTEIN 30
Eucgr.E03920.1	0,60	180,55	135,44	120,35	24,55	AT5G47550.1 Symbols: Cystatin/monellin superfamily protein chr5:19286596-19286964 REVERSE LENGTH=122
Eucgr.F02056.1	38,09	1,81	3,39	5,42	14,84	AT3G13130.1 Symbols: unknown protein;
Eucgr.A02615.1	0,54	0,00	0,02	0,00	0,00	AT3G51680.1 Symbols: NAD(P)-binding Rossmann-fold superfamily protein

Eucgr.F02145.1	0,45	20,73	14,00	6,64	6,80	AT4G24540.1 Symbols: AGL24 AGAMOUS-like 24 chr4:12671160-12673645 REVERSE LENGTH=220
Eucgr.G02938.1	48,93	2,52	5,91	9,39	13,24	AT2G39700.1 Symbols: ATEXPA4, ATEXP4, ATHEXP ALPHA 1.6, EXPA4 expansin A4
Eucgr.B02899.1	0,95	84,27	63,53	63,57	25,90	AT1G71000.1 Symbols: Chaperone DnaJ-domain superfamily protein chr1:26769336-26770111 REVERSE LENGTH=165
Eucgr.F03086.1	0,15	16,07	14,52	9,89	5,43	AT4G26270.1 Symbols: PFK3 phosphofructokinase 3 chr4:13301094-13304030 REVERSE LENGTH=489
Eucgr.H03385.1	6,95	375,36	306,61	226,60	105,02	AT5G50850.1 Symbols: MAB1 Transketolase family protein chr5:20689671-20692976 FORWARD LENGTH=363
Eucgr.L03245.1	0,07	190,81	124,71	44,86	2,79	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.G02490.2	60,32	11,22	6,20	4,76	8,48	AT1G69530.2 Symbols: ATEXPA1, EXP1, AT-EXP1, ATEXP1, ATHEXP ALPHA 1.2, EXPA1 expansin A1
Eucgr.H03770.1	4,62	31,62	25,84	22,19	15,19	AT4G15563.1 Symbols: unknown protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.D00485.1	0,00	0,82	0,69	0,20	0,74	AT2G40330.1 Symbols: PYL6, RCAR9 PYR1-like 6 chr2:16845177-16845824 REVERSE LENGTH=215
Eucgr.H02878.1	4,80	12,55	20,94	14,15	23,50	AT4G13550.1 Symbols: triglyceride lipases;triglyceride lipases chr4:7871251-7876160 REVERSE LENGTH=715
Eucgr.A01422.1	0,37	7,57	8,56	4,92	4,02	AT3G49610.1 Symbols: Domain of unknown function (DUF313) chr3:18390413-18391417 REVERSE LENGTH=334
Eucgr.F01661.1	0,15	2,82	2,16	2,39	1,33	AT4G34860.2 Symbols: Plant neutral invertase family protein chr4:16609955-16612019 REVERSE LENGTH=571
Eucgr.K02259.1	1,54	105,03	84,24	56,95	31,85	AT1G18070.2 Symbols: Translation elongation factor EF1A/initiation factor IF2gamma family protein
Eucgr.H02522.1	19,02	1,51	3,06	4,33	8,82	AT1G10750.1 Symbols: Protein of Unknown Function (DUF239) chr1:3574986-3578310 REVERSE LENGTH=467
Eucgr.I01660.1	7,51	15,20	18,90	17,21	19,46	AT3G01810.3 Symbols: unknown protein
Eucgr.I02316.4	7,86	33,41	27,76	28,59	19,16	AT5G66090.1 Symbols: unknown protein
Eucgr.B01913.1	20,02	60,50	48,86	44,37	52,10	AT3G05510.1 Symbols: Phospholipid/glycerol acyltransferase family protein
Eucgr.J01009.1	0,02	3,61	3,78	2,94	0,61	AT5G65550.1 Symbols: UDP-Glycosyltransferase superfamily protein chr5:26198410-26199810 REVERSE LENGTH=466
Eucgr.F02235.1	1,59	18,44	16,51	16,32	11,55	AT5G48570.1 Symbols: ROF2, ATFKBP65, FKBP65 FKBP-type peptidyl-prolyl cis-trans isomerase family protein
Eucgr.E00235.1	28,49	88,82	70,95	60,97	69,20	AT1G32260.1 Symbols: unknown protein
Eucgr.I01727.1	6,85	0,05	0,16	0,39	1,20	AT5G14450.1 Symbols: GDSL-like Lipase/Acylhydrolase superfamily protein
Eucgr.K00423.1	11,57	550,59	437,13	391,96	619,70	AT4G27670.1 Symbols: HSP21 heat shock protein 21 chr4:13819048-13819895 REVERSE LENGTH=227
Eucgr.A02367.1	10,37	49,70	36,95	46,72	33,54	AT1G01170.2 Symbols: Protein of unknown function (DUF1138) chr1:74105-74443 REVERSE LENGTH=83
Eucgr.L00540.1	0,55	0,00	0,02	0,15	0,03	AT3G51970.1 Symbols: ATSAT1, ASAT1, ATASAT1 acyl-CoA sterol acyl transferase 1
Eucgr.K02328.1	1,09	45,53	43,31	28,73	7,18	AT1G73390.3 Symbols: Endosomal targeting BRO1-like domain-containing protein
Eucgr.A00213.1	25,93	2,93	2,89	3,02	7,77	AT1G05805.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.K01196.1	37,71	2,02	3,50	5,81	14,19	AT1G78570.1 Symbols: RHM1, ROL1, ATRHM1 rhamnose biosynthesis 1
Eucgr.J00195.1	0,78	96,37	68,50	50,25	7,59	AT2G40170.1 Symbols: ATEM6, GEAM6, EM6 Stress induced protein chr2:16779792-16780167 REVERSE LENGTH=92
Eucgr.B03508.1	6,82	111,74	86,11	70,89	44,44	AT1G27300.1 Symbols: unknown protein

Eucgr.A02862.1	0,35	0,00	0,02	0,12	0,03	AT2G28100.1 Symbols: ATFUC1, FUC1 alpha-L-fucosidase 1 chr2:11974803-11976489 FORWARD LENGTH=506
Eucgr.H01377.1	0,46	65,76	46,42	35,69	12,76	AT5G36100.1 Symbols: unknown protein
Eucgr.F03878.1	2,58	218,30	167,32	133,63	46,63	AT1G80090.1 Symbols: Cystathionine beta-synthase (CBS) family protein
Eucgr.L03290.1	5,05	0,09	0,15	0,21	0,81	AT2G05920.1 Symbols: Subtilase family protein chr2:2269831-2272207 REVERSE LENGTH=754
Eucgr.I01700.1	25,91	0,69	1,52	3,71	6,37	AT3G27350.2 Symbols: unknown protein
Eucgr.I00668.1	3,32	51,85	37,57	37,58	23,88	AT2G14520.1 Symbols: CBS domain-containing protein with a domain of unknown function (DUF21)
Eucgr.C00873.1	1,92	24,38	18,82	14,75	7,94	AT2G26150.1 Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2
Eucgr.H04478.1	0,03	13,41	9,75	7,19	0,81	AT5G03250.1 Symbols: Phototropic-responsive NPH3 family protein chr5:774591-776855 FORWARD LENGTH=592
Eucgr.C04178.1	3,48	0,13	0,31	0,61	0,48	AT2G26700.1 Symbols: PID2 AGC (cAMP-dependent, cGMP-dependent and protein kinase C) kinase family protein
Eucgr.E00016.1	53,49	0,71	2,00	4,53	14,01	AT3G61920.1 Symbols: unknown protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.I02204.1	3,39	12,91	15,79	15,47	10,98	AT2G33620.4 Symbols: AT hook motif DNA-binding family protein chr2:14234749-14236563 FORWARD LENGTH=351
Eucgr.G01537.1	0,42	0,00	0,00	0,00	0,00	AT5G42650.1 Symbols: AOS, CYP74A, DDE2 allene oxide synthase chr5:17097803-17099359 REVERSE LENGTH=518
Eucgr.E00433.1	3,35	295,94	205,55	185,03	165,89	AT5G51440.1 Symbols: HSP20-like chaperones superfamily protein chr5:20891242-20892013 FORWARD LENGTH=210
Eucgr.C01715.1	3,32	63,28	54,72	35,80	28,00	AT5G20280.1 Symbols: ATSPS1F, SPS1F sucrose phosphate synthase 1F chr5:6844994-6849997 REVERSE LENGTH=1043
Eucgr.C03454.1	0,92	10,40	10,94	6,71	5,80	AT2G26150.1 Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2
Eucgr.A02804.1	24,80	78,79	76,07	72,67	89,08	AT5G62020.1 Symbols: AT-HSFB2A, HSFB2A heat shock transcription factor B2A
Eucgr.H00812.1	38,28	1,20	2,33	3,34	8,92	AT3G24660.1 Symbols: TMKL1 transmembrane kinase-like 1 chr3:9003641-9005751 FORWARD LENGTH=674
Eucgr.H02278.1	2,17	60,73	55,26	50,88	24,61	AT2G23110.2 Symbols: Late embryogenesis abundant protein, group 6 chr2:9840552-9840830 FORWARD LENGTH=92
Eucgr.D02134.1	0,50	0,00	0,01	0,00	0,05	AT3G10600.1 Symbols: CAT7 cationic amino acid transporter 7 chr3:3313969-3317162 FORWARD LENGTH=584
Eucgr.A02658.1	11,73	0,29	0,65	1,52	1,91	AT5G03870.1 Symbols: Glutaredoxin family protein chr5:1035813-1036967 REVERSE LENGTH=384
Eucgr.C01043.1	2,92	122,26	116,87	96,98	13,96	AT3G24520.1 Symbols: AT-HSFC1, HSFC1 heat shock transcription factor C1
Eucgr.E00444.1	25,26	3,62	4,88	6,07	11,61	AT4G17260.1 Symbols: Lactate/malate dehydrogenase family protein chr4:9674057-9675309 FORWARD LENGTH=353
Eucgr.H04092.1	28,01	0,78	2,01	2,33	10,01	AT1G48100.1 Symbols: Pectin lyase-like superfamily protein chr1:17766921-17769625 FORWARD LENGTH=475
Eucgr.F03621.1	0,60	53,74	50,05	47,35	6,91	AT1G21000.2 Symbols: PLATZ transcription factor family protein chr1:7338013-7339088 FORWARD LENGTH=243
Eucgr.I02278.1	0,18	20,21	23,14	33,84	2,38	AT4G35710.1 Symbols: Arabidopsis protein of unknown function (DUF241) chr4:16925301-16926152 FORWARD LENGTH=283
Eucgr.F02505.1	14,04	0,07	0,31	0,42	5,27	AT1G68360.1 Symbols: C2H2 and C2HC zinc fingers superfamily protein chr1:25621678-25622412 REVERSE LENGTH=244
Eucgr.E02655.1	0,52	0,00	0,00	0,00	0,05	AT2G34060.1 Symbols: Peroxidase superfamily protein chr2:14384914-14386530 FORWARD LENGTH=346
Eucgr.I01669.1	3,07	67,66	79,13	75,99	34,25	AT3G27250.1 Symbols: unknown protein
Eucgr.D00295.1	29,29	246,26	190,72	214,06	154,80	AT1G30070.1 Symbols: SGS domain-containing protein chr1:10546794-10548072 REVERSE LENGTH=222

Eucgr.E00015.2	22,81	1,66	2,21	3,70	7,74	No hits found
Eucgr.B02917.1	7,20	0,00	0,18	1,03	0,59	AT4G14060.1 Symbols: Polyketide cyclase/dehydrase and lipid transport superfamily protein
Eucgr.H03952.1	0,45	0,00	0,01	0,00	0,05	AT5G49130.1 Symbols: MATE efflux family protein chr5:19915904-19917525 FORWARD LENGTH=502
Eucgr.L00240.1	0,02	34,50	22,77	10,27	0,66	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.L03270.1	0,42	742,81	379,17	232,66	3,62	AT4G33467.2 Symbols: unknown protein
Eucgr.K01431.1	1,67	34,99	28,93	27,47	15,68	AT1G04520.1 Symbols: PDLP2 plasmodesmata-located protein 2 chr1:1231852-1233787 FORWARD LENGTH=307
Eucgr.H00798.1	58,69	1,20	4,03	6,42	24,86	AT5G48740.1 Symbols: Leucine-rich repeat protein kinase family protein chr5:19765324-19769314 REVERSE LENGTH=895
Eucgr.D01780.1	7,62	1,87	1,71	1,79	2,13	AT4G23720.1 Symbols: Protein of unknown function (DUF1191) chr4:12358707-12359648 FORWARD LENGTH=313
Eucgr.J02951.1	0,48	0,00	0,00	0,02	0,03	AT2G26490.1 Symbols: Transducin/WD40 repeat-like superfamily protein
Eucgr.D00534.1	0,00	0,90	1,04	0,71	0,29	AT5G45910.1 Symbols: GDSL-like Lipase/Acylhydrolase superfamily protein
Eucgr.E00347.1	3,00	226,36	192,13	201,85	132,87	AT4G33550.1 Symbols: Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.D01202.1	29,79	1,29	1,68	2,32	15,49	AT2G44690.1 Symbols: ARAC9, ATROP8, ROP8 Arabidopsis RAC-like 9 chr2:18429276-18430636 FORWARD LENGTH=209
Eucgr.A02926.1	0,06	4,08	5,38	5,33	1,12	AT5G07490.1 Symbols: unknown protein;
Eucgr.F03665.1	0,73	0,00	0,01	0,02	0,04	AT5G59810.1 Symbols: ATSBT5.4, SBT5.4 Subtilase family protein chr5:24096895-24100387 REVERSE LENGTH=778
Eucgr.L00251.1	0,21	325,56	149,21	61,46	3,38	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.C02145.1	7,05	63,91	50,64	32,15	20,87	AT4G31140.1 Symbols: O-Glycosyl hydrolases family 17 protein chr4:15141581-15143188 FORWARD LENGTH=484
Eucgr.C00112.1	0,70	12,29	14,05	17,42	3,57	AT5G57180.2 Symbols: CIA2 chloroplast import apparatus 2 chr5:23168393-23170763 FORWARD LENGTH=435
Eucgr.B04029.1	1,08	5,43	5,54	4,24	4,75	No hits found
Eucgr.K01574.1	2,12	14,23	17,44	12,16	9,54	AT2G41510.1 Symbols: ATCKX1, CKX1 cytokinin oxidase/dehydrogenase 1
Eucgr.J01011.1	24,97	0,49	1,02	2,06	5,93	AT5G04660.1 Symbols: CYP77A4 cytochrome P450, family 77, subfamily A, polypeptide 4
Eucgr.B01852.1	31,04	2,22	3,00	3,56	11,55	AT1G09070.1 Symbols: SRC2, (AT)SRC2 soybean gene regulated by cold-2 chr1:2927767-2928741 FORWARD LENGTH=324
Eucgr.D01381.1	17,58	1,64	1,70	1,92	4,85	AT4G05520.1 Symbols: ATEHD2, EHD2 EPS15 homology domain 2 chr4:2804522-2807833 FORWARD LENGTH=546
Eucgr.K02245.1	9,42	0,04	0,21	0,18	2,23	AT3G18180.1 Symbols: Glycosyltransferase family 61 protein chr3:6230270-6231878 FORWARD LENGTH=470
Eucgr.C02293.1	2,07	49,90	58,91	51,95	23,88	AT5G57480.1 Symbols: P-loop containing nucleoside triphosphate hydrolases superfamily protein
Eucgr.G01235.1	3,36	40,19	40,09	43,38	22,26	AT3G18990.1 Symbols: VRN1, REM39 AP2/B3-like transcriptional factor family protein
Eucgr.H02498.1	17,77	195,87	178,28	168,88	130,52	AT1G59960.1 Symbols: NAD(P)-linked oxidoreductase superfamily protein
Eucgr.K01344.1	5,46	109,37	132,11	131,68	77,29	AT3G12750.1 Symbols: ZIP1 zinc transporter 1 precursor chr3:4051950-4053156 REVERSE LENGTH=355
Eucgr.A00215.1	35,20	7,54	7,19	9,01	17,55	AT2G43130.1 Symbols: ARA4, ATRAB11F, ATRABA5C, ARA-4, RABA5C
Eucgr.H02273.1	0,11	17,75	13,64	12,12	2,51	AT2G23120.1 Symbols: Late embryogenesis abundant protein, group 6 chr2:9842102-9842353 FORWARD LENGTH=83

Eucgr.E01133.2	5,10	36,15	34,71	36,12	27,09	AT4G10430.3 Symbols: TMPIT-like protein chr4:6455110-6457996 REVERSE LENGTH=347
Eucgr.K02872.1	0,00	1,89	2,16	1,86	0,66	No hits found
Eucgr.G02197.1	8,32	23,70	36,64	34,02	25,13	AT1G23010.1 Symbols: LPR1 Cupredoxin superfamily protein chr1:8147353-8149580 FORWARD LENGTH=581
Eucgr.A02416.1	24,03	0,12	0,23	0,43	5,41	AT3G57830.1 Symbols: Leucine-rich repeat protein kinase family protein
Eucgr.I01670.1	5,31	43,82	33,92	30,65	22,35	AT3G48510.1 Symbols: unknown protein
Eucgr.B03684.1	49,42	9,83	22,21	38,16	22,44	AT5G12870.1 Symbols: ATMYB46, MYB46 myb domain protein 46 chr5:4062939-4064939 REVERSE LENGTH=280
Eucgr.K01425.1	10,37	0,33	0,93	1,13	5,35	AT4G28650.1 Symbols: Leucine-rich repeat transmembrane protein kinase family protein
Eucgr.K03318.1	49,40	3,62	3,07	4,48	16,55	AT3G54560.1 Symbols: HTA11 histone H2A 11 chr3:20196532-20197466 FORWARD LENGTH=136
Eucgr.K00150.1	45,43	2,66	4,30	5,27	10,07	AT5G51550.1 Symbols: EXL3 EXORDIUM like 3 chr5:20939793-20940806 REVERSE LENGTH=337
Eucgr.I02678.1	6,78	34,08	31,69	27,68	25,32	AT4G33500.1 Symbols: Protein phosphatase 2C family protein chr4:16112835-16116243 REVERSE LENGTH=724
Eucgr.C03434.1	0,31	6,75	6,22	6,19	2,03	AT2G26150.1 Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2
Eucgr.G02156.1	40,72	3,42	3,65	5,44	13,75	AT1G14890.1 Symbols: Plant invertase/pectin methylesterase inhibitor superfamily protein
Eucgr.K03526.1	15,24	174,63	154,97	121,95	59,79	AT4G02120.1 Symbols: CTP synthase family protein chr4:940873-944097 FORWARD LENGTH=556
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.A02229.1	34,74	0,51	1,19	1,73	4,71	AT2G01420.1 Symbols: PIN4, ATPIN4 Auxin efflux carrier family protein chr2:180478-183199 REVERSE LENGTH=612
Eucgr.F03514.1	189,06	3,00	6,50	5,98	31,95	AT5G55410.2 Symbols: Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein
Eucgr.A02008.1	0,16	1,07	1,89	1,66	0,83	AT1G06490.1 Symbols: ATGSL07, gsl07, atgsl7, GSL7 glucan synthase-like 7
Eucgr.B00580.1	7,83	0,48	0,83	1,39	1,06	AT5G07990.1 Symbols: TT7, CYP75B1, D501 Cytochrome P450 superfamily protein
Eucgr.E00360.1	5,58	623,57	416,31	243,00	88,39	AT4G17030.1 Symbols: ATEXLB1, EXPR, AT-EXPR, ATEXPR1, ATHEXP BETA 3.1, EXLB1 expansin-like B1
Eucgr.G03118.1	5,52	47,21	44,62	43,13	33,32	No hits found
Eucgr.K02961.3	16,85	174,50	121,56	94,41	87,15	AT2G40300.1 Symbols: ATFER4, FER4 ferritin 4 chr2:16831501-16833214 REVERSE LENGTH=259
Eucgr.B00412.1	30,69	0,47	0,95	1,72	12,73	AT5G62710.1 Symbols: Leucine-rich repeat protein kinase family protein chr5:25187438-25190325 FORWARD LENGTH=604
Eucgr.B03008.1	62,99	475,02	399,93	377,86	240,60	AT1G68300.1 Symbols: Adenine nucleotide alpha hydrolases-like superfamily protein chr1:25598518-25599261 REVERSE LENGTH=160
Eucgr.G01728.1	3,87	0,06	0,25	0,35	1,00	AT4G39010.1 Symbols: AtGH9B18, GH9B18 glycosyl hydrolase 9B18 chr4:18176162-18179102 REVERSE LENGTH=497
Eucgr.J00457.2	5,69	47,20	42,83	33,08	18,78	AT5G22510.1 Symbols: INV-E, At-A/N-InvE alkaline/neutral invertase chr5:7474974-7477479 REVERSE LENGTH=617
Eucgr.C00218.1	61,73	8,50	11,31	12,67	24,16	AT2G23810.1 Symbols: TET8 tetraspanin8 chr2:10135859-10137352 REVERSE LENGTH=273
Eucgr.F00301.1	28,54	4,88	5,57	7,81	14,63	AT1G78110.1 Symbols: unknown protein
Eucgr.K01055.1	42,72	337,71	362,68	582,22	200,35	No hits found
Eucgr.G02495.1	16,31	0,79	1,91	3,33	7,13	AT1G14190.1 Symbols: Glucose-methanol-choline (GMC) oxidoreductase family protein
Eucgr.H04815.2	8,01	133,50	102,82	79,74	77,03	AT2G22240.1 Symbols: ATMIPS2, MIPS2, ATIPS2 myo-inositol-1-phosphate synthase 2

Eucgr.B00170.1	1,13	0,22	0,00	0,07	0,09	No hits found
Eucgr.A02150.1	1,06	9,41	6,93	3,52	3,95	AT1G69560.1 Symbols: MYB105, LOF2, ATMYB105 myb domain protein 105
Eucgr.K01583.1	3,82	0,03	0,03	0,09	1,02	AT1G17020.1 Symbols: SRG1, ATSRG1 senescence-related gene 1 chr1:5820258-5821741 FORWARD LENGTH=358
Eucgr.K00778.1	2,61	0,06	0,26	0,71	0,60	AT2G32540.1 Symbols: ATCSLB04, CSLB04, ATCSLB4 cellulose synthase-like B4
Eucgr.I02514.1	3,73	0,20	0,33	0,38	0,53	AT4G17000.1 Symbols: unknown protein;
Eucgr.K01967.1	26,89	1,99	3,68	4,77	10,23	AT5G44530.1 Symbols: Subtilase family protein chr5:17937931-17941193 FORWARD LENGTH=840
Eucgr.J00948.1	0,00	2,58	2,03	1,20	0,23	AT2G28490.1 Symbols: RmlC-like cupins superfamily protein chr2:12178812-12180983 REVERSE LENGTH=511
Eucgr.A01877.1	0,00	0,31	0,79	1,75	0,22	AT4G35160.1 Symbols: O-methyltransferase family protein chr4:16730989-16732808 REVERSE LENGTH=382
Eucgr.H00467.1	6,20	79,40	69,57	83,40	24,29	AT1G60710.1 Symbols: ATB2 NAD(P)-linked oxidoreductase superfamily protein
Eucgr.F00649.1	1,21	0,06	0,13	0,17	0,23	AT5G55250.1 Symbols: IAMT1 IAA carboxylmethyltransferase 1 chr5:22407589-22410854 REVERSE LENGTH=386
Eucgr.A02858.1	0,77	45,90	34,68	20,99	8,40	AT5G51760.1 Symbols: AHG1 Protein phosphatase 2C family protein chr5:21026916-21028912 FORWARD LENGTH=416
Eucgr.H04619.1	0,34	29,24	24,75	23,01	2,43	No hits found
Eucgr.K01580.1	3,13	16,92	16,68	21,70	10,45	AT3G57190.1 Symbols: peptide chain release factor, putative chr3:21166468-21168198 FORWARD LENGTH=406
Eucgr.A01798.3	12,42	0,13	0,53	0,87	5,37	AT1G03620.1 Symbols: ELMO/CED-12 family protein chr1:904319-906013 REVERSE LENGTH=265
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.A02601.1	16,96	1,46	1,64	1,18	2,97	AT3G52500.1 Symbols: Eukaryotic aspartyl protease family protein chr3:19465644-19467053 REVERSE LENGTH=469
Eucgr.F01151.1	22,41	211,06	158,42	117,64	60,94	AT4G27520.1 Symbols: ENODL2, AtENODL2 early nodulin-like protein 2 chr4:13750668-13751819 REVERSE LENGTH=349
Eucgr.F02851.1	192,64	1,06	4,02	7,93	38,91	AT2G30810.1 Symbols: Gibberellin-regulated family protein chr2:13127945-13128630 REVERSE LENGTH=106
Eucgr.H02896.1	0,07	6,57	8,20	6,34	1,14	AT3G24310.1 Symbols: MYB305, ATMYB71 myb domain protein 305 chr3:8811336-8812330 REVERSE LENGTH=269
Eucgr.H02593.1	13,14	204,27	152,30	118,96	62,04	AT1G60420.1 Symbols: DC1 domain-containing protein chr1:22261978-22264243 FORWARD LENGTH=578
Eucgr.F02468.1	0,67	0,02	0,07	0,12	0,13	AT5G45480.1 Symbols: Protein of unknown function (DUF594) chr5:18426296-18428929 REVERSE LENGTH=877
Eucgr.B03303.1	43,12	3,33	6,88	9,78	19,35	AT1G10750.1 Symbols: Protein of Unknown Function (DUF239) chr1:3574986-3578310 REVERSE LENGTH=467
Eucgr.F03761.1	2,71	0,00	0,00	0,05	0,53	AT5G08640.2 Symbols: FLS1 flavonol synthase 1 chr5:2804009-2805175 FORWARD LENGTH=336
Eucgr.G00317.1	5,18	0,46	0,32	0,81	1,65	AT1G09812.1 Symbols: unknown protein
Eucgr.H02966.1	24,69	0,87	1,27	1,73	3,30	No hits found
Eucgr.I01646.1	48,55	1,72	2,40	2,82	14,36	AT3G27060.1 Symbols: TSO2, ATTSO2 Ferritin/ribonucleotide reductase-like family protein
Eucgr.J01089.1	28,49	0,44	2,27	6,81	14,22	AT2G39220.1 Symbols: PLP6, PLA IIB PATATIN-like protein 6 chr2:16375055-16376663 REVERSE LENGTH=499
Eucgr.B00827.1	6,35	0,57	2,11	5,48	2,27	No hits found
Eucgr.K01573.1	0,25	15,37	15,23	13,35	5,95	No hits found
Eucgr.G01350.2	23,73	146,29	126,30	144,55	62,96	AT4G34230.1 Symbols: CAD5, ATCAD5, CAD-5 cinnamyl alcohol dehydrogenase 5

Eucgr.F02055.1	2,82	0,21	0,18	0,43	0,66	AT3G13130.1 Symbols: unknown protein
Eucgr.E01110.2	4,34	59,21	49,17	51,45	27,12	AT1G72680.1 Symbols: ATCAD1, CAD1 cinnamyl-alcohol dehydrogenase chr1:27359346-27360876 REVERSE LENGTH=355
Eucgr.I01013.1	8,29	0,08	0,65	1,64	1,01	AT2G23610.1 Symbols: ATME53, ME53 methyl esterase 3 chr2:10044410-10046403 REVERSE LENGTH=263
Eucgr.A02860.1	4,54	0,07	0,11	0,14	0,99	AT2G16280.1 Symbols: KCS9 3-ketoacyl-CoA synthase 9 chr2:7051186-7052724 FORWARD LENGTH=512
Eucgr.K01314.2	2,51	222,00	178,21	124,80	132,76	AT3G08590.2 Symbols: Phosphoglycerate mutase, 2,3-bisphosphoglycerate-independent
Eucgr.F04059.1	20,67	2,20	3,84	5,47	10,52	AT3G13000.2 Symbols: Protein of unknown function, DUF547 chr3:4158214-4160989 REVERSE LENGTH=582
Eucgr.H03957.1	26,11	0,69	1,46	3,15	5,85	AT4G02340.1 Symbols: alpha/beta-Hydrolases superfamily protein chr4:1035722-1037403 FORWARD LENGTH=324
Eucgr.I01790.1	1,06	11,78	13,06	15,94	15,77	AT3G01500.2 Symbols: CA1 carbonic anhydrase 1 chr3:194853-197873 REVERSE LENGTH=347
Eucgr.G03283.1	6,98	23,73	17,15	14,04	15,53	AT3G60910.1 Symbols: S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
Eucgr.H03517.1	16,45	0,37	0,95	1,84	2,74	AT1G05370.1 Symbols: Sec14p-like phosphatidylinositol transfer family protein
Eucgr.F03754.1	12,93	0,24	0,68	1,63	5,89	AT1G38131.1 Symbols: O-fucosyltransferase family protein chr1:14293392-14296020 REVERSE LENGTH=589
Eucgr.C00378.1	2,87	0,02	0,08	0,16	0,68	AT2G24280.1 Symbols: alpha/beta-Hydrolases superfamily protein chr2:10334622-10336927 FORWARD LENGTH=494
Eucgr.E00847.1	32,93	1,08	1,02	1,82	4,93	AT4G23800.1 Symbols: HMG (high mobility group) box protein chr4:12390228-12392516 FORWARD LENGTH=456
Eucgr.E00078.3	18,23	242,77	193,91	127,40	69,04	AT2G46270.1 Symbols: GBF3 G-box binding factor 3 chr2:19000859-19002901 FORWARD LENGTH=382
Eucgr.I01891.1	13,34	0,50	1,23	2,02	3,26	AT3G28420.1 Symbols: Putative membrane lipoprotein chr3:10654674-10655324 REVERSE LENGTH=216
Eucgr.D01765.2	0,07	3,26	6,28	7,07	1,24	AT4G23990.1 Symbols: ATCSLG3, CSLG3 cellulose synthase like G3 chr4:12456491-12460498 FORWARD LENGTH=751
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.I00409.1	0,06	3,27	4,18	2,91	0,60	AT4G34131.1 Symbols: UGT73B3 UDP-glucosyl transferase 73B3 chr4:16343268-16344713 REVERSE LENGTH=481
Eucgr.G03282.1	0,09	10,41	8,85	6,96	1,53	AT2G28420.1 Symbols: Lactoylglutathione lyase / glyoxalase I family protein
Eucgr.I01627.3	8,50	57,85	46,76	38,03	19,29	AT3G01990.1 Symbols: ACR6 ACT domain repeat 6 chr3:330256-332066 FORWARD LENGTH=433
Eucgr.J00303.1	7,16	0,14	0,37	0,57	0,98	AT5G06470.1 Symbols: Glutaredoxin family protein chr5:1974659-1975378 REVERSE LENGTH=239
Eucgr.C04221.1	1,01	0,00	0,02	0,00	0,11	AT1G15360.1 Symbols: SHN1, WIN1 Integrase-type DNA-binding superfamily protein
Eucgr.G00501.1	0,71	0,03	0,02	0,07	0,09	AT5G42650.1 Symbols: AOS, CYP74A, DDE2 allene oxide synthase chr5:17097803-17099359 REVERSE LENGTH=518
Eucgr.C00666.1	5,19	63,12	75,80	71,30	30,32	AT2G26170.2 Symbols: CYP711A1 cytochrome P450, family 711, subfamily A, polypeptide 1
Eucgr.L02188.1	12,05	0,54	1,77	3,57	5,43	AT5G11000.1 Symbols: Plant protein of unknown function (DUF868) chr5:3479166-3480335 REVERSE LENGTH=389
Eucgr.B00302.1	27,47	0,98	1,66	2,43	6,41	AT2G47160.1 Symbols: BOR1 HCO3- transporter family chr2:19357740-19360787 REVERSE LENGTH=704
Eucgr.H01095.1	78,97	0,57	3,99	6,76	19,38	AT4G02290.1 Symbols: AtGH9B13, GH9B13 glycosyl hydrolase 9B13 chr4:1002654-1005125 REVERSE LENGTH=516
Eucgr.G03200.1	156,94	11,37	7,49	12,01	15,60	AT5G07050.1 Symbols: nodulin MtN21 /EamA-like transporter family protein
Eucgr.K03469.1	56,95	1,15	2,72	3,54	10,39	AT1G08560.1 Symbols: SYP111, KN, ATSYP111 syntaxin of plants 111 chr1:2709778-2710710 REVERSE LENGTH=310
Eucgr.E00812.1	8,79	0,24	0,87	2,51	2,99	AT1G64160.1 Symbols: Disease resistance-responsive (dirigent-like protein) family protein

Eucgr.C01388.1	79,86	0,45	1,28	1,19	21,24	AT2G10940.2 Symbols: Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein
Eucgr.J01066.1	37,91	812,10	857,97	610,25	159,86	AT2G39130.1 Symbols: Transmembrane amino acid transporter family protein
Eucgr.H02629.1	20,89	306,08	289,19	273,68	273,55	AT5G48570.1 Symbols: ROF2, ATKBP65, FKBP65 FKBP-type peptidyl-prolyl cis-trans isomerase family protein
Eucgr.B01279.1	10,18	0,08	0,22	0,34	5,20	AT3G26700.1 Symbols: Protein kinase superfamily protein chr3:9810669-9812356 FORWARD LENGTH=380
Eucgr.K02082.1	161,54	0,89	2,95	5,95	52,84	AT2G36570.1 Symbols: Leucine-rich repeat protein kinase family protein chr2:15335583-15337725 FORWARD LENGTH=672
Eucgr.A02790.1	15,66	1398,81	1140,99	647,25	272,46	AT5G52300.2 Symbols: RD29B, LTI65 CAP160 protein chr5:21237205-21239404 FORWARD LENGTH=618
Eucgr.E00266.1	34,34	2,38	6,52	11,42	13,52	No hits found
Eucgr.F02525.1	20,40	0,00	0,29	0,67	1,40	No hits found
Eucgr.F02150.1	9,89	0,25	0,43	0,60	1,79	AT5G60930.1 Symbols: P-loop containing nucleoside triphosphate hydrolases superfamily protein
Eucgr.H01202.1	12,87	0,05	0,33	2,39	1,37	AT2G47880.1 Symbols: Glutaredoxin family protein chr2:19605124-19605432 FORWARD LENGTH=102
Eucgr.I01639.1	11,06	69,22	57,03	66,79	28,53	AT5G40960.1 Symbols: Protein of unknown function (DUF 3339) chr5:16412745-16412951 REVERSE LENGTH=68
Eucgr.A02352.1	1,29	0,05	0,06	0,09	0,16	AT2G41790.1 Symbols: Insulinase (Peptidase family M16) family protein chr2:17429453-17436110 REVERSE LENGTH=970
Eucgr.K03244.2	29,50	382,43	351,93	254,16	266,20	AT3G53420.2 Symbols: PIP2A, PIP2, PIP2;1 plasma membrane intrinsic protein 2A
Eucgr.E00450.1	16,88	2,16	2,89	4,73	7,62	AT1G62520.1 Symbols: unknown protein
Eucgr.K01030.1	10,57	40,54	48,26	41,08	28,52	AT4G03560.1 Symbols: ATPC1, TPC1, ATCCH1, FOU2 two-pore channel 1 chr4:1580253-1585691 FORWARD LENGTH=733
Eucgr.B02566.1	2,55	53,48	47,33	37,84	27,61	AT1G24130.1 Symbols: Transducin/WD40 repeat-like superfamily protein chr1:8534183-8535430 REVERSE LENGTH=415
Eucgr.K02050.1	309,22	1,77	6,92	12,73	116,05	AT3G10080.1 Symbols: RmlC-like cupins superfamily protein chr3:3107476-3108159 REVERSE LENGTH=227
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.L02059.1	0,80	0,00	0,01	0,03	0,12	AT3G63470.1 Symbols: scpl40 serine carboxypeptidase-like 40 chr3:23438561-23440862 FORWARD LENGTH=502
Eucgr.H04021.1	5,29	0,08	0,17	0,19	0,34	AT4G16120.1 Symbols: ATSEB1, COBL7, SEB1 COBRA-like protein-7 precursor
Eucgr.I00918.1	1,14	0,08	0,07	0,30	0,19	AT4G36950.1 Symbols: MAPKKK21 mitogen-activated protein kinase kinase kinase 21
Eucgr.F02209.1	11,50	101,96	92,75	54,42	38,58	AT1G58520.1 Symbols: RXW8 lipases;hydrolases, acting on ester bonds
Eucgr.F03235.1	12,40	0,03	0,40	0,61	3,00	AT1G29050.1 Symbols: TBL38 TRICHOME BIREFRINGENCE-LIKE 38 chr1:10136376-10139082 REVERSE LENGTH=380
Eucgr.H00766.1	0,37	32,50	24,18	11,54	4,89	AT3G07250.1 Symbols: nuclear transport factor 2 (NTF2) family protein / RNA recognition motif (RRM)-containing protein
Eucgr.H04224.1	6,87	0,05	0,08	0,35	0,69	AT5G44040.1 Symbols: unknown protein
Eucgr.B03841.1	71,58	1,99	3,99	6,96	24,37	AT5G22400.1 Symbols: Rho GTPase activating protein with PAK-box/P21-Rho-binding domain
Eucgr.K03042.1	15,22	268,45	224,70	203,87	53,05	AT5G02020.1 Symbols: SIS Encodes a protein involved in salt tolerance, names SIS (Salt Induced Serine rich).
Eucgr.F02646.1	1,99	0,00	0,02	0,06	0,44	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.F04109.1	10,13	734,90	714,88	986,43	876,56	AT4G10250.1 Symbols: ATHSP22.0 HSP20-like chaperones superfamily protein
Eucgr.A00276.1	0,08	2,08	1,68	1,14	1,33	No hits found

Eucgr.B03207.1	3,55	0,14	0,15	0,28	0,55	AT1G26760.1 Symbols: ATXR1, SDG35 SET domain protein 35 chr1:9248304-9249941 REVERSE LENGTH=545
Eucgr.A01770.1	0,00	0,51	0,72	0,38	0,12	AT2G20340.1 Symbols: Pyridoxal phosphate (PLP)-dependent transferases superfamily protein
Eucgr.I01338.1	36,24	0,77	1,26	1,86	11,64	AT4G34760.1 Symbols: SAUR-like auxin-responsive protein family chr4:16582471-16582794 REVERSE LENGTH=107
Eucgr.E00448.1	580,44	10,96	19,88	34,02	221,46	AT4G12510.1 Symbols: Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein
Eucgr.B02529.2	50,46	4,34	8,64	14,31	25,22	AT5G60720.1 Symbols: Protein of unknown function, DUF547 chr5:24418436-24422177 REVERSE LENGTH=691
Eucgr.A01357.2	31,33	1,48	1,95	2,58	3,04	AT5G44040.1 Symbols: unknown protein
Eucgr.E01049.1	0,02	0,74	0,25	0,17	0,21	AT4G00870.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.I02135.1	9,97	0,49	0,58	0,90	3,12	AT1G72430.1 Symbols: SAUR-like auxin-responsive protein family chr1:27265150-27265509 REVERSE LENGTH=119
Eucgr.A01957.1	0,58	0,00	0,03	0,02	0,08	AT2G42800.1 Symbols: ATRLP29, RLP29 receptor like protein 29 chr2:17808157-17809545 REVERSE LENGTH=462
Eucgr.J01883.1	36,93	10,59	10,01	8,15	17,05	AT1G07220.1 Symbols: Arabidopsis thaliana protein of unknown function (DUF821)
Eucgr.J00621.1	21,67	0,30	0,32	0,70	8,21	AT3G43720.2 Symbols: Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein
Eucgr.B03715.1	16,06	0,00	0,00	0,00	1,00	AT1G74670.1 Symbols: Gibberellin-regulated family protein chr1:28053378-28053893 FORWARD LENGTH=101
Eucgr.F02337.4	7,27	113,34	113,21	82,20	37,16	AT1G45249.1 Symbols: ABF2, AREB1, ATAREB1 abscisic acid responsive elements-binding factor 2
Eucgr.J03131.1	7,34	0,58	0,62	0,67	2,26	AT1G79720.1 Symbols: Eukaryotic aspartyl protease family protein chr1:29997259-29998951 REVERSE LENGTH=484
Eucgr.H00807.1	0,85	0,00	0,00	0,00	0,20	AT3G24750.1 Symbols: unknown protein
Eucgr.D00426.1	11,10	0,28	0,39	0,45	1,34	AT5G45700.1 Symbols: Haloacid dehalogenase-like hydrolase (HAD) superfamily protein
Eucgr.C01078.1	16,16	0,25	0,57	0,70	1,61	AT4G13370.1 Symbols: Plant protein of unknown function (DUF936) chr4:7777916-7780334 REVERSE LENGTH=673
Eucgr.F04387.1	2,85	0,40	0,33	0,49	0,87	AT4G08550.1 Symbols: electron carriers;protein disulfide oxidoreductases
Eucgr.B01477.1	21,18	0,60	1,36	2,06	7,10	AT3G63430.1 Symbols: unknown protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H02960.1	122,70	2,60	7,75	12,67	57,78	AT1G04040.1 Symbols: HAD superfamily, subfamily IIIB acid phosphatase chr1:1042564-1043819 REVERSE LENGTH=271
Eucgr.F04160.1	17,17	0,04	0,25	0,26	3,08	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.G02547.1	0,01	1,16	0,81	0,39	0,24	AT1G27040.1 Symbols: Major facilitator superfamily protein chr1:9386893-9390018 REVERSE LENGTH=567
Eucgr.J03029.1	14,52	0,75	1,24	1,51	2,28	AT3G15550.1 Symbols: unknown protein
Eucgr.D00307.4	6,66	36,15	33,90	37,67	19,08	No hits found
Eucgr.I01402.1	289,10	9,62	19,51	27,72	61,09	AT4G34980.1 Symbols: SLP2 subtilisin-like serine protease 2 chr4:16656929-16659223 REVERSE LENGTH=764
Eucgr.J02407.1	74,28	18,89	17,82	19,95	38,28	AT3G51160.1 Symbols: MUR1, MUR_1, GMD2 NAD(P)-binding Rossmann-fold superfamily protein
Eucgr.E04312.2	19,54	0,52	1,00	1,71	5,69	AT4G20050.2 Symbols: QRT3 Pectin lyase-like superfamily protein chr4:10849911-10852090 REVERSE LENGTH=481
Eucgr.H02465.1	4,88	30,26	28,79	18,16	13,32	AT5G14420.2 Symbols: RGLG2 RING domain ligase2 chr5:4648355-4650563 REVERSE LENGTH=468
Eucgr.K01841.1	203,66	9,77	11,72	14,99	51,74	AT3G53750.1 Symbols: ACT3 actin 3 chr3:19915924-19917371 FORWARD LENGTH=377

Eucgr.G03262.1	7,68	0,49	0,53	0,77	0,76	AT3G51930.1 Symbols: Transducin/WD40 repeat-like superfamily protein
Eucgr.J00777.1	4,31	0,15	0,16	0,28	1,60	AT4G12620.1 Symbols: ORC1B, ATORC1B, UNE13 origin of replication complex 1B
Eucgr.D01627.1	90,32	2,93	4,91	8,21	19,08	AT1G30690.2 Symbols: Sec14p-like phosphatidylinositol transfer family protein
Eucgr.I01996.1	77,43	0,73	1,68	3,88	31,92	AT5G15350.1 Symbols: ENODL17, AtENODL17 early nodulin-like protein 17 chr5:4985184-4986154 REVERSE LENGTH=172
Eucgr.I00988.1	287,18	3,87	5,41	6,76	52,68	AT5G66920.1 Symbols: sks17 SKU5 similar 17 chr5:26722963-26725370 FORWARD LENGTH=546
Eucgr.J02867.3	36,68	1,42	2,88	3,69	14,74	AT3G13750.1 Symbols: BGAL1 beta galactosidase 1 chr3:4511192-4515756 FORWARD LENGTH=847
Eucgr.F03122.1	7,77	1,04	1,61	1,76	2,53	AT3G13690.1 Symbols: Protein kinase protein with adenine nucleotide alpha hydrolases-like domain
Eucgr.E02312.1	2,66	9,70	11,64	8,56	8,93	AT4G12020.3 Symbols: WRKY19, ATWRKY19, MAPKKK11, MEKK4 protein kinase family protein
Eucgr.K02088.1	16,98	1,60	2,05	2,25	6,28	AT5G48460.1 Symbols: Actin binding Calponin homology (CH) domain-containing protein
Eucgr.K02296.1	0,77	0,05	0,01	0,05	0,08	AT1G14670.1 Symbols: Endomembrane protein 70 protein family chr1:5037669-5040199 FORWARD LENGTH=592
Eucgr.C02605.1	5,32	0,00	0,19	0,53	0,72	AT2G25060.1 Symbols: ENODL14, AtENODL14 early nodulin-like protein 14
Eucgr.G02332.1	18,19	5,87	7,18	7,76	10,69	AT3G25860.1 Symbols: LTA2, PLE2 2-oxoacid dehydrogenases acyltransferase family protein
Eucgr.G02612.1	12,45	34,63	31,80	25,39	26,65	AT5G23390.1 Symbols: Plant protein of unknown function (DUF639) chr5:7870719-7874283 REVERSE LENGTH=730
Eucgr.H03764.1	26,53	6,32	6,53	8,88	14,82	AT2G45140.1 Symbols: PVA12 plant VAP homolog 12 chr2:18611029-18612971 FORWARD LENGTH=239
Eucgr.H02855.1	1,76	0,19	0,09	0,11	0,41	AT1G61820.1 Symbols: BGLU46 beta glucosidase 46 chr1:22835452-22838444 FORWARD LENGTH=516
Eucgr.F02578.1	20,17	0,22	0,70	0,50	2,27	AT3G15540.1 Symbols: IAA19, MSG2 indole-3-acetic acid inducible 19 chr3:5264100-5265378 FORWARD LENGTH=197
Eucgr.H02676.1	29,26	1,00	2,73	3,81	14,98	AT4G13950.1 Symbols: RALFL31 ralfl-like 31 chr4:8058268-8058609 REVERSE LENGTH=113
Eucgr.H00764.1	3,75	44,29	39,82	29,85	27,21	AT4G31940.1 Symbols: CYP82C4 cytochrome P450, family 82, subfamily C, polypeptide 4
Eucgr.A02888.1	7,36	0,04	0,08	0,08	1,55	AT1G62660.1 Symbols: Glycosyl hydrolases family 32 protein chr1:23199949-23203515 FORWARD LENGTH=648
Eucgr.J00695.1	1,65	0,02	0,04	0,15	0,17	AT5G19410.1 Symbols: ABC-2 type transporter family protein chr5:6545237-6547111 REVERSE LENGTH=624
Eucgr.H00729.1	3,66	84,10	81,01	106,79	20,81	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H04545.1	58,12	477,26	465,70	508,69	234,96	AT1G56220.2 Symbols: Dormancy/auxin associated family protein chr1:21043414-21044381 FORWARD LENGTH=112
Eucgr.A01499.1	0,25	7,54	6,14	3,12	1,56	AT2G36270.1 Symbols: ABI5, GIA1 Basic-leucine zipper (bZIP) transcription factor family protein
Eucgr.D01440.1	52,51	560,58	442,94	396,48	159,63	AT2G35760.1 Symbols: Uncharacterised protein family (UPF0497) chr2:15032059-15033094 FORWARD LENGTH=201
Eucgr.A00686.1	15,75	0,93	1,89	2,82	5,55	AT2G38370.1 Symbols: Plant protein of unknown function (DUF827) chr2:16072184-16074168 REVERSE LENGTH=522
Eucgr.F02993.1	70,48	0,65	1,11	0,99	9,15	AT1G72160.1 Symbols: Sec14p-like phosphatidylinositol transfer family protein
Eucgr.K02228.1	14,81	37,22	37,61	37,69	28,11	AT1G48840.1 Symbols: Plant protein of unknown function (DUF639) chr1:18061931-18064811 FORWARD LENGTH=691
Eucgr.B00987.1	2,83	0,05	0,06	0,37	0,82	AT1G18290.1 Symbols: unknown protein
Eucgr.C00762.2	6,50	1,87	0,87	1,00	2,01	AT5G20850.1 Symbols: ATRAD51, RAD51 RAS associated with diabetes protein 51

Eucgr.C04062.1	12,75	58,82	52,04	37,39	30,18	AT5G35180.2 Symbols: Protein of unknown function (DUF1336) chr5:13424538-13432699 FORWARD LENGTH=778
Eucgr.B00670.1	5,73	0,41	0,42	0,44	1,14	AT1G18550.1 Symbols: ATP binding microtubule motor family protein chr1:6381656-6384340 REVERSE LENGTH=725
Eucgr.I00528.1	4,99	0,36	0,46	0,72	1,49	AT4G39630.1 Symbols: unknown protein
Eucgr.D01730.1	12,19	0,11	0,82	1,16	3,11	AT4G23820.1 Symbols: Pectin lyase-like superfamily protein chr4:12397217-12400050 REVERSE LENGTH=444
Eucgr.I01145.1	35,01	260,99	270,42	233,44	201,24	AT4G34000.2 Symbols: ABF3, DPBF5 abscisic acid responsive elements-binding factor 3
Eucgr.B03636.1	10,43	0,82	0,87	1,65	4,69	AT5G06700.1 Symbols: TBR Plant protein of unknown function (DUF828) chr5:2063638-2065810 FORWARD LENGTH=608
Eucgr.G02632.1	30,54	89,77	92,26	73,60	63,06	AT3G20770.1 Symbols: EIN3, AtEIN3 Ethylene insensitive 3 family protein chr3:7260702-7262588 REVERSE LENGTH=628
Eucgr.D00364.1	0,31	0,00	0,00	0,02	0,06	AT4G19090.1 Symbols: Protein of unknown function (DUF594) chr4:10449900-10452757 FORWARD LENGTH=751
Eucgr.B02141.1	11,41	69,23	81,64	86,64	39,43	AT1G08650.1 Symbols: PPK1, ATPPPCK1 phosphoenolpyruvate carboxylase kinase 1
Eucgr.J00083.1	1,09	0,08	0,02	0,05	0,23	AT2G38720.1 Symbols: MAP65-5 microtubule-associated protein 65-5 chr2:16188047-16192102 FORWARD LENGTH=587
Eucgr.C03733.1	0,00	1,53	1,26	1,02	0,39	AT1G68180.1 Symbols: RING/U-box superfamily protein chr1:25554816-25555562 FORWARD LENGTH=248
Eucgr.B02161.1	0,00	3,55	3,16	1,33	0,47	AT5G18450.1 Symbols: Integrase-type DNA-binding superfamily protein chr5:6116097-6117020 REVERSE LENGTH=307
Eucgr.I01246.1	46,83	0,15	0,47	0,71	8,11	AT5G48485.1 Symbols: DIR1 Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein
Eucgr.H01151.1	115,54	644,91	537,09	547,36	429,41	AT2G47710.1 Symbols: Adenine nucleotide alpha hydrolases-like superfamily protein
Eucgr.E00270.1	30,29	3,21	3,29	4,03	8,00	AT2G23910.1 Symbols: NAD(P)-binding Rossmann-fold superfamily protein
Eucgr.B00229.1	27,71	1,88	5,18	8,16	11,72	AT4G01680.3 Symbols: MYB55 myb domain protein 55 chr4:716401-717415 REVERSE LENGTH=241
Eucgr.F04115.1	0,02	43,08	21,83	5,80	0,25	AT1G72100.1 Symbols: late embryogenesis abundant domain-containing protein / LEA domain-containing protein
Eucgr.K02271.1	38,30	5,26	10,63	13,16	17,95	AT1G73590.1 Symbols: PIN1, ATPIN1 Auxin efflux carrier family protein chr1:27659772-27662876 FORWARD LENGTH=622
Eucgr.L01670.1	1,15	6,44	8,37	8,55	3,22	AT4G11800.1 Symbols: Calcineurin-like metallo-phosphoesterase superfamily protein
Eucgr.B03139.1	36,26	0,41	1,99	4,46	15,63	AT1G69160.1 Symbols: unknown protein
Eucgr.J00914.1	2,96	0,20	0,21	0,24	0,78	AT2G28260.1 Symbols: ATCNGC15, CNGC15 cyclic nucleotide-gated channel 15
Eucgr.B02031.1	78,79	15,93	17,53	19,21	33,16	AT1G54690.1 Symbols: HTA3, H2AXB, G-H2AX, GAMMA-H2AX gamma histone variant H2AX
Eucgr.K00155.1	14,86	0,17	0,33	0,92	2,69	AT5G62360.1 Symbols: Plant invertase/pectin methylesterase inhibitor superfamily protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.J02110.2	5,44	122,97	79,25	49,75	19,29	AT1G07870.2 Symbols: Protein kinase superfamily protein chr1:2428942-2431843 REVERSE LENGTH=538
Eucgr.F00384.1	3,20	31,55	40,06	36,29	18,56	AT1G72200.1 Symbols: RING/U-box superfamily protein chr1:27169935-27171149 REVERSE LENGTH=404
Eucgr.C00694.1	1,43	0,00	0,00	0,02	0,07	AT5G51160.1 Symbols: Ankyrin repeat family protein chr5:20792280-20793681 FORWARD LENGTH=442
Eucgr.C03431.1	2,27	18,52	21,95	26,89	18,38	AT2G26150.1 Symbols: ATHSFA2, HSFA2 heat shock transcription factor A2
Eucgr.K02165.1	35,62	0,61	2,12	3,43	15,72	AT5G48800.1 Symbols: Phototropic-responsive NPH3 family protein chr5:19786881-19789003 FORWARD LENGTH=614
Eucgr.F00651.1	5,56	35,29	26,29	19,95	17,25	AT4G26780.1 Symbols: AR192 Co-chaperone GrpE family protein chr4:13485066-13486560 REVERSE LENGTH=327

Eucgr.F00721.2	10,47	2,01	2,11	3,28	5,55	AT3G20420.1 Symbols: RTL2, ATRL2 RNase THREE-like protein 2 chr3:7119371-7120895 REVERSE LENGTH=391
Eucgr.D02003.1	1,82	8,13	12,62	9,90	9,12	AT4G13550.1 Symbols: triglyceride lipases;triglyceride lipases chr4:7871251-7876160 REVERSE LENGTH=715
Eucgr.D01813.1	16,66	73,49	65,63	69,23	45,16	AT5G41560.1 Symbols: unknown protein
Eucgr.J00194.1	0,05	46,26	27,67	16,03	0,76	AT2G40170.1 Symbols: ATEM6, GEA6, EM6 Stress induced protein chr2:16779792-16780167 REVERSE LENGTH=92
Eucgr.B00829.2	40,14	9,60	10,61	14,94	20,66	AT1G73760.1 Symbols: RING/U-box superfamily protein chr1:27739366-27741161 REVERSE LENGTH=367
Eucgr.A02029.1	7,37	1,21	0,84	0,95	1,45	AT1G06660.1 Symbols: unknown protein
Eucgr.A01283.1	11,63	41,57	33,78	36,12	24,83	AT2G37678.1 Symbols: FHY1, PAT3, FRY1 far-red elongated hypocotyl 1 chr2:15801689-15802696 REVERSE LENGTH=202
Eucgr.F02300.1	15,32	0,62	0,79	1,36	1,68	AT3G16150.1 Symbols: N-terminal nucleophile aminohydrolases (Ntn hydrolases) superfamily protein
Eucgr.F03035.1	3,90	32,06	28,94	39,63	12,29	AT1G52240.2 Symbols: ATROPGEF11, ROPGEF11, PIRF1 RHO guanyl-nucleotide exchange factor 11
Eucgr.B03399.1	19,79	76,66	68,25	64,66	42,11	AT1G27150.1 Symbols: Tetratricopeptide repeat (TPR)-like superfamily protein
Eucgr.C01731.1	18,89	101,85	127,26	132,48	46,83	No hits found
Eucgr.I01495.1	7,74	1562,71	1211,97	880,37	92,12	AT3G12500.1 Symbols: ATHCHIB, PR3, PR-3, CHI-B, B-CHI, HCHIB basic chitinase
Eucgr.B02467.1	5,95	19,77	20,56	15,35	12,74	AT1G07910.2 Symbols: ATRNL, RNL, AtRLG1 RNAligase chr1:2446712-2454386 FORWARD LENGTH=1104
Eucgr.F01003.1	2,30	43,54	47,40	43,89	6,02	AT5G48150.2 Symbols: PAT1 GRAS family transcription factor chr5:19522497-19524053 REVERSE LENGTH=490
Eucgr.D02320.1	24,92	79,39	63,57	56,94	53,81	AT4G01280.2 Symbols: Homeodomain-like superfamily protein chr4:535288-536854 FORWARD LENGTH=303
Eucgr.K03370.1	13,37	112,45	118,15	116,79	43,19	AT3G63210.1 Symbols: MARD1 Protein of unknown function (DUF581) chr3:23354019-23354906 REVERSE LENGTH=263
Eucgr.F02465.1	0,35	0,01	0,01	0,02	0,09	AT3G46530.1 Symbols: RPP13 NB-ARC domain-containing disease resistance protein
Eucgr.G03232.1	16,14	0,72	1,57	2,15	5,91	AT5G12330.1 Symbols: LRP1 Lateral root primordium (LRP) protein-related chr5:3987677-3989074 REVERSE LENGTH=320
Eucgr.J02591.1	15,10	0,88	1,27	1,99	7,78	AT1G53820.1 Symbols: RING/U-box superfamily protein chr1:20091491-20092423 FORWARD LENGTH=310
Eucgr.F03575.1	62,99	0,76	1,47	2,24	19,19	AT5G24318.1 Symbols: O-Glycosyl hydrolases family 17 protein chr5:8282285-8283956 REVERSE LENGTH=458
Eucgr.F02000.1	6,72	0,16	0,26	0,40	0,90	AT5G56120.1 Symbols: unknown protein;
Eucgr.B03665.1	41,27	4,87	5,22	4,68	13,19	AT4G40030.2 Symbols: Histone superfamily protein chr4:18555840-18556827 REVERSE LENGTH=164
Eucgr.C00665.1	5,03	53,29	66,19	45,36	24,04	AT2G26170.1 Symbols: CYP711A1, MAX1 cytochrome P450, family 711, subfamily A, polypeptide 1
Eucgr.K02533.1	8,40	0,86	0,73	0,93	2,34	AT5G60980.1 Symbols: Nuclear transport factor 2 (NTF2) family protein with RNA binding (RRM-RBD-RNP motifs) domain
Eucgr.F01854.1	4,16	598,03	295,93	191,61	25,01	AT3G51030.1 Symbols: ATTRX1, ATTRX H1, TRX1 thioredoxin H-type 1 chr3:18951123-18951955 REVERSE LENGTH=114
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H01055.1	168,37	2,29	5,79	8,99	36,59	AT1G02640.1 Symbols: BXL2, ATBXL2 beta-xylosidase 2 chr1:564293-567580 FORWARD LENGTH=768
Eucgr.C00146.1	5,51	0,00	0,05	0,09	1,25	AT1G73300.1 Symbols: scpl2 serine carboxypeptidase-like 2 chr1:27559673-27562048 REVERSE LENGTH=441
Eucgr.I01275.1	6,36	0,00	0,03	0,05	0,51	AT1G29420.1 Symbols: SAUR-like auxin-responsive protein family chr1:10301483-10301908 REVERSE LENGTH=141
Eucgr.L02979.1	3,66	0,32	0,27	0,58	1,20	AT3G20290.2 Symbols: ATEHD1, EHD1 EPS15 homology domain 1 chr3:7075057-7078655 REVERSE LENGTH=545

Eucgr.D01672.1	4,58	0,48	0,70	0,94	1,71	AT4G10630.1 Symbols: Glutaredoxin family protein chr4:6566625-6567629 REVERSE LENGTH=334
Eucgr.I01413.2	2,11	0,20	0,39	0,30	0,80	AT2G20980.1 Symbols: MCM10 minichromosome maintenance 10 chr2:9011654-9013856 REVERSE LENGTH=396
Eucgr.A02158.1	3,13	8,15	9,16	8,09	7,08	AT3G03620.1 Symbols: MATE efflux family protein chr3:873904-876252 REVERSE LENGTH=500
Eucgr.G02325.1	2,51	28,24	20,16	21,06	9,55	AT1G68540.1 Symbols: NAD(P)-binding Rossmann-fold superfamily protein
Eucgr.I01172.2	11,08	34,85	26,74	24,74	22,72	AT4G33940.1 Symbols: RING/U-box superfamily protein chr4:16266085-16267612 FORWARD LENGTH=262
Eucgr.D02454.1	0,05	3,10	2,73	1,32	0,48	AT2G45400.1 Symbols: BEN1 NAD(P)-binding Rossmann-fold superfamily protein
Eucgr.D02266.1	0,01	0,63	0,55	0,39	0,22	AT3G61220.1 Symbols: NAD(P)-binding Rossmann-fold superfamily protein
Eucgr.C00147.1	10,43	0,02	0,09	0,16	2,72	AT5G09640.1 Symbols: SNG2, SCPL19 serine carboxypeptidase-like 19 chr5:2988373-2990966 FORWARD LENGTH=465
Eucgr.H00469.1	10,72	49,66	51,07	56,64	35,72	AT1G60710.1 Symbols: ATB2 NAD(P)-linked oxidoreductase superfamily protein
Eucgr.B00095.1	7,72	26,19	28,29	26,11	15,98	AT5G63520.1 Symbols: unknown protein
Eucgr.H00093.1	30,03	0,37	1,49	1,77	11,45	AT4G13710.2 Symbols: Pectin lyase-like superfamily protein chr4:7962550-7966012 FORWARD LENGTH=451
Eucgr.I00720.1	13,06	1,59	2,52	4,26	6,80	AT4G24230.5 Symbols: ACBP3 acyl-CoA-binding domain 3 chr4:12567240-12568754 REVERSE LENGTH=362
Eucgr.B02195.1	28,06	0,62	1,86	3,32	12,66	AT5G26670.1 Symbols: Pectinacetylerase family protein chr5:9318456-9320816 FORWARD LENGTH=416
Eucgr.A01180.1	21,59	1072,43	901,04	862,26	494,83	AT3G09640.2 Symbols: APX2, APX18 ascorbate peroxidase 2 chr3:2956301-2958163 FORWARD LENGTH=251
Eucgr.F00208.1	17,00	0,40	1,14	1,46	9,35	AT5G67090.1 Symbols: Subtilisin-like serine endopeptidase family protein
Eucgr.A02334.1	2,15	0,13	0,14	0,17	0,46	AT3G10310.1 Symbols: P-loop nucleoside triphosphate hydrolases superfamily protein
Eucgr.B03598.1	14,71	2,15	2,68	3,16	7,56	AT5G36110.1 Symbols: CYP716A1 cytochrome P450, family 716, subfamily A, polypeptide 1
Eucgr.H01197.1	19,97	0,40	2,28	4,02	7,96	AT1G03010.1 Symbols: Phototropic-responsive NPH3 family protein chr1:693480-696188 FORWARD LENGTH=634
Eucgr.J02961.1	0,07	17,02	7,45	5,89	1,20	No hits found
Eucgr.B00628.1	7,37	0,67	1,23	2,00	3,23	AT3G02210.1 Symbols: COBL1 COBRA-like protein 1 precursor chr3:409352-411478 REVERSE LENGTH=452
Eucgr.K02109.1	32,67	0,56	1,51	2,48	14,92	AT3G07340.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.H01519.1	1,16	227,04	274,22	267,45	16,07	AT3G63060.1 Symbols: EDL3 EID1-like 3 chr3:23300638-23301456 REVERSE LENGTH=272
Eucgr.A02002.1	0,20	1,24	2,13	1,38	0,79	AT1G06490.1 Symbols: ATGSL07, gsl07, atgsl7, GSL7 glucan synthase-like 7
Eucgr.H01011.1	11,23	30,28	36,66	29,95	25,22	AT4G02020.1 Symbols: EZA1, SWN, SDG10 SET domain-containing protein chr4:886693-891743 FORWARD LENGTH=856
Eucgr.I02738.1	157,61	5,61	7,44	9,62	60,30	AT2G46890.1 Symbols: Protein of unknown function (DUF1295) chr2:19266879-19268134 REVERSE LENGTH=322
Eucgr.K03279.1	36,62	2,45	5,72	8,68	15,81	AT5G51560.1 Symbols: Leucine-rich repeat protein kinase family protein
Eucgr.A01044.1	36,16	150,94	144,45	134,06	98,25	AT3G07770.1 Symbols: Hsp89.1, AtHsp90.6, AtHsp90-6 HEAT SHOCK PROTEIN 89.1
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.J02057.1	18,53	0,33	0,61	1,56	7,99	AT5G02440.1 Symbols: unknown protein
Eucgr.I01789.4	3,31	41,00	42,66	37,20	18,16	AT3G01510.1 Symbols: LSF1 like SEX4 1 chr3:198855-201682 REVERSE LENGTH=591

Eucgr.G02817.1	14,61	0,63	0,65	0,69	5,55	AT2G35770.1 Symbols: scpl28 serine carboxypeptidase-like 28 chr2:15034179-15036497 REVERSE LENGTH=462
Eucgr.J03191.1	3,93	15,89	16,74	13,92	10,60	AT1G78930.1 Symbols: Mitochondrial transcription termination factor family protein
Eucgr.F03345.1	22,16	1,38	1,91	2,12	9,18	AT5G54380.1 Symbols: THE1 protein kinase family protein chr5:22077313-22079880 REVERSE LENGTH=855
Eucgr.D00636.2	15,82	76,56	66,45	71,30	29,85	AT3G47160.2 Symbols: RING/U-box superfamily protein chr3:17360238-17361879 REVERSE LENGTH=257
Eucgr.B02770.1	13,08	2,54	3,55	4,43	6,83	AT2G38040.2 Symbols: CAC3 acetyl Co-enzyme a carboxylase carboxyltransferase alpha subunit
Eucgr.E00105.1	77,46	1,55	4,11	6,21	19,93	AT3G61490.3 Symbols: Pectin lyase-like superfamily protein chr3:22758439-22760137 FORWARD LENGTH=476
Eucgr.F02730.1	5,89	54,46	61,88	65,31	29,37	AT1G21360.1 Symbols: GLTP2 glycolipid transfer protein 2 chr1:7481365-7483237 FORWARD LENGTH=223
Eucgr.B03070.1	2,83	0,16	0,05	0,09	0,33	AT1G68740.1 Symbols: PHO1;H1 EXS (ERD1/XPR1/SYG1) family protein chr1:25812735-25816574 REVERSE LENGTH=784
Eucgr.F00495.1	7,54	83,26	77,38	55,20	28,40	No hits found
Eucgr.D01034.1	14,55	0,54	1,23	1,91	5,33	AT3G54950.1 Symbols: PLP7, PLA IIIA patatin-like protein 6 chr3:20359076-20360774 REVERSE LENGTH=488
Eucgr.A02438.1	4,71	0,34	0,59	0,95	1,89	No hits found
Eucgr.A02689.1	1,92	0,07	0,05	0,06	0,69	AT3G16520.1 Symbols: UGT88A1 UDP-glucosyl transferase 88A1 chr3:5618807-5620833 REVERSE LENGTH=451
Eucgr.K03540.1	13,07	42,64	35,67	35,25	31,50	AT4G02090.1 Symbols: unknown protein
Eucgr.F01784.1	3,28	10,77	10,55	9,53	8,03	No hits found
Eucgr.K02134.1	31,03	0,45	1,05	2,38	10,49	AT3G25130.1 Symbols: unknown protein
Eucgr.J00323.1	4,96	0,58	0,61	1,59	1,54	AT5G23440.1 Symbols: FTAA1 ferredoxin/thioredoxin reductase subunit A (variable subunit) 1
Eucgr.K01300.1	35,05	0,44	1,52	2,44	13,01	AT3G08600.1 Symbols: Protein of unknown function (DUF1191) chr3:2612646-2613596 FORWARD LENGTH=316
Eucgr.D02381.1	102,26	8,94	12,83	11,83	27,84	AT1G01300.1 Symbols: Eukaryotic aspartyl protease family protein chr1:117065-118522 FORWARD LENGTH=485
Eucgr.C00368.1	2,43	8,19	9,12	9,70	6,55	AT5G58050.1 Symbols: SVL4 SHV3-like 4 chr5:23494498-23497386 REVERSE LENGTH=753
Eucgr.B01111.1	58,85	12,93	15,61	17,59	32,48	AT4G14220.1 Symbols: RHF1A RING-H2 group F1A chr4:8196012-8198240 FORWARD LENGTH=371
Eucgr.G02550.2	20,18	102,70	103,74	103,49	56,62	AT1G07040.1 Symbols: unknown protein
Eucgr.D00296.1	15,53	3,16	3,29	4,46	8,41	AT2G34560.1 Symbols: P-loop containing nucleoside triphosphate hydrolases superfamily protein
Eucgr.H03285.1	53,31	489,01	664,72	513,12	242,38	AT1G15690.1 Symbols: AVP1, ATAVP3, AVP-3, AtVHP1;1 Inorganic H pyrophosphatase family protein
Eucgr.B03561.1	28,70	3,74	3,91	4,74	11,45	AT1G57820.1 Symbols: VIM1, ORTH2 Zinc finger (C3HC4-type RING finger) family protein
Eucgr.F01527.1	0,78	23,96	25,97	17,49	5,52	AT1G47530.1 Symbols: MATE efflux family protein chr1:17451724-17454110 FORWARD LENGTH=484
Eucgr.H04950.1	1,81	0,31	0,22	0,86	0,56	AT1G77120.1 Symbols: ADH1, ADH, ATADH, ATADH1 alcohol dehydrogenase 1
Eucgr.I01337.1	21,93	1,56	2,71	5,84	10,35	AT4G34750.2 Symbols: SAUR-like auxin-responsive protein family chr4:16577566-16578018 FORWARD LENGTH=150
Eucgr.F03278.2	36,02	5,60	6,03	8,41	16,58	AT1G16920.1 Symbols: RAB11, ATRABA1B, RABA1b RAB GTPase homolog A1B
Eucgr.K01646.1	25,16	4,47	5,36	7,15	14,82	AT3G56750.1 Symbols: unknown protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR

Eucgr.H03767.1	39,26	225,21	235,10	217,13	115,08	AT5G18670.1 Symbols: BMY3, BAM9 beta-amylase 3 chr5:6226138-6227999 FORWARD LENGTH=536
Eucgr.D01613.2	79,21	16,99	16,29	19,60	29,09	AT2G34770.1 Symbols: FAH1, ATFAH1 fatty acid hydroxylase 1 chr2:14666776-14668061 FORWARD LENGTH=237
Eucgr.B02615.1	7,49	0,32	0,25	0,36	2,19	AT1G69770.1 Symbols: CMT3 chromomethylase 3 chr1:26248496-26253519 REVERSE LENGTH=839
Eucgr.A01900.1	19,76	1,28	1,41	2,26	7,55	AT2G42570.1 Symbols: TBL39 TRICHOME BIREFRINGENCE-LIKE 39 chr2:17717498-17719921 REVERSE LENGTH=367
Eucgr.F04351.1	23,62	78,31	73,21	72,29	59,16	AT1G21780.2 Symbols: BTB/POZ domain-containing protein chr1:7652476-7653866 FORWARD LENGTH=326
Eucgr.H01465.1	0,00	2,60	2,10	0,55	0,14	AT3G22490.1 Symbols: Seed maturation protein chr3:7969785-7970738 REVERSE LENGTH=262
Eucgr.K01373.1	0,00	2,23	1,13	1,03	0,09	No hits found
Eucgr.G01861.1	0,00	9,28	6,63	2,41	0,22	AT2G23240.1 Symbols: Plant EC metallothionein-like protein, family 15 chr2:9895995-9896325 REVERSE LENGTH=85
Eucgr.G00824.1	0,00	0,81	0,61	0,30	0,10	AT5G45910.1 Symbols: GDSL-like Lipase/Acylhydrolase superfamily protein chr5:18620420-18622264 REVERSE LENGTH=372
Eucgr.H04654.1	69,13	4,69	10,72	14,51	33,14	AT1G67510.1 Symbols: Leucine-rich repeat protein kinase family protein chr1:25297477-25300184 REVERSE LENGTH=719
Eucgr.G00172.1	3,61	10,52	12,37	12,56	10,02	No hits found
Eucgr.H03574.1	7,17	0,47	0,87	1,20	3,68	AT5G19160.1 Symbols: TBL11 TRICHOME BIREFRINGENCE-LIKE 11 chr5:6430725-6432456 FORWARD LENGTH=464
Eucgr.A02652.1	3,53	0,24	0,86	1,17	1,59	AT4G21580.1 Symbols: oxidoreductase, zinc-binding dehydrogenase family protein
Eucgr.C00351.1	28,09	1,62	3,25	4,69	11,75	AT2G24230.1 Symbols: Leucine-rich repeat protein kinase family protein chr2:10301979-10304540 REVERSE LENGTH=853
Eucgr.A00199.1	7,15	0,43	0,52	0,45	1,24	AT3G58650.1 Symbols: unknown protein
Eucgr.I01272.1	19,58	2,97	3,65	3,72	7,31	AT2G16250.1 Symbols: Leucine-rich repeat protein kinase family protein chr2:7039682-7042933 REVERSE LENGTH=915
Eucgr.I02050.1	2,16	0,00	0,04	0,24	0,70	No hits found
Eucgr.I02034.1	1,63	0,08	0,03	0,08	0,32	AT3G02120.1 Symbols: hydroxyproline-rich glycoprotein family protein chr3:377122-377577 FORWARD LENGTH=126
Eucgr.E03946.1	0,56	0,05	0,01	0,07	0,18	AT1G21270.1 Symbols: WAK2 wall-associated kinase 2 chr1:7444997-7447345 FORWARD LENGTH=732
Eucgr.G03101.1	0,10	0,95	1,31	1,00	0,41	AT5G01660.1 Symbols: unknown protein
Eucgr.B02465.1	6,80	0,14	0,34	0,26	1,95	AT1G07900.1 Symbols: LBD1 LOB domain-containing protein 1 chr1:2442624-2443631 FORWARD LENGTH=190
Eucgr.C00688.1	1,89	0,00	0,00	0,06	0,12	AT5G51160.1 Symbols: Ankyrin repeat family protein chr5:20792280-20793681 FORWARD LENGTH=442
Eucgr.J01649.1	6,12	0,36	0,25	0,33	1,45	AT5G37010.1 Symbols: unknown protein
Eucgr.B00858.1	2,37	0,22	0,14	0,13	0,78	AT1G18090.1 Symbols: 5'-3' exonuclease family protein chr1:6224539-6227715 FORWARD LENGTH=577
Eucgr.C00602.1	2,41	0,15	0,26	0,57	0,90	AT5G25830.1 Symbols: GATA12 GATA transcription factor 12 chr5:9004398-9005502 REVERSE LENGTH=331
Eucgr.H04126.1	0,21	116,86	67,26	27,17	1,14	No hits found
Eucgr.F01532.1	3,53	0,55	0,25	0,49	0,60	AT5G42610.1 Symbols: Protein of unknown function (DUF607) chr5:17062507-17063934 FORWARD LENGTH=293
Eucgr.F03192.1	8,01	0,58	0,96	1,05	1,83	AT1G72250.2 Symbols: Di-glucose binding protein with Kinesin motor domain
Eucgr.E03164.1	1,54	0,00	0,00	0,00	0,20	AT2G34060.1 Symbols: Peroxidase superfamily protein chr2:14384914-14386530 FORWARD LENGTH=346
Eucgr.D02141.1	49,41	4,95	8,75	12,20	28,73	AT2G35190.1 Symbols: NPSN11, ATNPSN11, NSPN11 novel plant snare 11
GENE ID	FPKM					Annotation

	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H03412.1	2,02	25,37	23,40	18,00	8,94	AT3G22830.1 Symbols: AT-HSFA6B, HSFA6B heat shock transcription factor A6B
Eucgr.E02751.1	9,90	1,01	1,14	1,18	3,04	AT3G19390.1 Symbols: Granulin repeat cysteine protease family protein chr3:6723024-6724768 FORWARD LENGTH=452
Eucgr.E00653.1	20,86	998,82	820,18	566,16	475,38	AT5G52640.1 Symbols: HSP81-1, ATHS83, HSP81.1, HSP83, ATHSP90.1, AtHsp90-1, HSP90.1 heat shock protein 90.1
Eucgr.K02631.1	3,76	0,12	0,20	0,24	1,03	AT3G42660.1 Symbols: transducin family protein / WD-40 repeat family protein
Eucgr.H01935.1	0,21	16,09	12,48	11,36	1,34	No hits found
Eucgr.H00738.1	19,53	1,56	3,65	5,56	7,38	AT3G25290.2 Symbols: Auxin-responsive family protein chr3:9208955-9210353 FORWARD LENGTH=393
Eucgr.C03734.2	21,41	114,73	107,58	113,74	67,99	AT3G46500.1 Symbols: 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein
Eucgr.A00685.1	4,91	0,77	0,36	1,25	1,12	AT2G40475.1 Symbols: unknown protein;
Eucgr.G02914.1	2,55	0,04	0,04	0,03	0,28	AT2G39510.1 Symbols: nodulin MtN21 /EamA-like transporter family protein
Eucgr.I02446.1	25,27	76,29	54,91	49,57	50,51	AT5G13970.1 Symbols: unknown protein;
Eucgr.A01510.1	6,01	0,07	0,55	1,11	2,22	AT3G52780.1 Symbols: ATPAP20, PAP20 Purple acid phosphatases superfamily protein
Eucgr.J03116.1	35,53	243,22	220,93	205,31	109,05	AT5G25110.1 Symbols: CIPK25, SnRK3.25 CBL-interacting protein kinase 25 chr5:8657740-8659206 REVERSE LENGTH=488
Eucgr.G01719.1	6,80	0,33	0,44	0,55	2,35	AT2G21340.1 Symbols: MATE efflux family protein chr2:9132629-9136236 FORWARD LENGTH=559
Eucgr.K01326.1	4,13	0,23	0,50	0,59	1,79	AT1G47056.1 Symbols: VFB1 VIER F-box proteine 1 chr1:17276103-17277659 REVERSE LENGTH=518
Eucgr.A02419.1	4,27	0,23	0,20	0,54	1,02	AT2G42260.1 Symbols: UVI4, PYM uv-b-insensitive 4 chr2:17602240-17603283 REVERSE LENGTH=259
Eucgr.F02608.1	40,99	0,88	1,99	3,75	19,27	AT1G71970.1 Symbols: unknown protein
Eucgr.C03407.2	36,43	450,65	402,82	402,69	157,74	AT1G17550.1 Symbols: HAB2 homology to ABI2 chr1:6034917-6036939 FORWARD LENGTH=511
Eucgr.F03662.1	1,19	0,02	0,02	0,06	0,30	AT5G59810.1 Symbols: ATSBT5.4, SBT5.4 Subtilase family protein chr5:24096895-24100387 REVERSE LENGTH=778
Eucgr.G02658.1	20,17	0,47	1,02	1,26	3,58	AT3G03130.1 Symbols: unknown protein
Eucgr.B02372.1	7,27	32,99	36,44	42,25	23,11	AT3G46540.1 Symbols: ENTH/VHS family protein chr3:17134204-17135411 REVERSE LENGTH=307
Eucgr.A01404.1	0,64	9,23	6,74	5,80	2,56	AT5G53730.1 Symbols: Late embryogenesis abundant (LEA) hydroxyproline-rich glycoprotein family
Eucgr.A01247.1	1,43	0,14	0,08	0,07	0,51	AT5G02540.1 Symbols: NAD(P)-binding Rossmann-fold superfamily protein chr5:568609-570720 FORWARD LENGTH=331
Eucgr.K00764.1	0,95	7,77	4,52	3,85	3,04	AT2G22475.1 Symbols: GEM GRAM domain family protein chr2:9541523-9544778 FORWARD LENGTH=299
Eucgr.B02355.1	323,58	2,45	9,49	18,01	119,17	AT2G39700.1 Symbols: ATEXPA4, ATEXP4, ATHEXP ALPHA 1.6, EXPA4 expansin A4
Eucgr.A02869.1	21,72	5,23	5,88	8,03	13,66	AT5G51710.2 Symbols: KEA5, ATKEA5 K+ efflux antiporter 5 chr5:21004566-21008580 REVERSE LENGTH=565
Eucgr.G02884.1	22,55	0,40	0,29	1,47	10,73	AT5G04310.1 Symbols: Pectin lyase-like superfamily protein chr5:1203356-1207352 REVERSE LENGTH=518
Eucgr.J00542.1	101,89	11,05	9,82	10,65	36,38	AT3G44220.1 Symbols: Late embryogenesis abundant (LEA) hydroxyproline-rich glycoprotein family
Eucgr.F01953.2	15,06	0,43	1,14	1,84	8,49	AT1G21060.1 Symbols: Protein of unknown function, DUF547 chr1:7371799-7374085 FORWARD LENGTH=505
Eucgr.D02489.1	4,28	0,46	0,40	0,64	1,17	AT3G60660.1 Symbols: unknown protein;
Eucgr.J00903.1	13,90	0,47	0,99	1,29	7,35	AT5G04250.2 Symbols: Cysteine proteinases superfamily protein chr5:1176397-1178492 FORWARD LENGTH=345

Eucgr.F03301.1	1,14	0,05	0,20	0,56	0,33	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.F02302.1	26,54	0,78	1,24	1,69	6,64	AT4G34160.1 Symbols: CYCD3;1, CYCD3 CYCLIN D3;1 chr4:16357903-16359304 FORWARD LENGTH=376
Eucgr.E02910.1	17,99	3,07	2,73	3,54	8,95	No hits found
Eucgr.B01911.5	45,47	276,97	253,38	191,66	159,73	AT1G09140.2 Symbols: ATSRP30, SR30, At-SR30 SERINE-ARGININE PROTEIN 30
Eucgr.B00048.1	7,18	0,00	0,31	0,39	2,44	AT1G72210.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.K02942.1	28,02	4,38	5,40	5,70	11,94	AT5G01710.1 Symbols: methyltransferases chr5:263709-265250 REVERSE LENGTH=513
Eucgr.J03133.1	3,45	0,08	0,16	0,33	0,80	AT1G16070.1 Symbols: AtTLP8, TLP8 tubby like protein 8 chr1:5511899-5513779 REVERSE LENGTH=397
Eucgr.H00185.1	0,46	0,00	0,00	0,14	0,11	AT4G23990.1 Symbols: ATCSLG3, CSLG3 cellulose synthase like G3 chr4:12456491-12460498 FORWARD LENGTH=751
Eucgr.K00751.1	7,65	0,04	0,15	0,10	0,83	AT4G03965.1 Symbols: RING/U-box superfamily protein chr4:1889576-1890205 FORWARD LENGTH=209
Eucgr.G02816.1	16,72	1,26	1,78	1,84	5,78	AT3G07990.1 Symbols: SCPL27 serine carboxypeptidase-like 27 chr3:2552544-2554644 FORWARD LENGTH=459
Eucgr.E03124.1	9,69	2,73	2,82	4,77	5,27	AT4G04930.1 Symbols: DES-1-LIKE fatty acid desaturase family protein chr4:2509236-2510918 FORWARD LENGTH=332
Eucgr.A01002.1	9,58	0,94	1,05	0,99	2,24	AT4G01730.1 Symbols: DHHC-type zinc finger family protein chr4:749574-752034 FORWARD LENGTH=508
Eucgr.J00148.1	2,63	0,18	0,09	0,08	0,58	No hits found
Eucgr.D00751.1	0,02	7,90	3,84	1,41	0,21	AT4G21440.1 Symbols: ATMYB102, ATM4, MYB102 MYB-like 102 chr4:11418425-11419652 REVERSE LENGTH=350
Eucgr.G01894.3	28,14	3,39	3,61	5,80	12,54	AT1G47740.2 Symbols: PPPDE putative thiol peptidase family protein chr1:17567903-17569035 FORWARD LENGTH=279
Eucgr.K02233.1	7,39	0,26	0,40	0,40	1,27	AT1G18370.1 Symbols: HIK, ATNACK1, NACK1 ATP binding microtubule motor family protein
Eucgr.D01694.1	15,86	131,02	112,03	96,82	79,44	AT5G63830.1 Symbols: HIT-type Zinc finger family protein chr5:25543585-25544802 REVERSE LENGTH=405
Eucgr.F03980.1	29,88	1058,66	920,98	642,11	1318,09	AT3G12580.1 Symbols: HSP70, ATHSP70 heat shock protein 70 chr3:3991487-3993689 REVERSE LENGTH=650
Eucgr.H04428.2	37,29	7,59	7,86	8,52	17,19	AT3G06350.1 Symbols: EMB3004, MEE32 dehydroquinase dehydratase, putative / shikimate dehydrogenase, putative
Eucgr.G01342.1	193,47	1,29	2,45	4,01	67,16	AT4G15480.1 Symbols: UGT84A1 UDP-Glycosyltransferase superfamily protein
Eucgr.D01985.1	4,75	0,07	0,11	0,07	0,89	AT1G12370.2 Symbols: PHR1, UVR2 photolyase 1 chr1:4206500-4208842 REVERSE LENGTH=496
Eucgr.L03161.1	0,71	0,11	0,03	0,04	0,18	AT1G21270.1 Symbols: WAK2 wall-associated kinase 2 chr1:7444997-7447345 FORWARD LENGTH=732
Eucgr.K01547.1	8,25	26,70	32,27	37,79	24,48	AT1G08710.2 Symbols: F-box family protein chr1:2771720-2773011 FORWARD LENGTH=295
Eucgr.G02912.1	3,83	0,04	0,03	0,10	0,37	AT2G39510.1 Symbols: nodulin MtN21 /EamA-like transporter family protein
Eucgr.H00919.1	6,02	489,18	193,76	106,41	20,09	AT4G08570.1 Symbols: Heavy metal transport/detoxification superfamily protein
Eucgr.A01714.1	39,83	7,62	9,46	11,10	22,23	AT1G05210.1 Symbols: Transmembrane protein 97, predicted chr1:1510469-1511485 FORWARD LENGTH=168
Eucgr.C01067.1	14,85	0,50	0,74	0,72	6,92	AT3G24460.1 Symbols: Serinc-domain containing serine and sphingolipid biosynthesis protein
Eucgr.B00137.1	12,31	1,83	2,50	4,21	6,24	AT5G63380.1 Symbols: AMP-dependent synthetase and ligase family protein
Eucgr.J00969.2	34,52	2,05	2,92	4,23	13,80	AT2G39870.1 Symbols: unknown protein

Eucgr.J00271.1	8,34	299,91	192,56	122,45	31,18	AT5G04760.1 Symbols: Duplicated homeodomain-like superfamily protein chr5:1373752-1374529 REVERSE LENGTH=215
Eucgr.L03024.1	1,02	0,00	0,02	0,03	0,24	AT4G18780.1 Symbols: CESA8, IRX1, ATCESA8, LEW2 cellulose synthase family protein
Eucgr.C02352.1	11,51	0,00	0,19	0,14	2,09	AT1G33540.1 Symbols: scpl18 serine carboxypeptidase-like 18 chr1:12162349-12164700 REVERSE LENGTH=446
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.F02338.1	0,42	11,09	6,84	6,32	2,78	No hits found
Eucgr.E00226.1	10,13	0,53	0,78	0,68	2,11	AT3G03050.1 Symbols: CSLD3, KJK, ATCSLD3 cellulose synthase-like D3 chr3:687873-691629 FORWARD LENGTH=1145
Eucgr.K02618.1	2,22	0,26	0,18	0,32	0,64	AT4G31400.1 Symbols: CTF7 damaged DNA binding;DNA-directed DNA polymerases
Eucgr.A00994.1	20,90	2,91	3,97	4,36	13,00	AT2G46930.1 Symbols: Pectinacetylsterase family protein chr2:19283625-19286090 FORWARD LENGTH=416
Eucgr.G00557.1	16,97	46,33	47,35	51,16	34,05	AT1G60940.2 Symbols: SNRK2-10, SNRK2.10, SRK2B SNF1-related protein kinase 2.10
Eucgr.H03092.1	6,20	0,10	0,16	0,39	2,08	AT5G43420.1 Symbols: RING/U-box superfamily protein chr5:17451790-17452917 FORWARD LENGTH=375
Eucgr.A00530.1	41,00	2,35	4,67	5,83	16,20	AT5G01360.1 Symbols: TBL3 Plant protein of unknown function (DUF828) chr5:147608-149316 REVERSE LENGTH=434
Eucgr.L00249.1	0,34	280,16	135,36	52,37	1,25	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.A02410.1	25,38	1,89	3,12	5,75	12,43	AT5G06940.1 Symbols: Leucine-rich repeat receptor-like protein kinase family protein
Eucgr.J00169.1	14,01	0,35	0,55	1,29	3,34	AT2G38310.1 Symbols: PYL4, RCAR10 PYR1-like 4 chr2:16050251-16050874 FORWARD LENGTH=207
Eucgr.E00402.1	12,99	40,59	41,66	32,07	27,12	AT2G39810.1 Symbols: HOS1 ubiquitin-protein ligases chr2:16612941-16617802 FORWARD LENGTH=927
Eucgr.C04023.1	5,44	20,57	21,03	21,19	11,05	AT5G24970.2 Symbols: Protein kinase superfamily protein chr5:8604358-8608656 REVERSE LENGTH=761
Eucgr.F00103.2	21,10	0,55	1,06	1,51	6,47	AT1G58440.1 Symbols: XF1, SQE1 FAD/NAD(P)-binding oxidoreductase family protein
Eucgr.F02263.1	27,76	3,11	7,28	13,35	16,39	AT1G49890.1 Symbols: QWRF2 Family of unknown function (DUF566) chr1:18470282-18473463 FORWARD LENGTH=659
Eucgr.H03407.1	9,56	0,39	0,69	1,22	3,60	No hits found
Eucgr.A02144.1	6,82	17,97	18,26	15,96	12,00	AT3G09360.1 Symbols: Cyclin/Brf1-like TBP-binding protein chr3:2873796-2878432 FORWARD LENGTH=604
Eucgr.F01560.1	37,72	2,92	4,59	6,97	19,00	AT3G46940.1 Symbols: DUT1 DUTP-PYROPHOSPHATASE-LIKE 1 chr3:17288367-17288867 REVERSE LENGTH=166
Eucgr.D01775.1	1,57	0,04	0,52	0,56	0,55	AT4G23750.2 Symbols: CRF2, TMO3 cytokinin response factor 2 chr4:12376751-12377782 FORWARD LENGTH=343
Eucgr.F02691.1	2,11	16,62	17,60	13,57	7,69	AT5G50080.1 Symbols: ERF110 ethylene response factor 110 chr5:20365948-20366835 FORWARD LENGTH=220
Eucgr.F01369.1	5,98	1,26	1,44	1,99	3,30	AT1G78570.1 Symbols: RHM1, ROL1, ATRHM1 rhamnose biosynthesis 1
Eucgr.B03763.1	6,04	0,07	0,22	0,52	1,83	AT4G13870.2 Symbols: WRNEXO Werner syndrome-like exonuclease chr4:8023563-8025542 REVERSE LENGTH=288
Eucgr.K00770.2	5,60	33,27	23,33	23,26	13,75	AT2G32650.2 Symbols: RmlC-like cupins superfamily protein chr2:13851547-13851966 FORWARD LENGTH=139
Eucgr.J01951.2	42,69	16,98	14,79	15,70	24,39	AT2G29570.1 Symbols: PCNA2, ATPCNA2 proliferating cell nuclear antigen 2
Eucgr.I02584.1	13,10	2,99	2,97	3,32	5,76	AT2G04305.1 Symbols: Magnesium transporter CorA-like family protein chr2:1501679-1503448 REVERSE LENGTH=434
Eucgr.B00246.1	22,27	63,49	52,27	51,95	41,44	AT1G17100.1 Symbols: SOUL heme-binding family protein chr1:5844766-5845539 FORWARD LENGTH=232
Eucgr.C00213.3	0,28	9,02	5,83	5,10	1,59	No hits found

Eucgr.J00420.1	41,08	2,63	5,07	5,87	25,59	AT5G22740.1 Symbols: ATCSLA02, CSLA02, ATCSLA2, CSLA2 cellulose synthase-like A02
Eucgr.K01046.1	19,54	2,24	2,77	4,55	8,76	AT2G47070.1 Symbols: SPL1 squamosa promoter binding protein-like 1
Eucgr.E00628.2	225,84	1457,57	1141,60	1015,46	836,98	AT2G45960.3 Symbols: PIP1B, TMP-A, ATHH2, PIP1;2 plasma membrane intrinsic protein 1B
Eucgr.E02426.1	1,67	11,42	9,63	7,40	4,00	AT1G30220.1 Symbols: ATINT2, INT2 inositol transporter 2 chr1:10632957-10635439 REVERSE LENGTH=580
Eucgr.H00564.1	12,64	142,25	102,24	81,75	36,62	AT1G68090.1 Symbols: ANNAT5, ANNS annexin 5 chr1:25519442-25520774 REVERSE LENGTH=316
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.A01107.1	4,38	0,83	0,72	0,97	1,44	AT2G32590.1 unknown protein
Eucgr.J01598.1	19,43	0,43	0,94	1,51	8,01	AT5G16720.1 Symbols: Protein of unknown function, DUF593
Eucgr.J01910.1	0,54	7,08	6,35	4,03	2,69	No hits found
Eucgr.F01907.1	4,45	0,65	0,58	0,77	1,77	AT1G75150.1 Symbols: unknown protein;
Eucgr.C03593.1	0,18	5,75	2,92	2,40	1,04	AT5G20610.1 Symbols: unknown protein;
Eucgr.K01424.1	9,34	0,39	0,50	1,09	2,76	AT4G28660.1 Symbols: PSB28 photosystem II reaction center PSB28 protein
Eucgr.C02424.1	10,48	0,92	2,11	3,12	6,60	AT5G57670.1 Symbols: Protein kinase superfamily protein chr5:23360531-23363694 REVERSE LENGTH=579
Eucgr.L03349.1	0,74	0,00	0,00	0,05	0,17	AT3G63470.1 Symbols: scpl40 serine carboxypeptidase-like 40 chr3:23438561-23440862 FORWARD LENGTH=502
Eucgr.A02688.1	9,29	0,07	0,53	0,38	2,14	AT5G52860.1 Symbols: ABC-2 type transporter family protein chr5:21419776-21421545 REVERSE LENGTH=589
Eucgr.I01972.1	6,97	0,28	0,69	0,81	1,90	AT3G01015.1 Symbols: TPX2 (targeting protein for Xklp2) protein family chr3:1798-4017 REVERSE LENGTH=488
Eucgr.I01041.1	8,13	0,05	0,04	0,05	0,84	AT2G22990.1 Symbols: SNG1, SCPL8 sinapoylglucose 1 chr2:9786393-9789925 FORWARD LENGTH=433
Eucgr.A01014.3	20,05	0,97	1,28	2,21	11,70	AT1G75250.1 Symbols: ATRL6, RSM3, RL6 RAD-like 6 chr1:28245073-28245453 REVERSE LENGTH=126
Eucgr.B03001.1	10,78	0,37	0,87	0,87	4,40	AT1G68360.1 Symbols: C2H2 and C2HC zinc fingers superfamily protein chr1:25621678-25622412 REVERSE LENGTH=244
Eucgr.I02165.1	7,24	0,37	0,64	1,20	2,81	AT1G53820.1 Symbols: RING/U-box superfamily protein chr1:20091491-20092423 FORWARD LENGTH=310
Eucgr.E02665.1	1,68	0,03	0,01	0,04	0,30	AT1G53210.1 Symbols: sodium/calcium exchanger family protein / calcium-binding EF hand family protein
Eucgr.E03762.2	0,26	9,63	10,32	6,79	1,51	AT4G05200.1 Symbols: CRK25 cysteine-rich RLK (RECEPTOR-like protein kinase) 25
Eucgr.F03057.1	0,25	6,74	5,20	3,01	0,86	AT3G11560.4 Symbols: LETM1-like protein chr3:3639553-3645506 FORWARD LENGTH=872
Eucgr.E01053.1	56,62	8,07	14,73	24,68	31,55	AT2G46770.1 Symbols: NST1, EMB2301, ANAC043 NAC (No Apical Meristem) domain transcriptional regulator superfamily protein
Eucgr.K02999.1	5,34	0,11	0,55	0,75	1,56	AT5G38690.1 Symbols: Zinc-finger domain of monoamine-oxidase A repressor R1 protein
Eucgr.J00337.1	0,01	4,30	2,62	1,34	0,06	AT4G27420.1 Symbols: ABC-2 type transporter family protein chr4:13712434-13714797 REVERSE LENGTH=638
Eucgr.E01674.1	3,01	0,39	1,02	0,85	1,69	AT4G08850.2 Symbols: Leucine-rich repeat receptor-like protein kinase family protein
Eucgr.C00887.2	6,39	16,34	16,37	15,88	12,10	AT5G56900.2 Symbols: Cwfl-like family protein / zinc finger (CCCH-type) family protein
Eucgr.J00100.1	121,16	6,53	8,13	11,85	46,17	AT5G58300.2 Symbols: Leucine-rich repeat protein kinase family protein
Eucgr.H01073.1	12,56	0,90	0,95	1,16	4,15	No hits found

Eucgr.G01011.1	1,43	0,02	0,13	0,29	0,62	AT5G10530.1 Symbols: Concanavalin A-like lectin protein kinase family protein
Eucgr.G02554.1	0,56	17,38	14,94	10,37	3,41	AT1G69850.1 Symbols: ATNRT1:2, NTL1, NRT1:2 nitrate transporter 1:2 chr1:26296945-26300407 REVERSE LENGTH=585
Eucgr.D00605.1	9,78	0,45	0,64	0,98	3,08	AT1G28340.1 Symbols: ATRLP4, RLP4 receptor like protein 4 chr1:9940175-9943252 FORWARD LENGTH=626
Eucgr.B00988.1	8,49	1,21	1,60	2,75	4,84	AT3G17950.1 Symbols: unknown protein
Eucgr.K03267.1	3,62	0,07	0,21	0,22	1,30	AT5G43420.1 Symbols: RING/U-box superfamily protein chr5:17451790-17452917 FORWARD LENGTH=375
Eucgr.B00599.1	8,10	0,19	0,43	0,79	3,02	AT5G08020.1 Symbols: ATRPA70B, RPA70B RPA70-kDa subunit B chr5:2572107-2574879 FORWARD LENGTH=604
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.B00551.1	35,81	2,17	4,26	5,92	17,08	AT1G18670.1 Symbols: IBS1 Protein kinase superfamily protein chr1:6427242-6430696 REVERSE LENGTH=709
Eucgr.B01491.1	5,55	0,12	0,36	0,58	2,22	AT3G54650.1 Symbols: FBL17 RNI-like superfamily protein chr3:20226004-20228882 REVERSE LENGTH=593
Eucgr.E02340.1	1,98	0,13	0,08	0,15	0,36	AT1G79740.1 Symbols: hAT transposon superfamily chr1:30004367-30006715 REVERSE LENGTH=651
Eucgr.F02385.1	5,66	0,70	0,96	0,87	3,11	AT1G06870.1 Symbols: Peptidase S24/S26A/S26B/S26C family protein chr1:2108832-2110642 FORWARD LENGTH=367
Eucgr.E01067.3	79,45	2,95	7,99	12,36	30,56	AT1G32860.1 Symbols: Glycosyl hydrolase superfamily protein chr1:11907308-11908803 REVERSE LENGTH=426
Eucgr.K03005.1	78,79	14,60	19,22	25,88	44,34	AT5G01090.1 Symbols: Concanavalin A-like lectin family protein chr5:33055-34116 FORWARD LENGTH=353
Eucgr.B00050.1	49,41	3,54	5,42	8,21	18,44	AT4G40080.1 Symbols: ENTH/ANTH/VHS superfamily protein chr4:18579371-18580542 FORWARD LENGTH=365
Eucgr.J00390.1	23,19	0,68	0,47	1,32	6,52	AT5G22880.1 Symbols: H2B, HTB2 histone B2 chr5:7652130-7652567 REVERSE LENGTH=145
Eucgr.B00944.1	341,64	3,14	8,91	12,41	90,76	AT1G73620.1 Symbols: Pathogenesis-related thaumatin superfamily protein
Eucgr.L00518.1	2,43	0,00	0,19	0,66	0,55	AT2G02220.1 Symbols: ATPSKR1, PSKR1 phytosulfokin receptor 1 chr2:584098-587124 REVERSE LENGTH=1008
Eucgr.L00984.1	3,73	0,25	0,85	1,07	1,57	No hits found
Eucgr.B01142.1	22,88	0,88	1,77	2,57	8,87	AT3G06130.2 Symbols: Heavy metal transport/detoxification superfamily protein
Eucgr.B02240.1	10,68	1,20	1,78	2,19	6,12	AT5G58600.1 Symbols: PMR5, TBL44 Plant protein of unknown function (DUF828)
Eucgr.L01545.1	6,34	0,00	0,49	0,82	1,37	AT5G27550.1 Symbols: P-loop containing nucleoside triphosphate hydrolases superfamily protein
Eucgr.C00641.1	21,35	84,20	73,78	62,26	47,57	AT5G26040.2 Symbols: HDA2 histone deacetylase 2 chr5:9099321-9101598 REVERSE LENGTH=387
Eucgr.B00321.1	0,32	0,02	0,00	0,03	0,07	AT3G48150.1 Symbols: APC8, CDC23 anaphase-promoting complex subunit 8
Eucgr.H01542.1	14,48	59,67	54,43	56,83	28,25	AT1G03360.1 Symbols: ATRRP4, RRP4 ribosomal RNA processing 4 chr1:824653-826179 FORWARD LENGTH=322
Eucgr.I01044.2	4,41	0,02	0,04	0,07	1,17	AT1G33540.1 Symbols: scpl18 serine carboxypeptidase-like 18 chr1:12162349-12164700 REVERSE LENGTH=446
Eucgr.D01652.1	48,33	5,47	3,89	5,24	13,63	AT5G59970.1 Symbols: Histone superfamily protein chr5:24146352-24146663 REVERSE LENGTH=103
Eucgr.J02063.1	71,85	5,51	6,10	9,94	33,73	No hits found
Eucgr.A02875.1	13,92	0,77	0,89	1,23	3,14	AT5G51600.1 Symbols: PLE, ATMAP65-3, MAP65-3 Microtubule associated protein (MAP65/ASE1) family protein
Eucgr.C00778.1	6,85	0,47	1,06	1,55	3,55	AT1G74960.3 Symbols: FAB1 fatty acid biosynthesis 1 chr1:28152564-28155948 REVERSE LENGTH=541
Eucgr.I01037.1	40,91	0,25	0,80	1,36	16,04	AT1G73270.1 Symbols: scpl6 serine carboxypeptidase-like 6 chr1:27549021-27552517 REVERSE LENGTH=452

Eucgr.A02902.1	129,32	3,58	9,45	11,21	44,80	AT5G61480.1 Symbols: PXY Leucine-rich repeat protein kinase family protein
Eucgr.C02505.1	9,97	37,74	33,87	37,48	23,55	AT2G24060.1 Symbols: Translation initiation factor 3 protein chr2:10229453-10231390 FORWARD LENGTH=312
Eucgr.B02268.1	2,01	0,00	0,10	0,19	0,55	AT2G29680.2 Symbols: CDC6, ATCDC6 cell division control 6 chr2:12689586-12692854 FORWARD LENGTH=508
Eucgr.D00303.1	33,81	91,21	90,91	113,92	75,66	AT1G53560.1 Symbols: Ribosomal protein L18ae family chr1:19984908-19985960 REVERSE LENGTH=153
Eucgr.F03082.1	4,20	0,09	0,20	0,20	0,95	No hits found
Eucgr.H04906.1	9,33	0,25	0,09	0,34	2,38	No hits found
Eucgr.D01657.1	22,49	2,42	3,36	5,01	11,02	AT1G32930.1 Symbols: Galactosyltransferase family protein chr1:11931980-11934399 REVERSE LENGTH=399
Eucgr.F02441.1	16,28	0,11	0,50	0,79	6,95	AT1G78060.1 Symbols: Glycosyl hydrolase family protein chr1:29349796-29352868 REVERSE LENGTH=767
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.E01240.1	20,81	3,31	4,90	5,37	12,01	AT1G31850.2 Symbols: S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
Eucgr.F03071.1	26,13	1,39	4,09	6,57	18,16	AT1G79760.1 Symbols: DTA4 downstream target of AGL15-4 chr1:30012054-30013042 FORWARD LENGTH=299
Eucgr.K00289.1	9,82	0,57	0,69	1,25	2,03	No hits found
Eucgr.E00787.1	106,55	5,93	6,54	10,79	18,83	AT2G46150.1 Symbols: Late embryogenesis abundant (LEA) hydroxyproline-rich glycoprotein family
Eucgr.F01793.1	113,75	0,78	1,54	3,30	23,64	AT1G42970.1 Symbols: GAPB glyceraldehyde-3-phosphate dehydrogenase B subunit
Eucgr.I01534.1	41,79	346,80	294,74	241,43	181,85	AT4G35300.5 Symbols: TMT2 tonoplast monosaccharide transporter2 chr4:16796432-16798332 REVERSE LENGTH=542
Eucgr.A02961.1	5,51	0,48	0,37	0,33	1,37	AT1G50240.2 Symbols: FU Protein kinase family protein with ARM repeat domain
Eucgr.C02984.1	8,40	0,14	0,20	0,70	1,62	AT3G16500.1 Symbols: PAP1, IAA26 phytochrome-associated protein 1 chr3:5612801-5614208 REVERSE LENGTH=269
Eucgr.A02093.1	4,34	0,48	0,42	0,38	1,38	AT1G54385.2 Symbols: ARM repeat superfamily protein chr1:20301289-20303048 REVERSE LENGTH=560
Eucgr.B03344.1	9,80	0,49	0,67	0,85	2,25	AT1G23790.1 Symbols: Plant protein of unknown function (DUF936) chr1:8410099-8411917 FORWARD LENGTH=518
Eucgr.F02669.1	45,30	4,39	6,46	6,83	21,01	AT1G14290.1 Symbols: SBH2 sphingoid base hydroxylase 2 chr1:4880307-4881784 REVERSE LENGTH=259
Eucgr.G03410.1	0,88	0,00	0,00	0,00	0,13	AT2G27660.1 Symbols: Cysteine/Histidine-rich C1 domain family protein chr2:11798694-11800850 REVERSE LENGTH=718
Eucgr.D02102.1	3,92	52,28	42,04	29,15	15,81	AT2G35700.1 Symbols: ATERF38, ERF38 ERF family protein 38 chr2:15005205-15005789 FORWARD LENGTH=194
Eucgr.G00453.1	18,24	5,14	4,28	4,15	11,06	AT3G18830.1 Symbols: ATPLT5, PMT5, ATPMT5 polyol/monosaccharide transporter 5
Eucgr.G02470.1	50,46	5,37	8,59	14,80	31,89	No hits found
Eucgr.G03393.1	274,46	0,82	7,50	13,54	89,26	AT3G20820.1 Symbols: Leucine-rich repeat (LRR) family protein chr3:7280930-7282027 FORWARD LENGTH=365
Eucgr.J02497.1	1,00	5,25	6,14	6,11	2,68	AT1G65770.1 Symbols: AMR1 ascorbic acid mannose pathway regulator 1
Eucgr.B03204.1	13,62	55,51	54,92	47,91	32,70	AT2G03220.1 Symbols: FT1, ATFUT1, ATFT1, MUR2 fucosyltransferase 1 chr2:970401-972353 REVERSE LENGTH=558
Eucgr.F03752.1	2,46	0,00	0,07	0,31	0,83	AT1G75710.1 Symbols: C2H2-like zinc finger protein chr1:28428806-28431128 FORWARD LENGTH=462
Eucgr.F04123.1	4,43	24,12	27,70	19,16	14,12	AT1G35660.1 Symbols: unknown protein;
Eucgr.K02028.1	8,68	1,99	1,26	1,40	4,76	AT4G18590.1 Symbols: Nucleic acid-binding, OB-fold-like protein chr4:10236524-10236940 FORWARD LENGTH=106

Eucgr.A02614.1	25,53	108,74	109,69	117,90	62,20	AT5G06260.1 Symbols: TLD-domain containing nucleolar protein chr5:1902755-1904835 REVERSE LENGTH=424
Eucgr.A02807.2	29,92	7,05	7,88	8,57	19,60	AT5G07240.1 Symbols: IQD24 IQ-domain 24 chr5:2272028-2274051 FORWARD LENGTH=401
Eucgr.G02888.1	2,51	0,28	0,23	0,25	1,09	AT5G13550.1 Symbols: SULTR4;1 sulfate transporter 4.1 chr5:4355412-4359490 REVERSE LENGTH=685
Eucgr.D01969.1	10,57	0,00	0,03	0,07	1,75	AT4G22790.1 Symbols: MATE efflux family protein chr4:11975153-11976628 REVERSE LENGTH=491
Eucgr.B00460.1	35,05	2,61	4,89	9,65	16,17	AT5G08139.1 Symbols: RING/U-box superfamily protein chr5:2616487-2617617 FORWARD LENGTH=376
Eucgr.C01259.2	28,79	0,38	2,46	4,27	19,92	AT2G46225.2 Symbols: ABIL1 ABI-1-like 1 chr2:18982113-18984074 FORWARD LENGTH=329
Eucgr.C03720.3	12,49	64,35	70,19	66,15	36,01	AT5G19740.1 Symbols: Peptidase M28 family protein chr5:6673986-6676767 FORWARD LENGTH=681
Eucgr.F04353.1	31,13	7,91	6,54	7,72	14,84	AT1G21830.1 Symbols: unknown protein;
Eucgr.B02400.1	8,24	1,05	1,31	1,44	4,26	AT5G59790.1 Symbols: Domain of unknown function (DUF966) chr5:24090950-24092804 FORWARD LENGTH=423
Eucgr.K02419.1	12,75	57,17	49,61	53,84	27,59	AT5G03560.1 Symbols: Tetratricopeptide repeat (TPR)-like superfamily protein
GENE ID	FPKM					Annotation

	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.B00609.1	5,22	0,40	0,87	1,10	2,48	AT1G74160.1 Symbols: unknown protein
Eucgr.C00618.1	0,50	38,75	24,66	18,36	2,85	AT2G25890.1 Symbols: Oleosin family protein chr2:11037435-11037884 FORWARD LENGTH=149
Eucgr.F04267.1	0,80	0,04	0,13	0,10	0,28	AT4G25980.1 Symbols: Peroxidase superfamily protein chr4:13189393-13191507 FORWARD LENGTH=371
Eucgr.G01567.1	10,72	58,84	37,89	28,81	20,54	AT5G67210.1 Symbols: Protein of unknown function (DUF579) chr5:26819019-26819972 FORWARD LENGTH=317
Eucgr.K01750.1	14,82	1,65	3,32	4,93	8,91	AT2G41820.1 Symbols: Leucine-rich repeat protein kinase family protein
Eucgr.D02018.1	15,80	0,35	1,46	1,50	5,99	AT4G12390.1 Symbols: PME1 pectin methylesterase inhibitor 1 chr4:7336553-7337173 FORWARD LENGTH=206
Eucgr.G02689.1	8,40	50,00	41,57	43,99	26,38	AT2G25100.1 Symbols: Polynucleotidyl transferase, ribonuclease H-like superfamily protein
Eucgr.H03461.1	0,03	5,99	4,21	1,81	0,31	AT3G22490.1 Symbols: Seed maturation protein chr3:7969785-7970738 REVERSE LENGTH=262
Eucgr.L00712.1	7,42	0,38	0,48	0,71	2,38	AT4G19840.1 Symbols: ATPP2-A1, ATPP2A-1, PP2-A1 phloem protein 2-A1
Eucgr.D00432.1	30,00	1,79	4,78	8,24	14,94	AT1G17030.1 Symbols: unknown protein;
Eucgr.C02127.1	4,43	0,00	0,05	0,21	0,44	AT4G31330.1 Symbols: Protein of unknown function, DUF599 chr4:15202354-15203151 FORWARD LENGTH=239
Eucgr.K01221.2	23,90	2,62	4,73	7,69	14,67	AT3G59110.1 Symbols: Protein kinase superfamily protein chr3:21855673-21857847 FORWARD LENGTH=512
Eucgr.K03427.1	21,62	0,93	1,17	1,68	6,20	AT4G03100.1 Symbols: Rho GTPase activating protein with PAK-box/P21-Rho-binding domain
Eucgr.B03400.1	16,70	0,75	1,37	1,78	7,96	AT1G69910.1 Symbols: Protein kinase superfamily protein chr1:26330166-26332076 FORWARD LENGTH=636
Eucgr.H04811.1	6,24	0,71	0,65	0,83	2,40	AT5G10110.1 Symbols: unknown protein;
Eucgr.F03619.1	24,87	0,35	1,11	1,61	10,18	AT2G33570.1 Symbols: Domain of unknown function (DUF23) chr2:14217361-14219043 FORWARD LENGTH=496
Eucgr.H03555.1	15,82	0,81	1,89	2,64	8,84	AT1G47480.1 Symbols: alpha/beta-Hydrolases superfamily protein chr1:17417623-17419296 FORWARD LENGTH=314
Eucgr.D01977.1	20,18	62,67	46,61	52,79	34,33	AT1G12340.1 Symbols: Cornichon family protein chr1:4197396-4198032 FORWARD LENGTH=129
Eucgr.I00628.1	137,65	23,23	26,89	21,21	51,74	AT4G37870.1 Symbols: PCK1, PEPCCK phosphoenolpyruvate carboxykinase 1

Eucgr.C02915.1	0,61	0,00	0,02	0,00	0,17	AT4G28890.1 Symbols: RING/U-box superfamily protein chr4:14256437-14257735 REVERSE LENGTH=432
Eucgr.A02622.1	22,15	3,41	5,48	8,14	14,02	AT3G11590.1 Symbols: unknown protein;
Eucgr.E00596.1	22,69	1,22	4,29	7,27	14,35	AT4G11450.1 Symbols: Protein of unknown function (DUF3527) chr4:6959065-6961223 REVERSE LENGTH=694
Eucgr.G02657.1	13,90	3,76	5,04	5,27	9,26	AT5G04885.1 Symbols: Glycosyl hydrolase family protein chr5:1423369-1426628 FORWARD LENGTH=665
Eucgr.E01250.1	38,77	3,72	4,68	7,66	21,15	AT1G32100.1 Symbols: ATPRR1, PRR1 pinoreinol reductase 1 chr1:11546472-11547953 REVERSE LENGTH=317
Eucgr.F02355.1	8,00	0,84	0,74	0,60	2,71	AT4G33400.1 Symbols: Vacuolar import/degradation, Vid27-related protein
Eucgr.C03813.1	25,83	0,97	1,63	1,95	12,67	AT4G28950.1 Symbols: ARAC7, ATROP9, ATRAC7, RAC7, ROP9 RHO-related protein from plants 9
Eucgr.C00897.1	14,21	0,15	0,62	1,14	8,51	AT4G29720.1 Symbols: ATPAO5, PAO5 polyamine oxidase 5 chr4:14553456-14555057 REVERSE LENGTH=533
Eucgr.G02614.1	16,98	56,10	45,29	57,33	33,72	AT5G63670.1 Symbols: SPT42 SPT4 homolog 2 chr5:25488012-25489247 FORWARD LENGTH=116
Eucgr.G00958.1	31,51	9,46	10,73	12,03	19,74	AT2G07050.1 Symbols: CAS1 cycloartenol synthase 1 chr2:2924629-2930295 FORWARD LENGTH=759
Eucgr.C03290.1	3,32	0,04	0,12	0,11	0,81	AT5G20740.1 Symbols: Plant invertase/pectin methylesterase inhibitor superfamily protein
Eucgr.I01487.1	0,83	0,00	0,00	0,00	0,21	AT1G29420.1 Symbols: SAUR-like auxin-responsive protein family chr1:10301483-10301908 REVERSE LENGTH=141
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.C00584.1	12,41	1,03	1,94	2,62	8,01	AT1G27600.2 Symbols: IRX9-L, I9H Nucleotide-diphospho-sugar transferases superfamily protein
Eucgr.H01480.1	10,18	0,38	0,60	0,98	3,18	AT4G02800.1 Symbols: unknown protein;
Eucgr.K01099.1	14,76	131,81	86,85	75,72	59,12	AT2G31090.1 Symbols: unknown protein;
Eucgr.G01371.1	72,90	1,90	2,04	2,42	31,30	AT5G67260.1 Symbols: CYCD3;2 CYCLIN D3;2 chr5:26836313-26837665 FORWARD LENGTH=367
Eucgr.I02121.1	19,90	1,50	2,61	4,64	9,95	AT5G38560.1 Symbols: Protein kinase superfamily protein chr5:15439844-15443007 FORWARD LENGTH=681
Eucgr.I02501.1	7,81	1,31	1,75	2,21	5,01	AT3G22960.1 Symbols: PKP1, PKP-ALPHA Pyruvate kinase family protein chr3:8139369-8141771 FORWARD LENGTH=596
Eucgr.F01814.1	28,79	3,70	3,66	4,57	10,92	AT1G20930.1 Symbols: CDKB2;2 cyclin-dependent kinase B2;2 chr1:7292752-7294664 REVERSE LENGTH=315
Eucgr.J02626.1	11,01	1,99	3,04	3,28	5,86	AT5G25170.1 Symbols: PPPDE putative thiol peptidase family protein chr5:8693257-8694438 FORWARD LENGTH=218
Eucgr.K02806.1	25,05	1,53	4,51	7,19	13,06	AT5G26660.1 Symbols: ATMYB86, MYB86 myb domain protein 86 chr5:9331775-9333044 REVERSE LENGTH=352
Eucgr.H00370.1	124,23	5,85	6,17	6,60	26,26	AT2G43290.1 Symbols: MSS3 Calcium-binding EF-hand family protein chr2:17991308-17991955 REVERSE LENGTH=215
Eucgr.J01835.1	20,45	2,30	3,74	3,92	11,45	AT1G60630.1 Symbols: Leucine-rich repeat protein kinase family protein chr1:22334754-22336785 REVERSE LENGTH=652
Eucgr.H04041.1	1,64	0,21	0,12	0,19	0,82	AT3G17030.1 Symbols: Nucleic acid-binding proteins superfamily chr3:5804509-5808462 REVERSE LENGTH=668
Eucgr.D01080.1	6,28	0,25	0,65	0,89	2,36	AT5G17670.1 Symbols: alpha/beta-Hydrolases superfamily protein chr5:5821082-5822600 FORWARD LENGTH=309
Eucgr.H00778.1	2,36	0,19	0,19	0,15	0,81	AT3G25100.1 Symbols: CDC45 cell division cycle 45 chr3:9144292-9146082 FORWARD LENGTH=596
Eucgr.K03502.1	3,70	0,30	0,21	0,33	1,06	AT2G47560.1 Symbols: RING/U-box superfamily protein chr2:19511934-19512617 REVERSE LENGTH=227
Eucgr.G02798.1	22,08	0,71	0,34	0,26	4,04	AT3G54770.1 Symbols: RNA-binding (RRM/RBD/RNP motifs) family protein
Eucgr.A02602.2	4,29	0,73	0,82	0,92	1,96	AT2G36010.1 Symbols: E2F3, ATE2FA E2F transcription factor 3 chr2:15119688-15122893 FORWARD LENGTH=483

Eucgr.F04077.1	0,56	0,01	0,01	0,02	0,12	AT3G47570.1 Symbols: Leucine-rich repeat protein kinase family protein
Eucgr.G00446.1	23,59	3,61	5,87	10,39	14,73	AT4G34500.1 Symbols: Protein kinase superfamily protein chr4:16488005-16490792 REVERSE LENGTH=437
Eucgr.J03181.2	11,88	0,27	0,79	1,67	6,76	AT3G20260.1 Symbols: Protein of unknown function (DUF1666) chr3:7064190-7065751 REVERSE LENGTH=437
Eucgr.C04266.1	26,09	1317,97	1007,17	544,49	193,24	AT5G20250.4 Symbols: DIN10 Raffinose synthase family protein chr5:6833730-6836635 FORWARD LENGTH=844
Eucgr.I00199.1	3,29	0,98	0,83	0,63	2,04	AT4G33440.1 Symbols: Pectin lyase-like superfamily protein chr4:16092020-16094732 FORWARD LENGTH=475
Eucgr.B02583.1	2,50	0,15	0,18	0,22	1,22	AT5G23400.1 Symbols: Leucine-rich repeat (LRR) family protein chr5:7880603-7882372 FORWARD LENGTH=589
Eucgr.I00804.1	15,40	1,60	3,13	5,42	9,10	AT3G19420.1 Symbols: ATPEN2, PEN2 PTEN 2 chr3:6731824-6735354 FORWARD LENGTH=611
Eucgr.B03781.1	24,04	0,04	0,86	1,39	9,77	AT5G21105.1 Symbols: Plant L-ascorbate oxidase chr5:7172727-7177409 FORWARD LENGTH=588
Eucgr.F04121.1	58,25	2,56	4,77	3,79	29,98	AT1G71970.1 Symbols: unknown protein;
Eucgr.H01238.1	1,46	8,25	10,40	11,71	5,19	AT4G36790.1 Symbols: Major facilitator superfamily protein chr4:17336360-17338304 FORWARD LENGTH=489
Eucgr.D00086.1	5,95	0,22	0,63	1,05	2,53	AT4G20840.1 Symbols: FAD-binding Berberine family protein chr4:11157916-11159535 FORWARD LENGTH=539
Eucgr.J00152.1	7,28	0,10	0,32	0,98	2,78	AT3G28050.1 Symbols: nodulin MtN21 /EamA-like transporter family protein
Eucgr.J03121.1	12,12	130,82	102,99	110,44	30,65	AT1G80920.1 Symbols: J8 Chaperone DnaJ-domain superfamily protein chr1:30403863-30404549 REVERSE LENGTH=163
Eucgr.B03123.1	3,73	0,63	0,61	0,54	1,47	AT3G45070.1 Symbols: P-loop containing nucleoside triphosphate hydrolases superfamily protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H05115.1	149,35	0,20	0,11	0,27	2,51	AT5G63850.1 Symbols: AAP4 amino acid permease 4 chr5:25551494-25553374 FORWARD LENGTH=466
Eucgr.G02452.1	21,00	1,74	2,89	4,40	11,03	AT1G18650.1 Symbols: PDCB3 plasmodesmata callose-binding protein 3 chr1:6419036-6420413 REVERSE LENGTH=184
Eucgr.J01238.1	4,12	217,96	126,49	113,37	21,22	AT5G02220.1 Symbols: unknown protein;
Eucgr.J00928.1	1,04	69,46	45,49	33,12	5,61	AT5G60680.1 Symbols: Protein of unknown function, DUF584 chr5:24386247-24386738 FORWARD LENGTH=163
Eucgr.D00683.1	12,88	0,48	1,07	1,27	3,31	AT1G11090.1 Symbols: alpha/beta-Hydrolases superfamily protein chr1:3702655-3703629 FORWARD LENGTH=324
Eucgr.G01236.1	0,07	0,60	1,10	0,40	0,42	AT3G18990.1 Symbols: VRN1, REM39 AP2/B3-like transcriptional factor family protein
Eucgr.J00232.1	3,18	28,72	28,58	31,82	10,29	AT5G05700.1 Symbols: ATE1, DLS1, ATATE1 arginine-tRNA protein transferase 1
Eucgr.F04216.1	4,65	0,38	1,33	1,70	2,82	AT5G64740.1 Symbols: CESA6, IXR2, E112, PRC1 cellulose synthase 6 chr5:25881555-25886333 FORWARD LENGTH=1084
Eucgr.I01414.1	139,72	25,94	32,98	41,76	80,52	AT1G75840.1 Symbols: ARAC5, ATGP3, ROP4, ATROP4 RAC-like GTP binding protein 5
Eucgr.A02452.1	6,84	0,27	0,57	0,83	1,82	No hits found
Eucgr.G01455.1	5,99	0,26	0,30	0,29	1,55	AT3G19184.1 Symbols: AP2/B3-like transcriptional factor family protein chr3:6637555-6639035 FORWARD LENGTH=277
Eucgr.G01601.1	5,84	0,00	0,02	0,00	1,03	AT4G13440.1 Symbols: Calcium-binding EF-hand family protein chr4:7810215-7810679 FORWARD LENGTH=154
Eucgr.F02674.1	7,13	0,28	0,27	0,42	2,90	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.C02185.1	31,19	0,49	0,36	0,65	7,61	AT1G73270.1 Symbols: scpl6 serine carboxypeptidase-like 6 chr1:27549021-27552517 REVERSE LENGTH=452
Eucgr.K02684.1	0,94	59,52	31,31	17,24	6,17	AT5G23660.1 Symbols: MTN3, SWEET12, AtSWEET12 homolog of Medicago truncatula MTN3

Eucgr.K02629.1	0,50	16,38	15,25	13,46	3,42	AT5G23680.1 Symbols: Sterile alpha motif (SAM) domain-containing protein
Eucgr.F03763.1	0,73	0,00	0,02	0,11	0,26	AT5G08640.2 Symbols: FLS1 flavonol synthase 1 chr5:2804009-2805175 FORWARD LENGTH=336
Eucgr.F01637.1	43,10	5,05	6,51	7,10	25,95	AT1G19920.1 Symbols: APS2, ASA1 Pseudouridine synthase/archaeosine transglycosylase-like family protein
Eucgr.I02055.1	3,60	0,19	0,30	0,17	2,00	AT3G02230.1 Symbols: RGP1, ATRGP1 reversibly glycosylated polypeptide 1 chr3:415463-417304 FORWARD LENGTH=357
Eucgr.C01038.1	3,90	0,24	0,40	0,49	1,46	AT3G18830.1 Symbols: ATPLT5, PMT5, ATPMT5 polyol/monosaccharide transporter 5
Eucgr.F03951.1	4,70	0,00	0,03	0,12	1,68	AT5G10970.1 Symbols: C2H2 and C2HC zinc fingers superfamily protein chr5:3469268-3470086 FORWARD LENGTH=272
Eucgr.E02613.1	0,00	3,87	2,70	1,43	0,08	AT1G16770.1 Symbols: unknown protein;
Eucgr.B02320.1	0,00	3,18	1,89	0,55	0,17	No hits found
Eucgr.C00635.1	3,82	0,61	0,64	1,09	2,15	AT5G26010.1 Symbols: Protein phosphatase 2C family protein chr5:9085512-9087372 REVERSE LENGTH=331
Eucgr.F01221.1	42,25	141,23	147,83	152,96	85,38	AT1G53280.1 Symbols: Class I glutamine amidotransferase-like superfamily protein
Eucgr.F00754.1	1,41	0,00	0,10	0,05	0,51	No hits found
Eucgr.H04911.1	16,40	0,46	0,83	0,93	4,06	No hits found
Eucgr.F04010.1	7,51	0,11	0,35	0,47	3,06	AT4G31590.1 Symbols: ATCSLC05, CSLC05, ATCSLC5, CSLCS Cellulose-synthase-like C5
Eucgr.H02961.1	0,18	7,69	3,40	1,03	0,81	AT5G55240.1 Symbols: ATPXG2 ARABIDOPSIS THALIANA PEROXYGENASE 2
Eucgr.D01317.1	27,91	2,38	4,44	6,21	16,79	AT1G61900.1 Symbols: unknown protein
Eucgr.I01535.1	19,28	108,99	123,16	78,86	61,15	AT3G51490.2 Symbols: TMT3 tonoplast monosaccharide transporter3 chr5:19105018-19107562 REVERSE LENGTH=737
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.B03167.1	2,47	0,13	0,22	0,15	0,70	AT5G61210.1 Symbols: SNAP33, ATSNAP33, SNP33, ATSNAP33B soluble N-ethylmaleimide-sensitive factor
Eucgr.E00809.1	194,46	4,98	18,64	36,55	66,93	AT4G23690.1 Symbols: Disease resistance-responsive (dirigent-like protein) family protein
Eucgr.F03961.1	3,47	13,71	15,61	13,14	7,88	AT2G07680.1 Symbols: ATMRP11, MRP11, ABCC13 multidrug resistance-associated protein 11
Eucgr.H00187.1	0,35	0,00	0,01	0,04	0,11	AT4G23990.1 Symbols: ATCSLG3, CSLG3 cellulose synthase like G3 chr4:12456491-12460498 FORWARD LENGTH=751
Eucgr.K00318.2	62,82	6,58	10,43	15,19	31,35	AT5G53320.1 Symbols: Leucine-rich repeat protein kinase family protein chr5:21636453-21638337 REVERSE LENGTH=601
Eucgr.G03039.1	0,61	15,02	13,01	13,13	2,67	AT1G07280.3 Symbols: Tetratricopeptide repeat (TPR)-like superfamily protein
Eucgr.J00985.1	370,20	7,20	9,95	17,02	101,82	AT3G54260.1 Symbols: TBL36 TRICHOME BIREFRINGENCE-LIKE 36 chr3:20085097-20086745 REVERSE LENGTH=379
Eucgr.B00868.1	59,93	3,14	6,95	12,15	32,19	AT3G18050.1 Symbols: unknown protein;
Eucgr.K02307.1	28,04	0,00	0,03	0,03	1,42	AT1G73220.1 Symbols: AtOCT1, 1-Oct organic cation/carnitine transporter1
Eucgr.H04662.1	18,11	1,66	3,61	5,89	11,65	AT1G67560.1 Symbols: LOX6 PLAT/LH2 domain-containing lipoxygenase family protein
Eucgr.J02055.1	51,01	4,70	4,54	8,98	21,07	AT4G40030.2 Symbols: Histone superfamily protein chr4:18555840-18556827 REVERSE LENGTH=164
Eucgr.H01659.1	0,01	6,56	3,62	1,26	0,09	AT4G34410.1 Symbols: RRTF1 redox responsive transcription factor 1 chr4:16451992-16452798 FORWARD LENGTH=268
Eucgr.L01090.1	2,86	0,17	0,62	0,86	1,55	AT4G36890.1 Symbols: IRX14 Nucleotide-diphospho-sugar transferases superfamily protein

Eucgr.F03342.1	15,40	2,69	3,79	4,30	9,01	AT5G54400.1 Symbols: S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
Eucgr.D01942.1	17,85	0,09	0,73	1,24	7,34	AT4G22900.1 Symbols: Protein of unknown function (DUF1191) chr4:12010221-12011252 FORWARD LENGTH=343
Eucgr.H01211.1	5,26	0,65	0,67	1,50	1,87	AT1G48910.1 Symbols: YUC10 Flavin-containing monooxygenase family protein
Eucgr.J01214.2	27,51	0,61	2,09	2,85	13,06	AT3G28860.1 Symbols: ATMDR1, ATMDR11, PGP19, MDR11, MDR1, ATPGP19, ABCB19, ATABC19 ATP binding cassette subfamily B19
Eucgr.F02624.1	27,85	1,84	2,97	3,43	13,17	AT5G54160.1 Symbols: ATOMT1, OMT1 O-methyltransferase 1 chr5:21982075-21984167 FORWARD LENGTH=363
Eucgr.A01822.1	23,85	0,24	0,74	1,63	10,18	AT4G28310.1 Symbols: unknown protein
Eucgr.L02974.1	6,42	0,24	0,21	0,55	2,80	AT3G07880.1 Symbols: SCN1 Immunoglobulin E-set superfamily protein chr3:2514175-2515544 FORWARD LENGTH=240
Eucgr.F01185.1	3,31	38,69	29,25	22,48	7,98	AT5G54080.2 Symbols: HGO homogentisate 1,2-dioxygenase chr5:21945920-21948070 FORWARD LENGTH=461
Eucgr.B02771.1	16,28	0,60	1,42	1,49	4,75	AT1G58170.1 Symbols: Disease resistance-responsive (dirigent-like protein) family protein
Eucgr.I00910.1	4,61	0,17	0,17	0,32	1,58	AT5G67090.1 Symbols: Subtilisin-like serine endopeptidase family protein
Eucgr.K01836.1	0,22	140,01	103,05	43,74	1,07	AT5G06760.1 Symbols: LEA4-5 Late Embryogenesis Abundant 4-5 chr5:2089754-2090313 REVERSE LENGTH=158
Eucgr.H04042.1	8,92	0,60	0,92	1,36	5,22	AT3G17040.1 Symbols: HCF107 high chlorophyll fluorescent 107 chr3:5809378-5812605 REVERSE LENGTH=652
Eucgr.F00115.1	21,48	0,63	0,63	0,79	6,99	AT3G49940.1 Symbols: LBD38 LOB domain-containing protein 38 chr3:18514465-18515302 FORWARD LENGTH=247
Eucgr.H02335.1	7,53	0,15	0,46	1,08	5,03	AT4G37750.1 Symbols: ANT, DRG, CKC, CKC1 Integrase-type DNA-binding superfamily protein
Eucgr.H00130.1	21,52	3,91	4,30	6,49	14,76	AT1G04520.1 Symbols: PDLP2 plasmodesmata-located protein 2 chr1:1231852-1233787 FORWARD LENGTH=307
Eucgr.L03019.1	10,25	235,23	141,29	102,84	25,65	No hits found
Eucgr.C03048.1	1,24	0,00	0,01	0,09	0,31	AT4G26150.1 Symbols: CGA1, GATA22, GNL cytokinin-responsive gata factor 1
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.G03115.1	3,08	0,00	0,03	0,03	0,62	AT2G40435.1 Symbols: unknown protein
Eucgr.C02213.1	1,49	20,95	15,36	12,22	5,48	No hits found
Eucgr.H04937.1	13,67	0,02	0,15	0,21	4,66	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.C02874.1	158,39	1402,30	1448,07	1241,28	372,50	AT5G25560.2 Symbols: CHY-type/CTCHY-type/RING-type Zinc finger protein
Eucgr.I02220.2	27,62	84,32	76,11	78,92	49,88	AT4G35510.1 Symbols: unknown protein;
Eucgr.K02439.1	45,53	0,30	1,32	1,18	13,64	No hits found
Eucgr.C00963.1	1,30	23,61	22,96	34,80	2,98	AT5G56550.1 Symbols: OXS3, ATOXS3 oxidative stress 3 chr5:22895847-22896502 REVERSE LENGTH=172
Eucgr.H02314.1	34,55	1,35	5,36	11,28	18,37	AT5G67620.1 Symbols: unknown protein;
Eucgr.B00846.1	4,78	0,00	0,05	0,10	1,19	AT1G16390.1 Symbols: ATOCT3, 3-Oct organic cation/carnitine transporter 3 chr1:5602921-5604477 FORWARD LENGTH=518
Eucgr.G01751.1	4,83	0,23	0,42	1,17	2,23	AT4G34860.2 Symbols: Plant neutral invertase family protein chr4:16609955-16612019 REVERSE LENGTH=571
Eucgr.D01066.2	13,54	0,35	0,26	1,08	4,77	AT1G54730.2 Symbols: Major facilitator superfamily protein chr1:20424471-20429978 FORWARD LENGTH=470
Eucgr.A02407.1	13,58	4,89	4,17	5,41	8,98	AT3G12170.1 Symbols: Chaperone DnaJ-domain superfamily protein chr3:3881021-3882655 FORWARD LENGTH=262

Eucgr.D01021.1	4,04	0,52	0,34	0,41	1,37	AT3G25690.3 Symbols: CHUP1 Hydroxyproline-rich glycoprotein family protein
Eucgr.E03592.1	10,80	0,87	0,79	0,72	3,77	AT4G21270.1 Symbols: ATK1, KATA, KATAP kinesin 1 chr4:11329579-11333884 REVERSE LENGTH=793
Eucgr.K00540.1	0,31	4,88	2,14	1,31	1,08	AT1G28200.1 Symbols: FIP1 FH interacting protein 1 chr1:9850395-9852300 REVERSE LENGTH=259
Eucgr.J03010.1	1,00	0,00	0,03	0,00	0,34	AT1G12663.1 Symbols: Predicted to encode a PR (pathogenesis-related) protein.
Eucgr.H04817.1	8,33	0,08	0,15	0,58	1,43	No hits found
Eucgr.K00711.1	40,24	2,52	3,47	3,51	21,20	AT2G04780.2 Symbols: FLA7 FASCICLIN-like arabinogalactan 7 chr2:1677488-1678252 FORWARD LENGTH=254
Eucgr.B00946.1	52,67	1,74	2,12	2,74	11,37	AT3G18000.1 Symbols: NMT1, XPL1, PEAMT S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
Eucgr.J00336.1	14,73	0,87	1,28	1,61	8,15	AT5G23420.1 Symbols: HMGB6 high-mobility group box 6 chr5:7888712-7890111 REVERSE LENGTH=241
Eucgr.C03199.1	1532,75	66,91	100,38	132,04	213,15	AT3G43190.1 Symbols: SUS4, ATSUS4 sucrose synthase 4 chr3:15179204-15182577 REVERSE LENGTH=808
Eucgr.J02706.1	1,92	0,22	0,14	0,16	0,61	AT3G21340.1 Symbols: Leucine-rich repeat protein kinase family protein chr3:7511848-7515937 REVERSE LENGTH=899
Eucgr.K01509.1	0,01	0,74	0,64	0,62	0,09	AT5G13530.2 Symbols: KEG protein kinases;ubiquitin-protein ligases chr5:4345618-4354369 FORWARD LENGTH=1624
Eucgr.B01598.1	2,74	0,05	0,07	0,13	1,07	AT3G06880.2 Symbols: Transducin/WD40 repeat-like superfamily protein chr3:2170516-2175686 REVERSE LENGTH=1264
Eucgr.F01224.1	25,24	5,69	7,57	9,15	18,62	AT1G53290.1 Symbols: Galactosyltransferase family protein chr1:19871353-19873251 FORWARD LENGTH=345
Eucgr.A02891.1	173,15	26,43	29,77	40,45	84,89	AT5G62390.1 Symbols: ATBAG7, BAG7 BCL-2-associated athanogene 7 chr5:25052377-25054170 REVERSE LENGTH=446
Eucgr.J00806.1	39,40	2,72	3,85	4,81	15,48	AT4G40030.2 Symbols: Histone superfamily protein chr4:18555840-18556827 REVERSE LENGTH=164
Eucgr.B02618.1	4,68	0,26	0,46	0,52	1,43	AT5G13770.1 Symbols: Pentatricopeptide repeat (PPR-like) superfamily protein
Eucgr.J02473.1	8,98	0,98	1,05	1,19	4,88	AT4G35930.1 Symbols: F-box family protein chr4:17019482-17020744 FORWARD LENGTH=321
Eucgr.B03097.1	23,63	1,21	2,42	4,46	13,02	AT1G13600.1 Symbols: AtbZIP58, bZIP58 basic leucine-zipper 58 chr1:4650787-4651377 REVERSE LENGTH=196
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H00477.2	0,92	7,81	5,01	4,89	2,28	AT1G60730.2 Symbols: NAD(P)-linked oxidoreductase superfamily protein
Eucgr.D00242.1	5,10	0,16	0,38	0,49	2,13	AT4G21070.1 Symbols: ATBRCA1, BRCA1 breast cancer susceptibility1 chr4:11248174-11252633 FORWARD LENGTH=941
Eucgr.I00230.1	22,65	0,47	0,71	1,18	9,99	AT5G64660.1 Symbols: ATCMPG2, CMPG2 CYS, MET, PRO, and GLY protein 2
Eucgr.E00451.1	76,23	1,41	2,25	4,56	23,37	AT4G22570.1 Symbols: APT3 adenine phosphoribosyl transferase 3 chr4:11882310-11885250 REVERSE LENGTH=183
Eucgr.A00251.1	5,80	13,10	15,00	16,71	9,66	AT1G54380.1 Symbols: spliceosome protein-related chr1:20298287-20300899 REVERSE LENGTH=515
Eucgr.F00117.1	12,69	136,40	99,45	101,62	35,23	AT5G43150.1 Symbols: unknown protein
Eucgr.I01724.1	15,71	1,14	2,53	3,07	9,24	AT5G14430.1 Symbols: S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
Eucgr.B03932.1	29,45	0,53	1,78	3,44	15,22	AT5G22930.1 Symbols: Protein of unknown function (DUF1635) chr5:7668229-7669315 REVERSE LENGTH=238
Eucgr.K02904.1	1,98	7,48	10,54	8,73	5,68	AT1G74780.1 Symbols: Nodulin-like / Major Facilitator Superfamily protein
Eucgr.K01906.1	1,50	10,21	11,04	11,74	3,25	AT3G11900.1 Symbols: ANT1 aromatic and neutral transporter 1 chr3:3758523-3760103 FORWARD LENGTH=432
Eucgr.K00555.1	11,03	0,48	0,80	0,96	4,23	AT4G01680.3 Symbols: MYB55 myb domain protein 55 chr4:716401-717415 REVERSE LENGTH=241

Eucgr.I02367.1	8,30	0,45	0,78	0,58	3,89	AT4G36180.1 Symbols: Leucine-rich receptor-like protein kinase family protein
Eucgr.F01871.2	9,90	25,92	27,07	24,94	17,41	AT4G39170.1 Symbols: Sec14p-like phosphatidylinositol transfer family protein
Eucgr.H00932.1	6,18	17,51	17,06	16,40	10,28	AT1G11320.1 Symbols: unknown protein
Eucgr.F02126.1	15,99	81,63	67,38	51,81	31,20	AT5G10940.2 Symbols: transducin family protein / WD-40 repeat family protein
Eucgr.J00665.1	5,19	62,07	49,09	40,77	13,36	AT5G54620.1 Symbols: Ankyrin repeat family protein chr5:22187761-22189746 REVERSE LENGTH=431
Eucgr.C03057.1	2,25	0,18	0,19	0,21	0,72	AT2G26180.1 Symbols: IQD6 IQ-domain 6 chr2:11143433-11144982 REVERSE LENGTH=416
Eucgr.D00880.2	46,25	3,29	4,19	6,79	27,71	AT5G54530.1 Symbols: Protein of unknown function, DUF538 chr5:22152781-22154201 FORWARD LENGTH=161
Eucgr.J02310.1	0,80	6,66	5,90	2,79	1,78	AT5G40840.2 Symbols: SYN2 Rad21/Rec8-like family protein chr5:16359611-16363722 REVERSE LENGTH=810
Eucgr.D00630.1	6,01	0,44	0,87	1,34	3,37	AT3G15390.1 Symbols: SDE5 silencing defective 5 chr3:5196689-5199531 REVERSE LENGTH=490
Eucgr.F02356.1	1,39	37,24	36,27	28,89	5,34	No hits found
Eucgr.I00289.1	61,48	8,05	14,85	22,70	40,70	AT1G43130.1 Symbols: LCV2 like COV 2 chr1:16228661-16231158 REVERSE LENGTH=261
Eucgr.A01591.1	0,52	0,02	0,10	0,04	0,20	AT2G36660.1 Symbols: PAB7 poly(A) binding protein 7 chr2:15361476-15364398 REVERSE LENGTH=609
Eucgr.A01425.1	40,21	145,93	147,06	122,53	77,73	AT5G24400.1 Symbols: EMB2024, PGL3 NagB/RpiA/CoA transferase-like superfamily protein
Eucgr.B03382.1	32,12	1,67	3,57	5,27	16,24	AT1G27210.1 Symbols: ARM repeat superfamily protein chr1:9455669-9457917 REVERSE LENGTH=625
Eucgr.H01670.1	1,88	0,07	0,08	0,10	0,56	AT5G42930.1 Symbols: alpha/beta-Hydrolases superfamily protein chr5:17210738-17214152 REVERSE LENGTH=467
Eucgr.C03762.1	0,30	4,85	4,09	2,70	0,94	AT2G19900.1 Symbols: ATNADP-ME1, NADP-ME1 NADP-malic enzyme 1 chr2:8592106-8595403 REVERSE LENGTH=581
Eucgr.A01776.1	5,58	0,43	0,64	0,75	2,38	AT4G28230.1 Symbols: unknown protein;
Eucgr.E01169.1	69,42	4,13	5,46	6,43	17,75	AT4G19420.1 Symbols: Pectinacetylsterase family protein chr4:10587504-10589807 REVERSE LENGTH=397
Eucgr.A01210.1	2,39	22,87	13,48	11,77	1,30	AT2G37330.1 Symbols: ALS3 aluminum sensitive 3 chr2:15669190-15670172 FORWARD LENGTH=273
Eucgr.F02576.1	1,50	15,27	13,34	11,80	6,16	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.J02380.1	0,08	41,16	23,60	6,95	0,59	No hits found
Eucgr.J01346.1	5,03	23,40	19,88	12,57	9,84	AT3G28960.1 Symbols: Transmembrane amino acid transporter family protein
Eucgr.G03057.2	6,66	0,46	0,60	0,88	3,69	AT3G55490.1 Symbols: GINS complex protein chr3:20572171-20573415 REVERSE LENGTH=185
Eucgr.B01816.1	4,00	0,82	0,55	0,47	1,58	AT1G09450.1 Symbols: Protein kinase superfamily protein chr1:3049066-3052858 FORWARD LENGTH=599
Eucgr.J02488.1	0,26	4,09	3,62	2,37	1,00	AT4G35830.1 Symbols: ACO1 aconitase 1 chr4:16973007-16977949 REVERSE LENGTH=898
Eucgr.I02362.1	2,24	19,84	25,59	13,16	5,81	No hits found
Eucgr.I01253.1	2,11	0,00	0,00	0,00	0,41	AT1G29460.1 Symbols: SAUR-like auxin-responsive protein family chr1:10307609-10308055 REVERSE LENGTH=148
Eucgr.B03096.1	3,63	0,23	0,40	0,86	2,13	AT1G26220.1 Symbols: Acyl-CoA N-acyltransferases (NAT) superfamily protein
Eucgr.E03345.1	4,48	28,75	25,03	19,54	10,67	AT3G47160.1 Symbols: RING/U-box superfamily protein chr3:17360238-17361879 REVERSE LENGTH=245

Eucgr.B00548.1	63,74	3,94	17,82	25,52	37,72	AT5G61340.1 Symbols: unknown protein;
Eucgr.A02895.1	3,55	0,44	0,35	0,23	1,65	AT5G62410.1 Symbols: SMC2, TTN3, ATCAP-E1, ATSMC4 structural maintenance of chromosomes 2
Eucgr.D01982.1	45,48	5,08	7,26	9,83	28,05	No hits found
Eucgr.B04023.1	14,16	0,76	2,41	4,46	10,39	AT1G06920.1 Symbols: ATOFP4, OFP4 ovate family protein 4 chr1:2124854-2125801 REVERSE LENGTH=315
Eucgr.A02076.1	8,53	42,73	39,27	27,33	17,59	AT1G09230.1 Symbols: RNA-binding (RRM/RBD/RNP motifs) family protein chr1:2979637-2982563 REVERSE LENGTH=442
Eucgr.J01549.1	3,52	0,40	1,27	2,00	2,35	AT5G37790.1 Symbols: Protein kinase superfamily protein chr5:15008433-15011025 REVERSE LENGTH=552
Eucgr.F04163.1	6,22	0,15	0,32	0,57	1,82	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.E02407.1	38,10	0,67	4,63	7,79	25,72	AT4G18260.1 Symbols: Cytochrome b561/ferric reductase transmembrane protein family
Eucgr.J01393.1	50,50	3,41	9,74	12,85	28,42	AT5G15630.1 Symbols: COBL4, IRX6 COBRA-like extracellular glycosyl-phosphatidyl inositol-anchored protein family
Eucgr.A00785.1	22,93	1,66	4,06	6,20	12,75	AT5G58300.2 Symbols: Leucine-rich repeat protein kinase family protein
Eucgr.K02087.4	6,80	27,91	26,92	24,77	15,29	AT3G25410.1 Symbols: Sodium Bile acid symporter family chr3:9214565-9216331 REVERSE LENGTH=431
Eucgr.C02095.1	13,35	0,28	0,63	0,53	6,51	AT2G24490.2 Symbols: RPA2, ATRPA2, ROR1, ATRPA32A, RPA32A replicon protein A2
Eucgr.H00371.1	30,04	1,31	1,37	1,51	6,47	AT3G59400.1 Symbols: GUN4 enzyme binding;tetrapyrrole binding chr3:21948881-21949678 REVERSE LENGTH=265
Eucgr.J01789.1	133,07	11,99	12,71	14,40	54,74	AT1G58170.1 Symbols: Disease resistance-responsive (dirigent-like protein) family protein
Eucgr.G02893.1	14,23	63,60	65,88	61,38	27,70	AT2G39450.1 Symbols: MTP11, ATMTTP11 Cation efflux family protein chr2:16471744-16473735 REVERSE LENGTH=394
Eucgr.B03492.1	6,01	36,52	38,76	40,11	17,67	AT1G67340.1 Symbols: HCP-like superfamily protein with MYND-type zinc finger
Eucgr.D00694.1	3,64	0,04	0,14	0,09	1,26	AT1G11125.1 Symbols: unknown protein;
Eucgr.G00872.1	29,17	3,70	6,80	10,08	22,79	AT5G39420.1 Symbols: cdc2cAt CDC2C chr5:15772232-15774929 FORWARD LENGTH=644
Eucgr.J01624.1	14,50	0,60	0,58	0,57	3,57	AT1G66150.1 Symbols: TMK1 transmembrane kinase 1 chr1:24631503-24634415 FORWARD LENGTH=942
Eucgr.G01704.2	13,07	1,85	3,60	3,66	9,66	AT1G45130.1 Symbols: BGAL5 beta-galactosidase 5 chr1:17065447-17069110 FORWARD LENGTH=732
Eucgr.F00716.1	0,20	14,38	5,01	1,52	0,05	No hits found
Eucgr.E01665.1	2,29	0,00	0,25	0,71	1,11	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.F03030.1	15,78	1,87	2,56	3,48	8,19	AT2G25737.1 Symbols: Sulfite exporter TauE/SafE family protein chr2:10977174-10979677 FORWARD LENGTH=476
Eucgr.J00771.1	25,63	2,00	4,00	8,36	15,61	AT2G17380.1 Symbols: AP19 associated protein 19 chr2:7553122-7554887 FORWARD LENGTH=161
Eucgr.G02194.1	40,97	3,97	5,52	5,11	21,61	AT1G71020.1 Symbols: ARM repeat superfamily protein chr1:26790825-26793105 REVERSE LENGTH=628
Eucgr.B00537.2	12,12	52,77	53,21	46,71	7,08	AT5G61380.1 Symbols: TOC1, APRR1, PRR1, AtTOC1 CCT motif -containing response regulator protein
Eucgr.E02713.1	12,05	0,32	1,00	0,93	7,19	AT1G29050.1 Symbols: TBL38 TRICHOME BIREFRINGENCE-LIKE 38 chr1:10136376-10139082 REVERSE LENGTH=380
Eucgr.C01116.1	2,78	0,51	0,07	0,57	0,71	No hits found
Eucgr.A01406.1	4,55	52,32	50,28	61,46	9,65	AT5G24490.1 Symbols: 30S ribosomal protein, putative chr5:8365690-8367178 FORWARD LENGTH=308

Eucgr.K00387.1	9,89	66,76	54,47	39,19	19,93	AT5G24300.2 Symbols: SSI1 Glycogen/starch synthases, ADP-glucose type
Eucgr.E00328.1	29,66	4,61	4,12	5,48	15,95	AT4G17670.1 Symbols: Protein of unknown function (DUF581) chr4:9833948-9834663 REVERSE LENGTH=159
Eucgr.G03331.1	27,98	0,22	0,44	0,41	9,29	AT5G12470.1 Symbols: Protein of unknown function (DUF3411) chr5:4044950-4047290 REVERSE LENGTH=386
Eucgr.G01113.1	13,11	0,09	0,35	0,14	4,96	AT5G09640.1 Symbols: SNG2, SCPL19 serine carboxypeptidase-like 19 chr5:2988373-2990966 FORWARD LENGTH=465
Eucgr.D00356.1	3,25	0,25	0,48	0,48	1,53	AT1G29630.2 Symbols: 5'-3' exonuclease family protein chr1:10349587-10353538 FORWARD LENGTH=735
Eucgr.G01940.1	32,36	9,37	12,12	14,28	23,17	AT1G12850.1 Symbols: Phosphoglycerate mutase family protein chr1:4380022-4381577 REVERSE LENGTH=405
Eucgr.A01839.1	0,02	15,06	6,70	2,41	0,05	AT1G03790.1 Symbols: SOM Zinc finger C-x8-C-x5-C-x3-H type family protein
Eucgr.C01876.1	80,43	21,36	19,01	30,24	41,50	AT5G24930.1 Symbols: ATCOL4, COL4 CONSTANS-like 4 chr5:8589325-8590949 FORWARD LENGTH=406
Eucgr.E01075.1	22,00	1,02	0,93	1,08	6,25	AT5G59970.1 Symbols: Histone superfamily protein chr5:24146352-24146663 REVERSE LENGTH=103
Eucgr.J02275.1	20,40	1,65	2,51	3,90	11,31	AT3G27350.2 Symbols: unknown protein;
Eucgr.D01068.1	0,86	0,00	0,00	0,04	0,29	AT1G54730.2 Symbols: Major facilitator superfamily protein chr1:20424471-20429978 FORWARD LENGTH=470
Eucgr.E01150.1	5,09	1,09	0,72	1,50	2,41	AT4G19350.1 Symbols: EMB3006 embryo defective 3006 chr4:10562214-10563676 REVERSE LENGTH=180
Eucgr.A02408.1	1,36	0,05	0,05	0,07	0,66	AT1G51200.4 Symbols: A20/AN1-like zinc finger family protein chr1:18985690-18986211 FORWARD LENGTH=173
Eucgr.C00929.1	1,58	11,43	10,95	8,31	4,05	AT4G29420.1 Symbols: F-box/RNI-like superfamily protein chr4:14470715-14472204 REVERSE LENGTH=446
Eucgr.F01593.1	0,11	2,13	1,96	2,03	0,40	AT3G50940.1 Symbols: P-loop containing nucleoside triphosphate hydrolases superfamily protein
Eucgr.K03212.1	49,08	6,21	13,83	27,31	33,81	AT2G37080.1 Symbols: RIP3 ROP interactive partner 3 chr2:15581565-15584057 REVERSE LENGTH=583
Eucgr.F02437.1	1,95	0,16	0,06	0,08	0,64	AT1G78110.1 Symbols: unknown protein;
Eucgr.H02125.1	16,00	2,32	5,62	10,27	10,47	AT5G67210.1 Symbols: Protein of unknown function (DUF579) chr5:26819019-26819972 FORWARD LENGTH=317
Eucgr.E03246.1	586,63	21,03	39,85	53,99	216,49	No hits found
Eucgr.J00068.2	29,62	1,64	1,20	0,88	20,52	No hits found
Eucgr.K02905.1	2,66	10,24	11,71	9,55	5,16	AT3G28580.1 Symbols: P-loop containing nucleoside triphosphate hydrolases superfamily protein
Eucgr.E00030.1	15,04	73,32	53,16	49,43	31,23	AT3G61860.1 Symbols: ATRSP31, RSP31, At-RS31, RS31 RNA-binding (RRM/RBD/RNP motifs) family protein
Eucgr.H02448.1	4,40	0,03	0,08	0,12	1,43	AT1G67750.1 Symbols: Pectate lyase family protein chr1:25401660-25403165 FORWARD LENGTH=408
Eucgr.G00300.1	7,14	0,39	1,63	3,01	4,16	AT2G21100.1 Symbols: Disease resistance-responsive (dirigent-like protein) family protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.F01447.1	26,34	272,06	261,70	274,46	81,80	AT1G79040.1 Symbols: PSBR photosystem II subunit R chr1:29736085-29736781 FORWARD LENGTH=140
Eucgr.A02608.1	1,74	0,21	0,14	0,14	0,82	AT2G36026.1 Symbols: Ovate family protein chr2:15127265-15127816 FORWARD LENGTH=183
Eucgr.B03728.1	4,92	0,39	1,24	1,26	3,07	AT5G12340.1 Symbols: unknown protein;
Eucgr.D01823.1	0,27	27,86	24,42	19,89	1,52	AT4G25700.1 Symbols: BETA-OHASE 1, B1, chy1, BCH1 beta-hydroxylase 1
Eucgr.A02792.1	37,93	177,95	127,38	112,79	61,23	AT4G25570.1 Symbols: ACYB-2 Cytochrome b561/ferric reductase transmembrane protein family

Eucgr.K00285.1	0,37	0,02	0,02	0,04	0,16	AT5G23940.1 Symbols: EMB3009, PEL3, DCR HXXXD-type acyl-transferase family protein
Eucgr.F02906.1	10,02	80,27	56,19	35,86	22,01	AT5G17300.1 Symbols: RVE1 Homeodomain-like superfamily protein chr5:5690435-5692435 REVERSE LENGTH=387
Eucgr.H01043.1	9,20	0,13	0,47	0,60	4,58	AT4G02060.2 Symbols: PRL Minichromosome maintenance (MCM2/3/5) family protein
Eucgr.K02355.1	5,36	0,03	0,04	0,04	2,50	AT2G43840.2 Symbols: UGT74F1 UDP-glycosyltransferase 74 F1 chr2:18157681-18159166 FORWARD LENGTH=449
Eucgr.B02731.1	55,61	3,86	9,99	14,59	35,60	AT5G02010.1 Symbols: ATROPGEF7, ROPGEF7 RHO guanyl-nucleotide exchange factor 7
Eucgr.J01060.1	2,42	0,15	0,05	0,07	0,41	AT3G07510.2 Symbols: unknown protein
Eucgr.G03227.1	79,40	0,86	1,85	1,91	31,98	AT5G07030.1 Symbols: Eukaryotic aspartyl protease family protein chr5:2183600-2185717 REVERSE LENGTH=455
Eucgr.E02770.1	8,60	0,25	0,26	0,27	5,12	No hits found
Eucgr.C00451.1	13,11	75,21	62,76	49,39	25,17	AT2G12400.1 Symbols: unknown protein;
Eucgr.B02321.1	0,02	3,51	1,36	0,32	0,09	AT2G29090.2 Symbols: CYP707A2 cytochrome P450, family 707, subfamily A, polypeptide 2
Eucgr.H03469.1	59,65	3,72	4,50	5,26	30,06	AT4G03500.1 Symbols: Ankyrin repeat family protein chr4:1553453-1556571 FORWARD LENGTH=652
Eucgr.G00575.1	4,29	0,94	0,59	0,85	2,55	AT1G61000.1 Symbols: unknown protein
Eucgr.A01897.1	0,11	82,14	33,20	9,83	0,61	No hits found
Eucgr.K00573.1	6,39	0,60	1,09	1,72	3,91	AT3G62150.1 Symbols: PGP21 P-glycoprotein 21 chr3:23008755-23013579 REVERSE LENGTH=1296
Eucgr.G02934.1	43,45	2,01	5,66	12,10	29,12	AT3G55420.1 Symbols: unknown protein;
Eucgr.L00248.1	0,17	162,07	86,93	32,28	0,69	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.B02197.1	7,47	0,13	0,37	0,41	3,11	AT1G09540.1 Symbols: MYB61, ATMYB61 myb domain protein 61 chr1:3086333-3087689 FORWARD LENGTH=366
Eucgr.F00501.1	2,52	0,06	0,05	0,19	1,40	AT1G33110.1 Symbols: MATE efflux family protein chr1:12005084-12008618 FORWARD LENGTH=494
Eucgr.K01667.2	64,37	4,52	11,87	23,49	35,51	AT3G56480.1 Symbols: myosin heavy chain-related chr3:20936480-20938986 REVERSE LENGTH=490
Eucgr.B03348.1	26,89	0,21	0,15	0,07	9,47	AT1G10550.1 Symbols: XTH33, XET xyloglucan:xyloglucosyl transferase 33 chr1:3479257-3480957 REVERSE LENGTH=310
Eucgr.D00321.1	127,97	0,75	1,93	4,91	23,84	AT2G34430.1 Symbols: LHB1B1, LHCB1.4 light-harvesting chlorophyll-protein complex II subunit B1
Eucgr.G03135.1	10,32	1,20	2,05	2,76	6,88	AT2G40620.1 Symbols: Basic-leucine zipper (bZIP) transcription factor family protein
Eucgr.F03950.1	38,76	1,85	3,23	5,43	26,47	AT1G80690.1 Symbols: PPPDE putative thiol peptidase family protein chr1:30329283-30330524 REVERSE LENGTH=227
Eucgr.L00055.1	2,70	13,38	11,22	10,11	6,18	AT5G16990.1 Symbols: Zinc-binding dehydrogenase family protein chr5:5581831-5583849 REVERSE LENGTH=343
Eucgr.E04043.2	12,23	1,97	3,14	4,28	8,85	AT1G11790.1 Symbols: ADT1 arogenate dehydratase 1 chr1:3981476-3984962 FORWARD LENGTH=392
Eucgr.J02569.1	0,21	2,64	2,68	2,34	0,63	AT1G20925.1 Symbols: Auxin efflux carrier family protein chr1:7289973-7292507 FORWARD LENGTH=472
GENE ID	FPKM	Annotation				
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.A02086.1	23,71	7,91	6,93	8,18	17,77	AT4G32890.1 Symbols: GATA9 GATA transcription factor 9 chr4:15875598-15876615 FORWARD LENGTH=308
Eucgr.B02796.1	12,04	0,28	1,95	3,31	5,69	AT2G38080.1 Symbols: IRX12, LAC4, ATLMCO4, LMCO4 Laccase/Diphenol oxidase family protein
Eucgr.J00113.1	3,97	0,40	0,49	0,48	1,68	AT5G05240.1 Symbols: Uncharacterised conserved protein (UCP030365) chr5:1554023-1556230 REVERSE LENGTH=530

Eucgr.C04217.1	39,42	3,74	7,46	12,21	22,98	AT5G15050.1 Symbols: Core-2/l-branching beta-1,6-N-acetylglucosaminyltransferase family protein
Eucgr.D01955.1	0,05	0,17	0,84	0,76	0,22	AT2G45570.1 Symbols: CYP76C2 cytochrome P450, family 76, subfamily C, polypeptide 2
Eucgr.K02638.1	3,52	0,78	0,21	0,50	1,74	AT5G35520.1 Symbols: MIS12, ATMIS12 minichromosome instability 12 (mis12)-like
Eucgr.D00088.1	123,26	5,66	10,29	15,33	55,67	AT1G30760.1 Symbols: FAD-binding Berberine family protein chr1:10918321-10920441 FORWARD LENGTH=534
Eucgr.H02338.1	131,39	30,24	33,82	38,59	75,02	AT3G49720.2 Symbols: unknown protein;
Eucgr.C03090.1	0,21	0,02	0,00	0,01	0,07	AT1G79670.1 Symbols: RFO1, WAKL22 Wall-associated kinase family protein
Eucgr.D00322.1	364,73	15,53	18,75	41,58	83,31	AT2G34430.1 Symbols: LHB1B1, LHCB1.4 light-harvesting chlorophyll-protein complex II subunit B1
Eucgr.H04549.1	2,07	0,03	0,19	0,46	0,82	AT4G15920.1 Symbols: SWEET17, AtSWEET17 Nodulin MtN3 family protein chr4:9030742-9033343 REVERSE LENGTH=241
Eucgr.L03123.1	2,67	17,17	14,83	13,58	5,36	AT4G29420.1 Symbols: F-box/RNI-like superfamily protein chr4:14470715-14472204 REVERSE LENGTH=446
Eucgr.A02140.1	4,04	14,31	15,72	13,30	7,22	AT3G03710.1 Symbols: RIF10, PNP polyribonucleotide nucleotidyltransferase, putative
Eucgr.F01611.1	418,71	48,29	107,92	147,05	182,78	AT1G75500.2 Symbols: WAT1 Walls Are Thin 1 chr1:28338282-28340091 REVERSE LENGTH=389
Eucgr.I01231.1	2,43	160,13	93,47	46,77	8,13	AT2G15970.2 Symbols: COR413-PM1 cold regulated 413 plasma membrane 1
Eucgr.D01441.2	12,88	36,00	35,04	32,63	21,09	AT4G16440.1 Symbols: ferredoxin hydrogenases chr4:9269263-9271510 REVERSE LENGTH=474
Eucgr.K02422.1	3,70	20,83	17,92	15,51	7,45	AT5G03560.1 Symbols: Tetratricopeptide repeat (TPR)-like superfamily protein
Eucgr.J02173.1	77,27	4,25	5,33	4,84	37,11	AT5G40150.1 Symbols: Peroxidase superfamily protein chr5:16059750-16060736 REVERSE LENGTH=328
Eucgr.B02382.1	0,24	6,79	4,92	3,65	0,93	AT1G07645.1 Symbols: ATDSI-1VOC, DSI-1VOC dessication-induced 1VOC superfamily protein
Eucgr.H00791.1	7,86	0,92	1,30	2,29	5,65	AT5G48720.2 Symbols: XRI, XRI1 x-ray induced transcript 1 chr5:19759233-19761621 FORWARD LENGTH=300
Eucgr.J02841.1	3,06	0,67	0,34	0,92	1,71	AT1G17520.1 Symbols: Homeodomain-like/winged-helix DNA-binding family protein chr1:6024959-6027224 REVERSE LENGTH=296
Eucgr.D01777.1	3,63	0,14	0,17	0,24	1,36	AT4G23740.1 Symbols: Leucine-rich repeat protein kinase family protein chr4:12367063-12369159 FORWARD LENGTH=638
Eucgr.H04936.1	8,51	0,04	0,03	0,08	2,50	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.J01431.1	11,18	0,12	0,16	0,31	3,93	AT3G30530.1 Symbols: ATBZIP42, bZIP42 basic leucine-zipper 42 chr3:12139512-12140033 FORWARD LENGTH=173
Eucgr.K00151.4	15,32	2,94	3,48	4,89	11,48	AT4G25290.1 Symbols: DNA photolyases;DNA photolyases chr4:12941486-12944994 REVERSE LENGTH=692
Eucgr.G03233.1	7,44	0,38	0,69	0,51	3,70	No hits found
Eucgr.B01963.1	0,36	0,00	0,00	0,06	0,13	AT3G14620.1 Symbols: CYP72A8 cytochrome P450, family 72, subfamily A, polypeptide 8
Eucgr.H01495.1	3,96	0,04	0,10	0,23	1,46	AT3G53810.1 Symbols: Concanavalin A-like lectin protein kinase family protein
Eucgr.F04162.1	1,95	0,09	0,04	0,06	0,72	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.C01045.1	11,70	0,52	1,49	2,57	8,06	AT4G32350.1 Symbols: Regulator of Vps4 activity in the MVB pathway protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.A02929.1	45,35	195,33	180,49	171,77	70,43	AT4G25130.1 Symbols: PMSR4 peptide met sulfoxide reductase 4 chr4:12898802-12899998 REVERSE LENGTH=258
Eucgr.A02121.2	41,44	3,24	6,66	12,13	26,68	AT5G32450.1 Symbols: RNA binding (RRM/RBD/RNP motifs) family protein

Eucgr.E00070.1	32,68	1,90	4,13	6,97	20,34	AT1G01430.1 Symbols: TBL25 TRICHOME BIREFRINGENCE-LIKE 25 chr1:156953-158536 REVERSE LENGTH=456
Eucgr.K00075.1	14,32	44,40	42,17	39,60	22,57	AT5G16750.1 Symbols: TOZ Transducin family protein / WD-40 repeat family protein
Eucgr.K01353.1	1,93	12,72	12,64	11,28	4,29	AT5G60870.1 Symbols: Regulator of chromosome condensation (RCC1) family protein
Eucgr.E01876.1	3,68	39,38	33,04	35,68	9,70	AT5G45040.1 Symbols: Cytochrome c chr5:18175566-18176851 REVERSE LENGTH=175
Eucgr.B00601.1	2,95	0,16	0,23	0,35	1,44	AT5G08020.1 Symbols: ATRPA70B, RPA70B RPA70-kDa subunit B chr5:2572107-2574879 FORWARD LENGTH=604
Eucgr.G03122.1	259,31	5,64	10,62	21,19	94,55	AT2G40480.1 Symbols: Plant protein of unknown function (DUF827) chr2:16910425-16912444 FORWARD LENGTH=518
Eucgr.J03127.1	8,27	1576,75	1347,55	881,92	925,42	AT1G16030.1 Symbols: Hsp70b heat shock protein 70B chr1:5502386-5504326 REVERSE LENGTH=646
Eucgr.K00315.1	0,76	0,10	0,05	0,07	0,29	AT5G52950.1 Symbols: unknown protein;
Eucgr.E00460.1	0,08	7,37	6,22	4,26	0,35	AT4G12430.1 Symbols: TPPF Haloacid dehalogenase-like hydrolase (HAD) superfamily protein
Eucgr.J02714.1	0,74	0,13	0,00	0,09	0,28	AT2G19190.1 Symbols: FRK1 FLG22-induced receptor-like kinase 1 chr2:8326067-8329893 REVERSE LENGTH=876
Eucgr.G03078.1	35,58	1,80	4,90	11,75	22,17	AT3G11090.1 Symbols: LBD21 LOB domain-containing protein 21 chr3:3475036-3475533 FORWARD LENGTH=165
Eucgr.A01720.1	8,57	0,69	0,78	1,00	3,94	AT1G01380.1 Symbols: ETC1 Homeodomain-like superfamily protein chr1:147267-147740 FORWARD LENGTH=83
Eucgr.F03729.1	0,77	0,03	0,07	0,11	0,40	AT4G08500.1 Symbols: MEK1, ATMEK1, MAPKKK8, ARAKIN MAPK/ERK kinase kinase 1
Eucgr.E03922.1	7,89	119,99	97,52	76,35	18,49	No hits found
Eucgr.H01054.1	4,39	0,00	0,04	0,39	1,57	AT4G02090.1 Symbols: unknown protein;
Eucgr.I00191.1	18,91	304,30	179,39	171,43	32,65	AT5G64530.1 Symbols: ANAC104, XND1 xylem NAC domain 1 chr5:25795360-25796699 FORWARD LENGTH=187
Eucgr.D01367.1	2,05	0,00	0,00	0,00	0,62	AT5G01840.1 Symbols: ATOFP1, OFP1 ovate family protein 1 chr5:324552-325364 FORWARD LENGTH=270
Eucgr.D01770.1	9,52	146,22	130,43	116,67	25,55	AT4G24015.1 Symbols: RING/U-box superfamily protein chr4:12469887-12471197 REVERSE LENGTH=174
Eucgr.K02218.1	33,80	4,07	3,06	6,15	18,24	No hits found
Eucgr.B03813.1	16,93	52,12	51,45	58,11	24,66	AT5G22100.1 Symbols: RNA cyclase family protein chr5:7329015-7330718 FORWARD LENGTH=375
Eucgr.J01834.1	9,39	3,15	3,11	3,89	6,70	AT3G02820.1 Symbols: zinc knuckle (CCHC-type) family protein chr3:611573-613294 FORWARD LENGTH=282
Eucgr.G01707.1	11,60	0,20	0,68	0,95	6,59	AT2G16440.1 Symbols: MCM4 Minichromosome maintenance (MCM2/3/5) family protein
Eucgr.H03450.2	16,30	1,23	2,24	3,51	10,18	AT3G22550.1 Symbols: Protein of unknown function (DUF581) chr3:7991827-7992805 REVERSE LENGTH=267
Eucgr.K01388.2	14,26	52,88	45,91	42,56	23,17	AT5G48240.3 Symbols: unknown protein;
Eucgr.K00903.1	0,38	3,91	4,61	3,45	1,26	AT1G03940.1 Symbols: HXXXD-type acyl-transferase family protein chr1:1009542-1010951 REVERSE LENGTH=469
Eucgr.G01683.1	15,59	1,40	2,46	3,13	10,99	AT4G34530.1 Symbols: ClB1 cryptochrome-interacting basic-helix-loop-helix 1
Eucgr.I00196.1	14,36	0,40	0,47	0,67	5,42	AT1G49780.1 Symbols: PUB26 plant U-box 26 chr1:18429024-18430289 REVERSE LENGTH=421
Eucgr.B02755.1	22,23	1,12	1,22	3,87	8,75	AT3G54050.2 Symbols: HCEF1 high cyclic electron flow 1 chr3:20016951-20018527 FORWARD LENGTH=417
Eucgr.D00371.1	0,17	0,00	0,00	0,03	0,07	AT4G19090.1 Symbols: Protein of unknown function (DUF594) chr4:10449900-10452757 FORWARD LENGTH=751
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR

Eucgr.B00072.1	4,34	0,24	0,30	0,28	2,65	AT2G31900.1 Symbols: XIF, ATXIF, ATMYO5 myosin-like protein XIF chr2:13560760-13569623 REVERSE LENGTH=1556
Eucgr.G01052.1	19,72	2,20	3,15	4,89	13,64	AT4G17980.1 Symbols: anac071, NAC071 NAC domain containing protein 71
Eucgr.H01669.1	5,20	0,09	0,39	0,93	2,37	AT5G67050.1 Symbols: alpha/beta-Hydrolases superfamily protein chr5:26759482-26761165 REVERSE LENGTH=477
Eucgr.G01558.1	0,26	0,00	0,00	0,06	0,10	AT5G10770.1 Symbols: Eukaryotic aspartyl protease family protein chr5:3403331-3405331 REVERSE LENGTH=474
Eucgr.D01019.1	58,87	10,87	11,41	15,76	36,62	AT3G23290.2 Symbols: LSH4 Protein of unknown function (DUF640) chr3:8326958-8328010 FORWARD LENGTH=195
Eucgr.B00238.2	5,78	0,61	0,51	1,16	2,96	AT4G24670.2 Symbols: TAR2 tryptophan aminotransferase related 2 chr4:12727940-12730694 REVERSE LENGTH=440
Eucgr.H05117.4	3,47	0,98	0,88	1,11	2,44	AT5G63860.1 Symbols: UVR8 Regulator of chromosome condensation (RCC1) family protein
Eucgr.F01643.1	3,53	0,00	0,17	0,36	2,30	AT1G75710.1 Symbols: C2H2-like zinc finger protein chr1:28428806-28431128 FORWARD LENGTH=462
Eucgr.E04300.1	14,03	37,96	39,12	36,51	23,84	AT2G35060.1 Symbols: KUP11 K+ uptake permease 11 chr2:14775184-14778184 REVERSE LENGTH=792
Eucgr.G02143.1	0,03	4,40	1,90	1,18	0,04	AT1G14870.1 Symbols: PCR2 PLANT CADMIUM RESISTANCE 2 chr1:5128591-5129458 REVERSE LENGTH=152
Eucgr.H02450.1	18,67	76,82	72,34	64,69	35,91	AT1G13160.1 Symbols: ARM repeat superfamily protein chr1:4484634-4488388 FORWARD LENGTH=804
Eucgr.F03668.1	39,12	1,79	5,22	7,67	22,63	AT3G44730.1 Symbols: ATKP1, KP1 kinesin-like protein 1 chr3:16285888-16290852 FORWARD LENGTH=1087
Eucgr.D02639.1	0,40	37,14	14,82	4,44	1,20	AT5G62150.1 Symbols: peptidoglycan-binding LysM domain-containing protein
Eucgr.J01777.2	22,81	2,90	5,14	7,05	16,94	AT2G02170.2 Symbols: Remorin family protein chr2:556595-558610 REVERSE LENGTH=486
Eucgr.A01623.1	10,29	25,55	28,96	25,95	16,84	AT4G39160.2 Symbols: Homeodomain-like superfamily protein chr4:18236828-18240433 FORWARD LENGTH=624
Eucgr.F04485.1	0,10	11,70	7,83	4,51	0,42	AT2G21820.1 Symbols: unknown protein;
Eucgr.K00648.1	14,25	0,45	0,21	0,76	4,92	AT5G35740.1 Symbols: Carbohydrate-binding X8 domain superfamily protein
Eucgr.K02300.1	29,89	4,72	6,31	10,05	21,14	AT3G48750.1 Symbols: CDKA;1, CDC2AAT, CDK2, CDC2, CDC2A, CDKA1 cell division control 2
Eucgr.C01575.1	3,50	0,09	0,11	0,09	1,44	AT5G25580.1 Symbols: unknown protein
Eucgr.A00208.1	0,02	3,66	1,60	0,24	0,06	AT3G62560.1 Symbols: Ras-related small GTP-binding family protein chr3:23137539-23138880 FORWARD LENGTH=193
Eucgr.D00412.1	0,02	1,92	1,24	0,20	0,05	AT5G45690.1 Symbols: Protein of unknown function (DUF1264) chr5:18535139-18536259 REVERSE LENGTH=247
Eucgr.K01851.1	0,01	0,83	0,66	0,28	0,03	AT2G41970.1 Symbols: Protein kinase superfamily protein chr2:17520517-17522304 REVERSE LENGTH=365
Eucgr.G03284.1	0,01	0,88	0,67	0,79	0,04	AT5G13790.1 Symbols: AGL15 AGAMOUS-like 15 chr5:4449128-4450802 REVERSE LENGTH=268
Eucgr.F00591.1	0,02	2,18	1,55	0,20	0,05	AT1G75750.1 Symbols: GASA1 GAST1 protein homolog 1 chr1:28441813-28442284 REVERSE LENGTH=98
Eucgr.A00995.1	7,47	98,88	59,96	43,40	13,77	AT2G46940.1 Symbols: unknown protein;
Eucgr.D00583.1	16,76	0,59	0,95	1,93	8,70	No hits found
Eucgr.H02351.1	21,03	3,01	6,16	6,08	14,74	AT5G65710.1 Symbols: HSL2 HAESA-like 2 chr5:26292372-26295440 FORWARD LENGTH=993
Eucgr.K02329.1	4,76	23,72	33,52	24,98	12,17	AT1G73380.1 Symbols: unknown protein;
Eucgr.C00156.1	19,40	0,91	1,38	2,83	12,05	AT4G30230.1 Symbols: unknown protein;
Eucgr.K02060.1	4,73	21,67	20,39	19,59	8,75	AT1G19490.1 Symbols: Basic-leucine zipper (bZIP) transcription factor family protein
Eucgr.B02239.1	1,35	372,30	180,88	56,90	3,52	AT3G47340.1 Symbols: ASN1, DIN6, AT-ASN1 glutamine-dependent asparagine synthase 1

GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	
Eucgr.F02999.1	13,23	0,37	1,71	2,57	7,82	AT1G22610.1 Symbols: C2 calcium/lipid-binding plant phosphoribosyltransferase family protein
Eucgr.E00446.1	10,53	0,08	0,53	0,74	4,93	AT1G62500.1 Symbols: Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein
Eucgr.D00059.1	7,86	22,20	23,11	21,70	12,79	AT5G18820.1 Symbols: EMB3007 TCP-1/cpn60 chaperonin family protein chr5:6271549-6274153 FORWARD LENGTH=575
Eucgr.F03226.1	3,54	0,51	0,59	1,04	2,37	AT3G14740.2 Symbols: RING/FYVE/PHD zinc finger superfamily protein chr3:4952185-4953306 REVERSE LENGTH=343
Eucgr.D02618.1	18,64	41,27	44,62	39,80	29,03	AT4G31120.1 Symbols: SKB1, ATPRMT5 SHK1 binding protein 1 chr4:15132185-15136568 REVERSE LENGTH=642
Eucgr.A01416.1	0,08	12,40	7,60	4,34	0,29	AT4G27670.1 Symbols: HSP21 heat shock protein 21 chr4:13819048-13819895 REVERSE LENGTH=227
Eucgr.E02733.1	0,90	0,00	0,03	0,02	0,40	AT3G49340.1 Symbols: Cysteine proteinases superfamily protein chr3:18293347-18294577 REVERSE LENGTH=341
Eucgr.B01665.1	7,24	0,70	1,44	1,58	4,68	AT3G62630.1 Symbols: Protein of unknown function (DUF1645) chr3:23163937-23165079 REVERSE LENGTH=380
Eucgr.F01247.1	62,51	5,65	5,19	8,11	34,14	AT1G78780.2 Symbols: pathogenesis-related family protein chr1:29621447-29622431 REVERSE LENGTH=238
Eucgr.J01592.1	3,62	0,18	0,17	0,45	1,49	AT1G70780.1 Symbols: unknown protein;
Eucgr.F02449.1	0,21	0,00	0,00	0,05	0,09	AT1G02390.1 Symbols: ATPAT2, GPAT2 glycerol-3-phosphate acyltransferase 2
Eucgr.I01944.1	0,07	1,86	2,17	1,75	0,27	AT5G39785.1 Symbols: Protein of unknown function (DUF1666) chr5:15929257-15932215 FORWARD LENGTH=606
Eucgr.D01116.1	2,47	0,11	0,03	0,18	0,78	AT3G17350.1 Symbols: unknown protein
Eucgr.H02856.1	0,36	0,00	0,01	0,00	0,15	AT1G61820.1 Symbols: BGLU46 beta glucosidase 46 chr1:22835452-22838444 FORWARD LENGTH=516
Eucgr.C04122.1	13,12	1,84	1,88	3,31	7,45	AT3G24460.1 Symbols: Serinc-domain containing serine and sphingolipid biosynthesis protein
Eucgr.I01141.1	1,91	0,04	0,00	0,00	0,58	AT5G49730.1 Symbols: ATFRO6, FRO6 ferric reduction oxidase 6 chr5:20201355-20204455 REVERSE LENGTH=738
Eucgr.L00250.1	0,08	24,46	8,06	2,35	0,03	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.I02395.1	36,67	7955,76	4612,95	3696,64	424,49	No hits found
Eucgr.I01970.1	1,48	0,02	0,14	0,12	0,81	AT3G29060.1 Symbols: EXS (ERD1/XPR1/SYG1) family protein chr3:11044990-11048465 REVERSE LENGTH=800
Eucgr.K02663.1	4,01	55,92	37,45	23,69	8,00	AT1G73920.1 Symbols: alpha/beta-Hydrolases superfamily protein chr1:27791194-27794698 FORWARD LENGTH=704
Eucgr.F03947.1	12,82	0,29	0,60	0,85	5,97	AT5G25190.1 Symbols: Integrase-type DNA-binding superfamily protein chr5:8707007-8707655 REVERSE LENGTH=181
Eucgr.F02971.1	9,35	27,60	31,49	27,79	16,25	AT1G71490.1 Symbols: Tetraatricopeptide repeat (TPR)-like superfamily protein
Eucgr.G02591.1	14,31	50,74	52,41	62,14	23,31	AT5G08520.1 Symbols: Duplicated homeodomain-like superfamily protein chr5:2755470-2757741 REVERSE LENGTH=298
Eucgr.G02992.1	2,45	0,00	0,08	0,11	1,25	AT2G37820.1 Symbols: Cysteine/Histidine-rich C1 domain family protein
Eucgr.J02002.1	13,21	0,81	0,99	1,03	6,89	AT1G07420.1 Symbols: ATSMO2, SMO2-1 sterol 4-alpha-methyl-oxidase 2-1
Eucgr.D02220.1	10,52	0,86	1,06	1,60	5,86	AT1G32250.1 Symbols: Calcium-binding EF-hand family protein chr1:11639843-11640343 FORWARD LENGTH=166
Eucgr.H03598.1	26,97	1,37	3,01	6,08	17,56	AT3G06145.1 Symbols: unknown protein;
Eucgr.K00264.1	1191,00	71,53	132,92	190,87	475,74	AT5G23860.2 Symbols: TUB8 tubulin beta 8 chr5:8042962-8044528 FORWARD LENGTH=449
Eucgr.J02238.1	2,25	0,29	0,13	0,21	1,36	AT3G27640.1 Symbols: Transducin/WD40 repeat-like superfamily protein

Eucgr.G03119.1	52,32	0,71	0,44	0,43	12,97	AT2G40460.1 Symbols: Major facilitator superfamily protein chr2:16897123-16901171 FORWARD LENGTH=583
Eucgr.B00615.2	11,42	0,62	0,78	1,56	4,90	AT1G26355.1 Symbols: SP1L1 SPIRAL1-like1 chr1:9117950-9118611 REVERSE LENGTH=113
GENE ID	FPKM	Annotation				
CT	CH	NAR	CHSTOP	NARSTOP	TAIR	
Eucgr.H00748.1	40,86	3,51	3,54	3,08	24,78	AT3G07340.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.L00235.1	0,04	73,19	38,88	15,60	0,03	AT1G56600.1 Symbols: AtGolS2, GoIS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.F01599.1	18,17	0,64	0,62	0,66	8,13	AT1G75390.1 Symbols: AtbZIP44, bZIP44 basic leucine-zipper 44 chr1:28292224-28292745 FORWARD LENGTH=173
Eucgr.F00741.2	49,64	158,03	144,26	149,25	88,78	AT1G11910.1 Symbols: APA1, ATAPA1 aspartic proteinase A1 chr1:4017119-4019874 REVERSE LENGTH=506
Eucgr.B02318.1	3,10	0,14	0,04	0,00	1,34	AT2G29125.1 Symbols: RTFL2, DVL13 ROTUNDIFOLIA like 2 chr2:12523472-12523819 FORWARD LENGTH=115
Eucgr.I01316.1	12,00	2,71	2,61	4,89	7,64	AT4G34660.1 Symbols: SH3 domain-containing protein chr4:16545595-16548294 REVERSE LENGTH=368
Eucgr.A01602.1	45,92	4,95	10,83	17,13	31,40	AT1G63300.1 Symbols: Myosin heavy chain-related protein chr1:23482193-23486067 FORWARD LENGTH=1029
Eucgr.K01496.1	0,04	0,59	1,21	2,78	0,02	AT1G55370.2 Symbols: NDF5 NDH-dependent cyclic electron flow 5 chr1:20674852-20676071 FORWARD LENGTH=354
Eucgr.K01352.1	338,38	22,20	30,35	15,76	184,16	AT3G22120.1 Symbols: CWLP cell wall-plasma membrane linker protein chr3:7795283-7796287 REVERSE LENGTH=334
Eucgr.E02410.1	0,00	32,77	24,51	6,33	0,10	AT1G07900.1 Symbols: LBD1 LOB domain-containing protein 1 chr1:2442624-2443631 FORWARD LENGTH=190
Eucgr.L03254.1	0,00	2,32	1,51	0,40	0,04	AT1G73190.1 Symbols: ALPHA-TIP, TIP3;1 Aquaporin-like superfamily protein
Eucgr.B03510.1	9,53	159,19	65,25	39,11	19,24	AT3G48660.1 Symbols: Protein of unknown function (DUF 3339) chr3:18029659-18030133 FORWARD LENGTH=89
Eucgr.I01142.1	4,35	0,03	0,08	0,03	1,54	AT5G49740.1 Symbols: ATFRO7, FRO7 ferric reduction oxidase 7 chr5:20205549-20208628 REVERSE LENGTH=747
Eucgr.E03147.1	2,67	26,90	16,87	14,49	6,25	AT3G47160.2 Symbols: RING/U-box superfamily protein chr3:17360238-17361879 REVERSE LENGTH=257
Eucgr.D01150.1	2,55	0,06	0,09	0,39	1,38	AT2G44580.1 Symbols: zinc ion binding chr2:18402081-18403412 REVERSE LENGTH=386
Eucgr.F03300.1	13,76	1,85	1,93	2,95	7,82	AT5G53900.2 Symbols: Serine/threonine-protein kinase WNK (With No Lysine)-related
Eucgr.G02154.1	0,09	3,17	1,68	1,45	0,05	AT1G14870.1 Symbols: PCR2 PLANT CADMIUM RESISTANCE 2 chr1:5128591-5129458 REVERSE LENGTH=152
Eucgr.B03746.1	45,97	0,29	0,54	0,50	12,30	AT5G07990.1 Symbols: TT7, CYP75B1, D501 Cytochrome P450 superfamily protein
Eucgr.E04058.1	36,99	2,23	1,91	2,85	62,13	No hits found
Eucgr.B02749.1	5,57	1,08	1,14	1,82	4,23	AT4G03120.1 Symbols: C2H2 and C2HC zinc fingers superfamily protein chr4:1385747-1387331 FORWARD LENGTH=207
Eucgr.E01278.1	25,24	2,15	3,90	6,43	18,03	AT3G07040.1 Symbols: RPM1, RPS3 NB-ARC domain-containing disease resistance protein
Eucgr.B00857.1	5,71	19,53	23,20	19,36	10,32	AT3G18100.1 Symbols: MYB4R1, AtMYB4R1 myb domain protein 4r1 chr3:6200689-6204583 FORWARD LENGTH=847
Eucgr.A00596.1	1,52	9,49	13,79	14,11	3,29	AT2G38290.1 Symbols: ATAMT2, AMT2;1, AMT2 ammonium transporter 2
Eucgr.B00330.1	24,60	0,65	2,92	6,17	15,66	AT1G12220.2 Symbols: RP55 Disease resistance protein (CC-NBS-LRR class) family
Eucgr.G03123.1	1,81	0,08	0,04	0,05	0,81	AT5G05330.1 Symbols: HMG-box (high mobility group) DNA-binding family protein
Eucgr.I02361.1	5,43	55,55	51,56	32,51	12,50	No hits found
Eucgr.E03059.1	3,55	0,42	0,23	0,37	1,85	No hits found

Eucgr.B02000.1	31,38	0,43	0,84	2,08	13,48	AT1G72210.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.F03538.2	110,08	2,56	3,99	4,35	61,28	AT1G76160.1 Symbols: sks5 SKU5 similar 5 chr1:28578211-28581020 REVERSE LENGTH=541
Eucgr.B00523.1	77,15	4,49	6,63	14,78	44,66	AT3G47470.1 Symbols: LHCA4, CAB4 light-harvesting chlorophyll-protein complex l subunit A4
Eucgr.J01663.1	97,41	1,21	1,87	3,31	75,71	AT5G13870.1 Symbols: EXGT-A4, XTH5 xyloglucan endotransglucosylase/hydrolase 5
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.F03555.1	13,95	1,61	1,58	2,17	7,86	No hits found
Eucgr.A02867.1	21,50	77,97	60,27	57,79	33,36	AT5G62200.1 Symbols: Embryo-specific protein 3, (ATS3) chr5:24984463-24985674 REVERSE LENGTH=190
Eucgr.H02220.1	18,05	0,74	0,78	1,29	10,03	AT1G47128.1 Symbols: RD21, RD21A Granulin repeat cysteine protease family protein
Eucgr.D00363.1	0,44	0,09	0,04	0,09	0,24	AT5G45470.1 Symbols: Protein of unknown function (DUF594) chr5:18422164-18424764 REVERSE LENGTH=866
Eucgr.I01755.1	2,80	56,99	38,16	17,61	1,21	AT5G40390.1 Symbols: SIP1 Raffinose synthase family protein chr5:16161720-16165085 FORWARD LENGTH=783
Eucgr.H03017.2	43,78	3,51	4,59	5,87	21,12	AT3G23410.1 Symbols: ATFAO3, FAO3 fatty alcohol oxidase 3 chr3:8382860-8386024 FORWARD LENGTH=746
Eucgr.H02628.2	25,01	3,03	4,72	4,59	16,98	AT5G48560.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.G02193.1	4,85	0,09	0,39	0,56	2,25	AT2G01340.1 Symbols: At17.1 Encodes a protein whose expression is responsive to nematode infection.
Eucgr.G02743.1	1,23	16,29	14,12	10,00	2,99	AT5G03530.1 Symbols: ATRAB ALPHA, ATRAB, ATRAB18B, ATRABC2A, RABC2A RAB GTPase homolog C2A
Eucgr.J00363.1	88,68	3,76	16,43	29,57	53,35	AT5G41040.1 Symbols: HXXXD-type acyl-transferase family protein chr5:16431448-16433086 FORWARD LENGTH=457
Eucgr.C00061.1	7,18	0,77	0,94	0,91	3,86	AT2G07170.1 Symbols: ARM repeat superfamily protein chr2:2975565-2978692 FORWARD LENGTH=820
Eucgr.F01455.1	36,69	6,23	9,26	12,83	29,36	AT5G45030.2 Symbols: Trypsin family protein chr5:18172981-18175238 FORWARD LENGTH=607
Eucgr.L03276.1	1,64	0,10	0,00	0,40	0,66	No hits found
Eucgr.C03867.1	6,81	0,81	1,11	1,30	3,93	AT3G24460.1 Symbols: Serinc-domain containing serine and sphingolipid biosynthesis protein
Eucgr.I02197.1	15,27	1,38	2,05	1,80	10,08	AT1G20925.1 Symbols: Auxin efflux carrier family protein chr1:7289973-7292507 FORWARD LENGTH=472
Eucgr.H01385.1	19,50	0,96	1,35	1,75	14,54	AT1G65295.1 Symbols: unknown protein;
Eucgr.J00741.2	10,27	0,13	0,71	0,82	6,34	AT3G51950.2 Symbols: Zinc finger (CCCH-type) family protein / RNA recognition motif (RRM)-containing protein
Eucgr.G00858.1	0,54	0,05	0,02	0,17	0,25	AT2G34690.1 Symbols: ACD11 Glycolipid transfer protein (GLTP) family protein
Eucgr.I01042.1	1,98	0,00	0,00	0,02	0,70	AT2G22990.1 Symbols: SNG1, SCPL8 sinapoylglucose 1 chr2:9786393-9789925 FORWARD LENGTH=433
Eucgr.K00901.1	0,83	0,03	0,98	1,18	1,51	AT1G03940.1 Symbols: HXXXD-type acyl-transferase family protein chr1:1009542-1010951 REVERSE LENGTH=469
Eucgr.E03836.1	23,59	3,23	5,20	10,23	17,29	AT5G47770.1 Symbols: FPS1 farnesyl diphosphate synthase 1 chr5:19345297-19347415 FORWARD LENGTH=384
Eucgr.D00128.1	5,05	0,18	0,46	0,86	3,25	AT4G20940.1 Symbols: Leucine-rich receptor-like protein kinase family protein
Eucgr.G02320.1	54,05	12,58	15,55	26,78	45,51	AT1G13380.1 Symbols: Protein of unknown function (DUF1218) chr1:4589218-4590362 REVERSE LENGTH=188
Eucgr.J01387.1	112,85	18,00	22,85	43,46	75,96	AT1G26355.1 Symbols: SP1L1 SPIRAL1-like1 chr1:9117950-9118611 REVERSE LENGTH=113
Eucgr.H04935.1	1,58	0,00	0,03	0,17	0,89	No hits found

Eucgr.J00011.1	8,59	1,18	1,09	1,64	6,23	AT3G12530.1 Symbols: PSF2 PSF2 chr3:3972604-3973864 REVERSE LENGTH=210
Eucgr.F04412.1	0,55	0,06	0,02	0,00	0,25	AT4G09770.2 Symbols: TRAF-like family protein chr4:6154534-6155859 REVERSE LENGTH=297
Eucgr.E00641.1	1,91	0,00	0,03	0,24	0,84	AT3G61460.1 Symbols: BRH1 brassinosteroid-responsive RING-H2 chr3:22741701-22742213 REVERSE LENGTH=170
Eucgr.I01393.1	2,27	156,93	131,40	134,97	6,62	AT2G17880.1 Symbols: Chaperone DnaJ-domain superfamily protein chr2:7767176-7767658 REVERSE LENGTH=160
Eucgr.F04055.2	102,46	11,17	19,96	31,01	67,83	AT3G13040.2 Symbols: myb-like HTH transcriptional regulator family protein
Eucgr.I01737.1	24,79	140,09	119,59	108,71	45,70	AT5G14500.1 Symbols: aldose 1-epimerase family protein chr5:4674503-4676368 REVERSE LENGTH=306
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.K02704.1	0,05	0,88	0,69	0,56	0,06	AT5G50700.1 Symbols: HSD1 hydroxysteroid dehydrogenase 1 chr5:20623259-20624995 REVERSE LENGTH=349
Eucgr.B01107.1	170,31	16,77	31,62	65,19	105,63	AT3G06035.1 Symbols: Glycoprotein membrane precursor GPI-anchored chr3:1823172-1824110 REVERSE LENGTH=200
Eucgr.F02909.1	0,66	0,00	0,02	0,03	1,35	AT1G65920.1 Symbols: Regulator of chromosome condensation (RCC1) family with FYVE zinc finger domain
Eucgr.G01970.1	1,27	12,23	6,28	3,64	2,57	AT5G13330.1 Symbols: Rap2.6L related to AP2 6l chr5:4272384-4274461 FORWARD LENGTH=212
Eucgr.E00177.1	2,68	12,37	11,79	9,56	4,38	AT1G01770.1 Symbols: unknown protein;
Eucgr.J01448.1	293,20	54,21	46,52	29,50	177,22	AT3G25572.1 Symbols: CPuORF11 conserved peptide upstream open reading frame 11
Eucgr.H04687.1	0,92	0,00	0,00	0,02	0,32	AT1G24430.1 Symbols: HXXXD-type acyl-transferase family protein chr1:8658190-8659497 REVERSE LENGTH=435
Eucgr.J00984.1	5,33	0,07	0,16	0,51	2,90	AT3G10080.1 Symbols: RmlC-like cupins superfamily protein chr3:3107476-3108159 REVERSE LENGTH=227
Eucgr.H04601.1	5,36	0,12	0,72	2,17	2,67	AT5G14070.1 Symbols: ROXY2 Thioredoxin superfamily protein chr5:4541915-4542337 FORWARD LENGTH=140
Eucgr.H00446.1	12,90	0,94	1,77	3,32	8,62	AT5G12890.1 Symbols: UDP-Glycosyltransferase superfamily protein chr5:4069658-4071124 REVERSE LENGTH=488
Eucgr.I02753.1	49,34	6,76	9,82	18,33	37,24	AT1G58070.1 Symbols: unknown protein;
Eucgr.C03749.1	2,33	14,24	13,94	10,11	5,14	AT1G35190.2 Symbols: 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein
Eucgr.C03181.1	30,87	83,50	73,81	89,50	25,44	AT5G20885.1 Symbols: RING/U-box superfamily protein chr5:7084133-7084663 REVERSE LENGTH=176
Eucgr.L00245.1	1,83	287,87	164,66	80,81	5,28	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.H02635.1	6,82	0,04	0,35	0,43	3,64	AT3G23740.1 Symbols: unknown protein;
Eucgr.J01371.2	4,99	0,70	1,03	1,45	3,69	AT5G15530.1 Symbols: BCCP2, CAC1-B biotin carboxyl carrier protein 2 chr5:5038955-5040437 FORWARD LENGTH=255
Eucgr.H01080.1	0,60	6,95	6,64	8,41	1,61	AT1G24020.2 Symbols: MLP423 MLP-like protein 423 chr1:8500653-8501458 REVERSE LENGTH=155
Eucgr.I01757.1	1,24	15,94	10,85	4,78	0,59	AT5G40390.1 Symbols: SIP1 Raffinose synthase family protein chr5:16161720-16165085 FORWARD LENGTH=783
Eucgr.L03103.1	4,74	0,00	0,09	0,47	2,39	AT1G12663.1 Symbols: Predicted to encode a PR (pathogenesis-related) protein.
Eucgr.L02809.1	3,37	0,21	0,81	1,00	2,25	AT5G26910.1 Symbols: unknown protein;
Eucgr.D02595.1	93,74	5,16	8,96	13,91	65,47	AT4G19120.2 Symbols: ERD3 S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
Eucgr.G02484.1	8,11	0,04	0,24	0,38	4,24	AT1G26761.1 Symbols: Arabinanase/levansucrase/invertase chr1:9250269-9251603 REVERSE LENGTH=444
Eucgr.D01668.1	63,62	0,94	2,21	3,72	41,65	AT1G32860.1 Symbols: Glycosyl hydrolase superfamily protein chr1:11907308-11908803 REVERSE LENGTH=426

Eucgr.H03535.1	23,32	2,21	3,07	2,75	17,46	AT3G22400.1 Symbols: LOX5 PLAT/LH2 domain-containing lipoxygenase family protein
Eucgr.H03413.1	0,22	9,42	7,54	6,04	0,54	AT2G26400.1 Symbols: ARD, ATARD3, ARD3 acireductone dioxygenase 3 chr2:11232058-11233274 REVERSE LENGTH=199
Eucgr.H01468.1	26,78	2,36	3,11	4,58	19,24	AT3G04860.1 Symbols: Plant protein of unknown function (DUF868) chr3:1339349-1340218 REVERSE LENGTH=289
Eucgr.B01196.1	1,29	0,02	0,09	0,32	0,68	AT3G52970.1 Symbols: CYP76G1 cytochrome P450, family 76, subfamily G, polypeptide 1
Eucgr.I00200.3	47,74	3,92	3,37	6,51	24,57	AT5G64040.1 Symbols: PSAN photosystem I reaction center subunit PSI-N, chloroplast, putative / PSI-N, putative (PSAN)
Eucgr.E03364.1	30,20	0,87	0,81	1,36	17,97	AT1G11120.2 Symbols: unknown protein;
Eucgr.K02869.1	32,32	4,32	8,35	11,07	25,57	AT3G48040.1 Symbols: ARAC8, ATROP10, ROP10, ATRAC8 RHO-related protein from plants 10
Eucgr.K02974.1	109,96	5,80	13,01	25,21	62,01	AT5G01360.1 Symbols: TBL3 Plant protein of unknown function (DUF828) chr5:147608-149316 REVERSE LENGTH=434
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.C02732.1	33,22	0,68	2,05	3,07	23,80	AT2G25270.1 Symbols: unknown protein;
Eucgr.C00797.1	72,88	7,59	11,44	10,90	34,37	AT2G05920.1 Symbols: Subtilase family protein chr2:2269831-2272207 REVERSE LENGTH=754
Eucgr.G02493.1	13,14	4,01	3,24	2,52	19,02	AT5G39660.2 Symbols: CDF2 cycling DOF factor 2 chr5:15878920-15880712 FORWARD LENGTH=457
Eucgr.J00105.1	2,70	20,19	18,01	14,17	5,09	AT2G38560.1 Symbols: TFIIS transcript elongation factor IIS chr2:16134802-16136319 FORWARD LENGTH=378
Eucgr.H01622.1	0,31	0,00	0,00	0,02	0,14	AT4G12910.1 Symbols: scpl20 serine carboxypeptidase-like 20 chr4:7550576-7553051 REVERSE LENGTH=497
Eucgr.A01860.1	6,07	0,07	0,25	0,34	3,31	AT5G46090.1 Symbols: Protein of unknown function (DUF679) chr5:18693485-18694129 FORWARD LENGTH=214
Eucgr.F03309.1	1,53	0,00	0,04	0,57	0,85	AT5G53850.2 Symbols: haloacid dehalogenase-like hydrolase family protein
Eucgr.D01819.1	80,34	4,38	7,93	13,49	59,59	AT1G63910.1 Symbols: AtMYB103, MYB103 myb domain protein 103 chr1:23719968-23721625 REVERSE LENGTH=370
Eucgr.B02453.1	13,56	1,98	3,25	3,47	10,76	AT2G28550.3 Symbols: RAP2.7 related to AP2.7 chr2:12226168-12228251 REVERSE LENGTH=464
Eucgr.K01387.2	3,33	26,78	22,63	18,02	2,56	AT3G07650.4 Symbols: COL9 CONSTANS-like 9 chr3:2442494-2443901 FORWARD LENGTH=372
Eucgr.I02211.1	1,65	70,32	142,40	284,67	0,90	AT1G20823.1 Symbols: RING/U-box superfamily protein chr1:7238880-7239473 FORWARD LENGTH=197
Eucgr.C00728.1	38,12	1,77	4,02	6,06	26,18	AT1G02180.1 Symbols: ferredoxin-related chr1:413619-414505 REVERSE LENGTH=226
Eucgr.B03079.1	0,46	7,39	8,38	9,72	1,15	AT1G13480.1 Symbols: Protein of unknown function (DUF1262) chr1:4622783-4624043 REVERSE LENGTH=387
Eucgr.L01116.1	3,48	0,20	0,43	0,79	2,08	AT5G44530.1 Symbols: Subtilase family protein chr5:17937931-17941193 FORWARD LENGTH=840
Eucgr.B03117.1	4,21	0,11	0,15	0,41	2,33	No hits found
Eucgr.K03327.1	2,94	7,53	10,15	7,92	5,01	AT3G63340.2 Symbols: Protein phosphatase 2C family protein chr3:23392181-23397999 REVERSE LENGTH=1075
Eucgr.D01344.1	9,02	0,04	0,07	0,04	4,47	AT4G22010.1 Symbols: sks4 SKU5 similar 4 chr4:11663429-11666463 FORWARD LENGTH=541
Eucgr.D00234.3	23,67	0,91	0,97	3,13	10,72	AT1G30380.1 Symbols: PSAK photosystem I subunit K chr1:10722325-10723013 FORWARD LENGTH=130
Eucgr.H01314.3	81,51	4,51	13,74	28,23	51,70	AT1G56720.3 Symbols: Protein kinase superfamily protein chr1:21263630-21265559 REVERSE LENGTH=492
Eucgr.K03183.1	1,01	0,00	0,28	0,35	1,90	AT2G37330.1 Symbols: ALS3 aluminum sensitive 3 chr2:15669190-15670172 FORWARD LENGTH=273
Eucgr.B03111.1	6,22	0,15	0,91	1,39	3,83	AT3G30380.2 Symbols: alpha/beta-Hydrolases superfamily protein chr3:11974424-11976571 FORWARD LENGTH=377

Eucgr.K01639.1	47,77	8,08	7,88	10,74	38,75	AT3G57040.1 Symbols: ARR9, ATRR4 response regulator 9 chr3:21110059-21111228 FORWARD LENGTH=234
Eucgr.K01853.1	15,53	0,92	1,43	1,67	10,58	AT3G58060.1 Symbols: Cation efflux family protein chr3:21497778-21499676 REVERSE LENGTH=411
Eucgr.E02504.1	2,20	0,00	0,09	0,13	1,42	No hits found
Eucgr.C00826.1	13,21	0,89	0,69	0,73	9,60	AT1G22640.1 Symbols: ATMYB3, MYB3 myb domain protein 3 chr1:8006289-8007219 FORWARD LENGTH=257
Eucgr.I02368.2	31,02	10,48	7,28	6,84	45,32	AT4G38960.1 Symbols: B-box type zinc finger family protein chr4:18161576-18163045 FORWARD LENGTH=183
Eucgr.I01572.1	2,80	0,46	0,51	0,47	1,99	AT3G26680.3 Symbols: SNM1 DNA repair metallo-beta-lactamase family protein
Eucgr.F01077.1	50,23	2,67	3,23	4,98	29,15	AT5G54270.1 Symbols: LHCB3, LHCB3*1 light-harvesting chlorophyll B-binding protein 3
Eucgr.H02038.1	1,00	0,00	0,02	0,21	0,55	AT5G42930.1 Symbols: alpha/beta-Hydrolases superfamily protein chr5:17210738-17214152 REVERSE LENGTH=467
Eucgr.K00655.1	13,42	1,65	2,00	2,91	9,97	AT2G05160.1 Symbols: CCCH-type zinc fingerfamily protein with RNA-binding domain
Eucgr.I00574.2	13,15	1,29	2,31	2,03	11,55	AT2G22620.1 Symbols: Rhamnogalacturonate lyase family protein chr2:9604902-9610156 REVERSE LENGTH=677
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H03420.1	59,57	1,15	1,08	1,40	23,29	AT3G51895.1 Symbols: SULTR3;1, AST12 sulfate transporter 3;1 chr3:19251503-19255677 REVERSE LENGTH=658
Eucgr.B03112.1	90,16	4,80	18,07	34,72	60,93	AT1G13635.2 Symbols: DNA glycosylase superfamily protein chr1:4674248-4675784 FORWARD LENGTH=311
Eucgr.D01901.1	0,59	11,36	5,67	3,21	1,34	AT5G59220.1 Symbols: HAI1 highly ABA-induced PP2C gene 1 chr5:23894672-23896497 REVERSE LENGTH=413
Eucgr.D01065.1	5,47	0,67	0,56	1,04	3,52	AT1G08930.2 Symbols: ERD6 Major facilitator superfamily protein chr1:2873604-2876979 FORWARD LENGTH=496
Eucgr.A01383.1	0,43	8,83	7,37	5,16	0,99	AT4G12680.1 Symbols: unknown protein;
Eucgr.K00892.1	1,68	0,00	0,02	0,02	0,77	AT1G18010.1 Symbols: Major facilitator superfamily protein chr1:6199980-6201359 FORWARD LENGTH=459
Eucgr.E00800.1	24,98	2,16	4,34	7,41	38,51	AT2G46150.1 Symbols: Late embryogenesis abundant (LEA) hydroxyproline-rich glycoprotein family
Eucgr.B00458.1	92,46	9,79	21,72	35,60	60,49	AT1G74690.1 Symbols: IQD31 IQ-domain 31 chr1:28061498-28063924 REVERSE LENGTH=587
Eucgr.H01946.1	2,08	8,74	16,45	8,26	4,46	AT3G50380.1 Symbols: Protein of unknown function (DUF1162) chr3:18686527-18700533 REVERSE LENGTH=3072
Eucgr.K03248.1	6,20	185,16	141,68	125,30	13,51	AT2G28400.1 Symbols: Protein of unknown function, DUF584 chr2:12148513-12149001 REVERSE LENGTH=162
Eucgr.G02119.1	120,61	17,32	38,68	69,41	90,39	AT1G14870.1 Symbols: PCR2 PLANT CADMIUM RESISTANCE 2 chr1:5128591-5129458 REVERSE LENGTH=152
Eucgr.K00177.1	2,04	0,00	0,04	0,05	0,95	AT1G65680.1 Symbols: ATEXPB2, EXPB2, ATHEXP BETA 1.4 expansin B2
Eucgr.J01893.1	23,61	0,43	0,97	1,32	13,26	AT2G30010.1 Symbols: TBL45 TRICHOME BIREFRINGENCE-LIKE 45 chr2:12805833-12809226 FORWARD LENGTH=398
Eucgr.J00309.1	2,26	0,04	0,16	0,49	1,74	AT3G03770.2 Symbols: Leucine-rich repeat protein kinase family protein chr3:945303-948436 REVERSE LENGTH=802
Eucgr.E00644.1	2,51	0,05	0,08	0,06	1,45	AT3G05170.1 Symbols: Phosphoglycerate mutase family protein chr3:1466738-1468219 FORWARD LENGTH=316
Eucgr.J02976.1	1,02	0,06	0,03	0,10	0,50	AT1G79890.1 Symbols: RAD3-like DNA-binding helicase protein chr1:30048655-30052203 FORWARD LENGTH=882
Eucgr.D01294.1	10,82	1,01	1,54	1,66	7,36	AT1G11545.1 Symbols: XTH8 xyloglucan endotransglucosylase/hydrolase 8 chr1:3878689-3880286 REVERSE LENGTH=305
Eucgr.B01744.1	0,34	0,00	0,02	0,13	0,18	AT5G54160.1 Symbols: ATOMT1, OMT1 O-methyltransferase 1 chr5:21982075-21984167 FORWARD LENGTH=363
Eucgr.A02802.1	0,30	11,17	20,62	25,92	0,14	No hits found

Eucgr.A02803.1	0,02	2,72	2,87	3,18	0,00	No hits found
Eucgr.I01808.1	0,03	1,35	1,33	0,98	0,00	AT1G23740.1 Symbols: Oxidoreductase, zinc-binding dehydrogenase family protein
Eucgr.J00290.1	85,69	14,15	29,81	56,15	76,26	AT5G04840.1 Symbols: bZIP protein chr5:1406005-1407648 FORWARD LENGTH=307
Eucgr.J00838.1	0,06	0,80	0,40	0,31	0,13	AT1G23030.1 Symbols: ARM repeat superfamily protein chr1:8156745-8158842 FORWARD LENGTH=612
Eucgr.L00241.1	0,02	8,29	5,04	1,18	0,00	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.F02332.1	7,23	19,30	17,84	15,66	10,47	AT3G49180.1 Symbols: RID3 Transducin/WD40 repeat-like superfamily protein
Eucgr.H05077.1	39,05	7,29	16,20	30,22	32,75	AT3G05270.2 Symbols: Plant protein of unknown function (DUF869) chr3:1500803-1502926 REVERSE LENGTH=615
Eucgr.C00212.1	27,23	0,59	0,67	0,98	14,51	AT4G30410.1 Symbols: sequence-specific DNA binding transcription factors
Eucgr.F00650.1	42,22	3,35	5,96	9,14	32,11	AT5G55230.1 Symbols: ATMAP65-1, MAP65-1 microtubule-associated proteins 65-1
Eucgr.C04173.1	39,88	12,41	10,96	16,18	35,36	AT4G17486.1 Symbols: PPPDE putative thiol peptidase family protein chr4:9749992-9751201 REVERSE LENGTH=224
Eucgr.G02236.1	7,69	21,24	21,64	19,84	11,07	AT1G49540.1 Symbols: ELP2, AtELP2 elongator protein 2 chr1:18333767-18337382 REVERSE LENGTH=838
Eucgr.A01426.1	1,11	16,49	10,30	8,79	2,09	No hits found
GENE ID	FPKM					Annotation

	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.H04616.1	8,31	0,40	0,52	0,64	5,00	AT1G67320.2 Symbols: DNA primase, large subunit family chr1:25205520-25208966 REVERSE LENGTH=454
Eucgr.L01040.1	25,93	131,16	128,06	85,19	43,84	AT2G12400.1 Symbols: unknown protein;
Eucgr.J00129.2	5,26	0,38	0,37	0,99	3,07	AT3G56290.1 Symbols: unknown protein;
Eucgr.H00366.1	108,13	13,22	11,41	12,49	62,24	AT1G06040.1 Symbols: STO B-box zinc finger family protein chr1:1828662-1829659 REVERSE LENGTH=248
Eucgr.D00408.2	0,38	0,00	0,00	0,01	0,25	AT5G45650.1 Symbols: subtilase family protein chr5:18513520-18518790 REVERSE LENGTH=791
Eucgr.J01434.1	6,83	0,06	0,19	0,44	3,55	No hits found
Eucgr.I02455.1	14,29	1,86	1,93	2,06	10,28	AT5G66750.1 Symbols: DDM1, CHR01, CHR1, CHA1, SOM4, SOM1, ATDDM1 chromatin remodeling 1
Eucgr.B03245.1	6,37	33,40	31,78	22,98	10,70	AT5G14420.2 Symbols: RGLG2 RING domain ligase2 chr5:4648355-4650563 REVERSE LENGTH=468
Eucgr.I02090.1	11,49	0,17	2,19	3,07	6,60	AT3G30530.1 Symbols: ATBZIP42, bZIP42 basic leucine-zipper 42 chr3:12139512-12140033 FORWARD LENGTH=173
Eucgr.F04356.1	16,92	0,00	0,14	0,24	11,14	AT1G44760.1 Symbols: Adenine nucleotide alpha hydrolases-like superfamily protein
Eucgr.G03317.1	4,73	0,06	0,06	0,07	2,77	AT5G12470.1 Symbols: Protein of unknown function (DUF3411) chr5:4044950-4047290 REVERSE LENGTH=386
Eucgr.E02667.1	78,18	13,06	25,76	45,74	59,90	AT2G33990.1 Symbols: iqd9 IQ-domain 9 chr2:14360502-14361630 REVERSE LENGTH=263
Eucgr.A01039.1	0,89	0,00	0,03	0,04	0,53	AT5G48170.1 Symbols: SLY2 F-box family protein chr5:19532729-19533202 REVERSE LENGTH=157
Eucgr.J02254.1	4,42	0,71	0,43	0,46	2,51	AT5G14490.1 Symbols: anac085, NAC085 NAC domain containing protein 85
Eucgr.A00996.1	15,85	0,99	1,76	2,42	11,61	AT4G01680.3 Symbols: MYB55 myb domain protein 55 chr4:716401-717415 REVERSE LENGTH=241
Eucgr.A02890.1	0,23	8,26	4,84	2,03	0,16	AT5G23950.1 Symbols: Calcium-dependent lipid-binding (CaLB domain) family protein
Eucgr.B01827.1	5,07	0,16	0,81	1,35	3,68	AT4G01680.3 Symbols: MYB55 myb domain protein 55 chr4:716401-717415 REVERSE LENGTH=241

Eucgr.I00052.1	5,17	0,77	0,64	0,49	8,13	AT5G08720.1 Symbols: unknown protein
Eucgr.A01373.1	2,40	16,15	18,85	18,13	4,53	No hits found
Eucgr.D01123.1	8,79	0,10	0,30	0,31	4,77	AT3G17350.1 Symbols: unknown protein;
Eucgr.J01758.2	14,12	1,01	0,70	1,08	9,09	AT1G55480.1 Symbols: ZKT protein containing PDZ domain, a K-box domain, and a TPR region
Eucgr.I00229.1	2,36	0,08	0,08	0,16	1,34	No hits found
Eucgr.G01772.1	67,33	1,02	3,12	4,42	40,59	AT4G38660.1 Symbols: Pathogenesis-related thaumatin superfamily protein
Eucgr.D02287.1	5,85	0,12	0,42	0,64	4,19	AT1G63650.3 Symbols: EGL3 basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.F04348.1	7,69	2,28	2,27	2,42	6,21	AT1G77580.2 Symbols: Plant protein of unknown function (DUF869) chr1:29144191-29146793 REVERSE LENGTH=779
Eucgr.D00536.1	9,72	0,76	1,09	1,87	7,05	AT5G45920.1 Symbols: SGNH hydrolase-type esterase superfamily protein
Eucgr.D02360.1	7,94	0,05	0,02	0,16	3,87	AT5G53200.1 Symbols: TRY Homeodomain-like superfamily protein chr5:21582913-21583960 FORWARD LENGTH=106
Eucgr.K01100.1	0,20	27,44	17,99	3,77	0,43	AT2G42560.1 Symbols: late embryogenesis abundant domain-containing protein / LEA domain-containing protein
Eucgr.D02070.1	1,27	25,95	13,55	4,74	0,68	AT4G25200.1 Symbols: ATHSP23.6-MITO, HSP23.6-MITO mitochondrion-localized small heat shock protein 23.6
Eucgr.E02892.1	0,77	0,03	0,07	0,29	0,52	AT1G17020.1 Symbols: SRG1, ATSRG1 senescence-related gene 1 chr1:5820258-5821741 FORWARD LENGTH=358
Eucgr.F02333.1	1,19	0,04	0,08	0,37	0,78	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.I01048.1	7,39	0,30	0,43	0,43	5,01	AT1G70210.1 Symbols: CYCD1;1, ATCYCD1;1 CYCLIN D1;1 chr1:26440015-26441980 FORWARD LENGTH=339
Eucgr.J02034.1	2,07	17,67	20,57	28,18	1,51	No hits found
Eucgr.B02872.1	4,92	0,56	1,01	1,32	3,73	AT1G15170.1 Symbols: MATE efflux family protein chr1:5220690-5222756 FORWARD LENGTH=481
Eucgr.E03358.1	24,39	4,93	8,11	10,91	20,70	AT1G61100.1 Symbols: disease resistance protein (TIR class), putative chr1:22508831-22512122 REVERSE LENGTH=808
Eucgr.K02268.1	30,11	103,55	90,24	79,91	43,26	No hits found
Eucgr.L01175.1	0,63	0,04	0,06	0,25	0,44	AT5G63020.1 Symbols: Disease resistance protein (CC-NBS-LRR class) family
Eucgr.J02694.1	1,36	0,04	0,02	0,14	0,72	AT5G51260.1 Symbols: HAD superfamily, subfamily IIIB acid phosphatase
Eucgr.F00357.1	57,27	1,46	2,05	2,82	35,06	AT4G08950.1 Symbols: EXO Phosphate-responsive 1 family protein chr4:5740378-5741322 FORWARD LENGTH=314
Eucgr.K00384.1	9,00	0,28	0,64	0,99	6,32	AT4G00990.1 Symbols: Transcription factor jumonji (jmc) domain-containing protein
Eucgr.E02931.1	0,11	5,24	1,25	0,63	0,26	AT1G16770.1 Symbols: unknown protein;
Eucgr.J01057.1	28,85	0,98	0,40	0,82	18,36	AT3G07510.2 Symbols: unknown protein;
Eucgr.H00037.1	1,32	0,02	0,10	0,09	0,83	AT2G01120.1 Symbols: ORC4, ATORC4 origin recognition complex subunit 4
Eucgr.H04541.2	31,01	2,69	7,10	13,17	24,12	AT1G28570.1 Symbols: SGNH hydrolase-type esterase superfamily protein
Eucgr.G02885.1	137,27	2,76	9,13	23,45	108,13	AT1G09070.1 Symbols: SRC2, (AT)SRC2 soybean gene regulated by cold-2 chr1:2927767-2928741 FORWARD LENGTH=324
Eucgr.F02583.1	5,14	31,22	29,00	22,63	3,68	AT1G22640.1 Symbols: ATMYB3, MYB3 myb domain protein 3 chr1:8006289-8007219 FORWARD LENGTH=257

Eucgr.B00225.1	0,00	0,17	0,21	0,08	0,00	AT5G50170.1 Symbols: C2 calcium/lipid-binding and GRAM domain containing protein
Eucgr.E00198.1	0,02	3,23	2,24	0,60	0,00	AT2G45760.1 Symbols: BAP2, BAL BON association protein 2 chr2:18847125-18847748 REVERSE LENGTH=207
Eucgr.L02067.1	0,02	1,90	1,51	0,18	0,00	No hits found
Eucgr.F02697.1	0,02	1,22	1,11	0,65	0,00	No hits found
Eucgr.K00891.1	0,60	0,02	0,02	0,01	0,36	AT1G18010.1 Symbols: Major facilitator superfamily protein chr1:6199980-6201359 FORWARD LENGTH=459
Eucgr.G03224.1	4,84	19,98	15,95	17,07	7,02	AT3G06730.1 Symbols: TRX P, TRX z Thioredoxin z chr3:2124276-2125845 FORWARD LENGTH=183
Eucgr.D00314.1	9,08	0,82	0,89	1,04	6,04	AT5G49010.1 Symbols: SLD5 DNA replication protein-related chr5:19866464-19868074 FORWARD LENGTH=220
Eucgr.E03993.2	0,09	1,57	1,73	3,13	0,10	AT4G04330.1 Symbols: Chaperonin-like RbcX protein chr4:2116700-2118563 REVERSE LENGTH=174
Eucgr.I02808.1	37,46	7,02	12,32	24,06	32,73	AT2G31930.1 Symbols: unknown protein;
Eucgr.G00660.1	6,11	15,46	16,90	15,98	8,68	AT3G26410.1 Symbols: TRM11, AtTRM11 methyltransferases;nucleic acid binding
Eucgr.E04327.1	11,72	0,18	0,28	0,41	7,85	AT5G44680.1 Symbols: DNA glycosylase superfamily protein chr5:18024461-18025893 REVERSE LENGTH=353
Eucgr.C01374.1	3,70	0,00	0,39	0,75	6,50	AT5G52600.1 Symbols: AtMYB82, MYB82 myb domain protein 82 chr5:21343197-21343968 REVERSE LENGTH=201
Eucgr.A02016.1	9,94	0,13	0,85	1,93	6,84	AT4G18260.1 Symbols: Cytochrome b561/ferric reductase transmembrane protein family
Eucgr.G02426.1	11,24	0,16	0,50	1,03	7,23	AT1G69080.1 Symbols: Adenine nucleotide alpha hydrolases-like superfamily protein
Eucgr.I00735.1	53,06	0,45	0,88	0,65	36,95	AT2G39540.1 Symbols: Gibberellin-regulated family protein chr2:16500866-16501241 FORWARD LENGTH=87
Eucgr.J00725.1	0,11	6,36	3,63	1,74	0,11	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.E01052.1	22,26	0,53	1,73	3,34	16,27	AT1G64690.1 Symbols: BLT branchless trichome chr1:24038069-24038890 FORWARD LENGTH=273
Eucgr.K00441.1	1,79	0,04	0,25	0,41	1,19	AT5G38280.1 Symbols: PR5K PR5-like receptor kinase chr5:15293325-15295838 REVERSE LENGTH=665
Eucgr.B03158.1	0,20	3,04	2,65	1,65	0,42	No hits found
Eucgr.G02486.1	0,39	3,91	2,48	1,76	0,71	AT1G69490.1 Symbols: NAP, ANAC029, ATNAP NAC-like, activated by AP3/PI
Eucgr.F02085.1	19,98	0,09	0,55	1,20	12,89	AT1G75590.1 Symbols: SAUR-like auxin-responsive protein family chr1:28383250-28383714 REVERSE LENGTH=154
Eucgr.G02347.5	17,99	2,45	2,54	3,97	12,90	AT1G68830.1 Symbols: STN7 STT7 homolog STN7 chr1:25872654-25875473 REVERSE LENGTH=562
Eucgr.I02161.1	10,99	1,54	2,13	5,30	9,49	AT3G14280.1 Symbols: unknown protein;
Eucgr.D01725.1	1,95	0,18	0,19	0,16	1,40	AT5G41850.1 Symbols: alpha/beta-Hydrolases superfamily protein chr5:16756698-16757791 REVERSE LENGTH=224
Eucgr.H02963.1	70,93	3,44	2,65	3,31	47,81	AT2G05620.1 Symbols: PGR5 proton gradient regulation 5 chr2:2081204-2081687 REVERSE LENGTH=133
Eucgr.C00105.1	29,83	7,01	7,08	11,79	24,54	AT4G29930.1 Symbols: basic helix-loop-helix (bHLH) DNA-binding superfamily protein
Eucgr.H04382.1	22,55	1,11	3,43	4,89	18,19	AT3G04810.2 Symbols: ATNEK2, NEK2 NIMA-related kinase 2 chr3:1318096-1321101 FORWARD LENGTH=606
Eucgr.C00725.1	13,96	0,00	0,16	0,35	8,81	AT3G13540.1 Symbols: ATMYB5, MYB5 myb domain protein 5 chr3:4420239-4421443 FORWARD LENGTH=249
Eucgr.C02807.1	4,74	0,08	0,54	2,81	8,18	AT3G15680.1 Symbols: Ran BP2/NZF zinc finger-like superfamily protein chr3:5315437-5316048 FORWARD LENGTH=164

Eucgr.I01248.1	14,63	3,98	4,37	5,55	12,42	AT4G36750.1 Symbols: Quinone reductase family protein chr4:17324642-17326215 FORWARD LENGTH=273
Eucgr.F02320.1	101,08	0,80	3,34	5,57	85,41	AT1G46480.1 Symbols: WOX4 WUSCHEL related homeobox 4 chr1:17236903-17237953 REVERSE LENGTH=251
Eucgr.B03435.1	2,43	0,05	0,21	0,36	1,77	AT1G26840.1 Symbols: ORC6, ATORC6 origin recognition complex protein 6
Eucgr.K02194.1	3,98	19,91	15,22	16,23	5,89	AT5G50930.1 Symbols: Histone superfamily protein chr5:20722069-20724317 FORWARD LENGTH=242
Eucgr.A02563.1	12,92	1,21	1,83	3,51	10,92	AT5G01420.1 Symbols: Glutaredoxin family protein chr5:174886-176091 REVERSE LENGTH=401
Eucgr.H01076.1	34,40	91,65	87,28	88,04	48,24	AT3G06760.1 Symbols: Drought-responsive family protein chr3:2132971-2134355 FORWARD LENGTH=224
Eucgr.J01082.1	42,67	2,92	7,55	12,94	33,44	AT5G03250.1 Symbols: Phototropic-responsive NPH3 family protein chr5:774591-776855 FORWARD LENGTH=592
Eucgr.I00762.1	6,66	0,81	1,22	1,77	5,56	AT5G67280.1 Symbols: RLK receptor-like kinase chr5:26842430-26845126 REVERSE LENGTH=751
Eucgr.K00929.1	17,01	1,98	2,92	3,77	14,14	AT4G28220.1 Symbols: NDB1 NAD(P)H dehydrogenase B1 chr4:13993078-13995651 FORWARD LENGTH=571
Eucgr.H04081.1	78,04	0,92	1,14	1,64	49,01	No hits found
Eucgr.K02470.1	0,61	19,17	10,27	4,78	0,98	AT3G47600.1 Symbols: MYB94, ATMYBCP70, ATMYB94 myb domain protein 94
Eucgr.J01901.1	66,36	6,44	19,72	43,79	54,74	AT3G51780.1 Symbols: ATBAG4, BAG4 BCL-2-associated athanogene 4 chr3:19207029-19208178 REVERSE LENGTH=269
Eucgr.B03203.1	0,04	1,66	1,03	0,34	0,06	AT1G69480.1 Symbols: EXS (ERD1/XPR1/SYG1) family protein chr1:26114187-26117479 REVERSE LENGTH=777
Eucgr.G02011.1	0,00	0,59	0,34	0,09	0,00	AT4G13420.1 Symbols: HAK5, ATHAK5 high affinity K+ transporter 5 chr4:7797038-7802174 REVERSE LENGTH=785
Eucgr.A00205.1	0,00	4,14	2,98	0,70	0,00	AT3G62560.1 Symbols: Ras-related small GTP-binding family protein chr3:23137539-23138880 FORWARD LENGTH=193
Eucgr.F02242.1	0,00	7,53	3,96	0,98	0,00	AT1G27461.1 Symbols: unknown protein;
Eucgr.K00224.1	0,00	0,47	0,27	0,19	0,00	AT4G23030.1 Symbols: MATE efflux family protein chr4:12072857-12074365 FORWARD LENGTH=502
Eucgr.F01607.1	0,00	1,71	1,16	0,04	0,00	AT1G75490.1 Symbols: Integrase-type DNA-binding superfamily protein
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.L03185.1	0,00	5,78	2,76	1,18	0,00	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.A01560.1	0,00	1,45	2,30	1,08	0,00	AT3G10020.2 Symbols: unknown protein;
Eucgr.B01991.1	0,00	15,76	9,07	3,42	0,00	AT1G54860.1 Symbols: Glycoprotein membrane precursor GPI-anchored chr1:20457522-20458434 REVERSE LENGTH=200
Eucgr.G00593.1	0,00	0,57	0,11	0,00	0,00	AT3G14690.1 Symbols: CYP72A15 cytochrome P450, family 72, subfamily A, polypeptide 15
Eucgr.B01792.1	0,00	2,16	0,51	0,00	0,00	AT1G56600.1 Symbols: AtGolS2, GolS2 galactinol synthase 2 chr1:21207620-21209291 FORWARD LENGTH=335
Eucgr.E02621.1	0,00	4,20	2,55	0,21	0,00	AT2G18540.1 Symbols: RmlC-like cupins superfamily protein chr2:8042382-8045008 REVERSE LENGTH=707
Eucgr.C00859.1	0,00	1,60	0,83	0,28	0,00	AT5G49690.1 Symbols: UDP-Glycosyltransferase superfamily protein chr5:20189968-20191350 REVERSE LENGTH=460
Eucgr.I02269.1	0,00	9,89	1,64	0,00	0,00	AT3G12500.1 Symbols: ATHCHIB, PR3, PR-3, CHI-B, B-CHI, HCHIB basic chitinase
Eucgr.J00396.1	0,00	1,28	0,82	0,48	0,00	AT1G08170.1 Symbols: Histone superfamily protein chr1:2562941-2563672 REVERSE LENGTH=243
Eucgr.F02406.1	0,00	0,59	0,17	0,06	0,00	AT4G31940.1 Symbols: CYP82C4 cytochrome P450, family 82, subfamily C, polypeptide 4
Eucgr.E02381.1	583,82	14,25	21,98	52,95	210,10	AT2G34430.1 Symbols: LHB1B1, LHCb1.4 light-harvesting chlorophyll-protein complex II subunit B1

Eucgr.K03437.1	12,80	0,59	0,65	0,65	9,94	No hits found
Eucgr.A01301.1	9,53	21,38	24,00	24,64	12,97	AT5G11240.1 Symbols: transducin family protein / WD-40 repeat family protein
Eucgr.H04316.1	101,04	26,20	35,12	40,53	84,26	AT1G54460.1 Symbols: TPX2 (targeting protein for Xklp2) protein family chr1:20340907-20342900 REVERSE LENGTH=338
Eucgr.F04176.1	7,68	37,72	32,00	34,90	11,57	AT1G71697.1 Symbols: ATCK1, CK, CK1 choline kinase 1 chr1:26971537-26973177 FORWARD LENGTH=346
Eucgr.K01523.1	2,70	10,18	10,24	12,48	2,06	AT5G35220.1 Symbols: EGY1 Peptidase M50 family protein chr5:13484606-13487546 REVERSE LENGTH=548
Eucgr.F03689.1	42,16	9,06	13,58	12,25	55,76	AT5G55730.2 Symbols: FLA1 FASCICLIN-like arabinogalactan 1 chr5:22558375-22560392 REVERSE LENGTH=424
Eucgr.K00027.1	0,31	0,00	0,00	0,10	0,22	AT4G18250.1 Symbols: receptor serine/threonine kinase, putative chr4:10087343-10091963 REVERSE LENGTH=853
Eucgr.J02152.1	0,53	0,00	0,01	0,02	0,39	AT5G40030.1 Symbols: Protein kinase superfamily protein chr5:16026227-16028283 FORWARD LENGTH=499
Eucgr.I01028.1	25,13	0,20	0,32	0,07	29,67	AT2G23690.1 Symbols: unknown protein;
Eucgr.A02070.1	1,00	17,38	8,86	5,29	1,76	AT3G04070.1 Symbols: anac047, NAC047 NAC domain containing protein 47
Eucgr.K01575.1	0,71	0,08	0,04	0,11	0,52	AT5G03960.1 Symbols: IQD12 IQ-domain 12 chr5:1068333-1069683 REVERSE LENGTH=403
Eucgr.E03963.1	0,50	0,03	0,03	0,07	0,37	AT1G61370.1 Symbols: S-locus lectin protein kinase family protein chr1:22642096-22645147 REVERSE LENGTH=814
Eucgr.H01178.1	3,52	0,19	0,21	0,55	2,53	AT1G02970.1 Symbols: WEE1, ATWEE1 WEE1 kinase homolog chr1:673408-676127 FORWARD LENGTH=500
Eucgr.H04752.1	1,96	0,00	0,00	0,12	1,40	AT5G10310.1 Symbols: unknown protein;
Eucgr.F01004.1	0,15	3,50	4,05	3,97	0,17	No hits found
Eucgr.G01335.1	15,84	0,00	0,18	0,60	13,02	AT2G15220.1 Symbols: Plant basic secretory protein (BSP) family protein chr2:6608689-6609366 FORWARD LENGTH=225
Eucgr.K00375.1	4,48	0,16	0,16	0,39	3,38	AT5G24270.2 Symbols: SOS3 Calcium-binding EF-hand family protein chr5:8238781-8240179 REVERSE LENGTH=222
Eucgr.K02858.1	218,59	19,50	43,49	66,15	141,91	AT3G48000.1 Symbols: ALDH2B4, ALDH2, ALDH2A aldehyde dehydrogenase 2B4
Eucgr.J00814.1	0,86	39,56	39,35	42,76	0,63	AT3G20810.1 Symbols: JMJD5 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein
Eucgr.H02790.1	3,07	0,05	0,22	0,28	2,33	AT4G31980.1 Symbols: unknown protein;
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.B00400.3	49,34	23,50	19,02	18,39	42,32	AT5G62670.1 Symbols: AHA11, HA11 H(+)-ATPase 11 chr5:25159495-25164957 FORWARD LENGTH=956
Eucgr.H04727.1	2,46	0,12	0,11	0,24	1,87	AT5G65400.1 Symbols: alpha/beta-Hydrolases superfamily protein chr5:26131448-26133062 FORWARD LENGTH=252
Eucgr.L01215.1	0,52	0,03	0,02	0,04	0,40	AT5G60900.1 Symbols: RLK1 receptor-like protein kinase 1 chr5:24498467-24501494 REVERSE LENGTH=748
Eucgr.C00184.1	45,26	0,45	0,72	0,22	37,20	AT4G25810.1 Symbols: XTR6, XTH23 xyloglucan endotransglycosylase 6
Eucgr.B02604.1	1,57	0,16	0,20	0,33	1,32	AT5G60900.1 Symbols: RLK1 receptor-like protein kinase 1 chr5:24498467-24501494 REVERSE LENGTH=748
Eucgr.F02442.1	0,37	0,00	0,00	0,03	0,27	AT1G78060.1 Symbols: Glycosyl hydrolase family protein chr1:29349796-29352868 REVERSE LENGTH=767
Eucgr.G01033.1	4,60	0,00	0,64	1,92	3,56	AT3G27890.1 Symbols: NQR NADPH:quinone oxidoreductase chr3:10350807-10351938 REVERSE LENGTH=196
Eucgr.K01122.1	16,74	1,68	3,16	4,48	14,64	AT2G42580.1 Symbols: TTL3, VIT tetraatricopeptide-repeat thioredoxin-like 3
Eucgr.J03079.2	4,27	0,21	0,24	0,57	3,52	AT1G52510.1 Symbols: alpha/beta-Hydrolases superfamily protein chr1:19563039-19565260 REVERSE LENGTH=380

Eucgr.J02006.1	0,69	0,03	0,03	0,00	1,06	AT1G07440.1 Symbols: NAD(P)-binding Rossmann-fold superfamily protein chr1:2286436-2287665 REVERSE LENGTH=266
Eucgr.K03116.2	21,97	1,81	0,90	0,98	20,94	AT5G02230.2 Symbols: Haloacid dehalogenase-like hydrolase (HAD) superfamily protein
Eucgr.H04927.1	0,64	0,02	0,08	0,08	0,51	AT5G60900.1 Symbols: RLK1 receptor-like protein kinase 1 chr5:24498467-24501494 REVERSE LENGTH=748
Eucgr.C04145.1	0,36	0,02	0,06	0,10	0,29	AT2G26910.1 Symbols: PDR4, ATPDR4 pleiotropic drug resistance 4 chr2:11481623-11487874 FORWARD LENGTH=1420
Eucgr.J01268.1	4,76	0,04	0,38	0,46	4,09	AT3G63530.2 Symbols: BB, BB2 RING/U-box superfamily protein chr3:23456407-23457787 REVERSE LENGTH=248
Eucgr.F00223.1	2,30	63,00	63,86	76,36	2,12	AT5G42900.2 Symbols: cold regulated gene 27 chr5:17198366-17199712 REVERSE LENGTH=246
Eucgr.A00882.1	246,28	23,21	64,91	87,97	141,86	AT1G79620.1 Symbols: Leucine-rich repeat protein kinase family protein chr1:29957633-29962174 REVERSE LENGTH=971
Eucgr.C00960.1	24,76	1,91	4,06	5,27	21,47	AT4G29240.1 Symbols: Leucine-rich repeat (LRR) family protein chr4:14418826-14420073 FORWARD LENGTH=415
Eucgr.B02677.1	1,47	0,03	0,05	0,07	1,24	AT1G14820.3 Symbols: Sec14p-like phosphatidylinositol transfer family protein
Eucgr.A01353.1	0,50	0,03	0,00	0,04	0,45	AT4G08850.1 Symbols: Leucine-rich repeat receptor-like protein kinase family protein
Eucgr.I00190.2	27,64	1,25	1,27	1,92	23,73	AT2G13610.1 Symbols: ABC-2 type transporter family protein chr2:5673827-5675776 REVERSE LENGTH=649
Eucgr.G00960.1	14,22	0,68	1,32	2,31	12,29	AT5G64330.1 Symbols: NPH3, RPT3, JK218 Phototropic-responsive NPH3 family protein
Eucgr.J02113.1	25,87	3,02	5,53	8,49	22,64	AT2G28550.3 Symbols: RAP2.7 related to AP2.7 chr2:12226168-12228251 REVERSE LENGTH=464
Eucgr.E03919.1	0,57	25,27	9,68	0,90	0,82	AT5G47550.1 Symbols: Cystatin/monellin superfamily protein chr5:19286596-19286964 REVERSE LENGTH=122
Eucgr.E02995.1	4,23	21,74	21,81	19,90	5,59	No hits found
Eucgr.J02625.1	6,60	0,33	0,53	0,75	5,93	AT5G11000.1 Symbols: Plant protein of unknown function (DUF868) chr5:3479166-3480335 REVERSE LENGTH=389
Eucgr.B03632.1	2,65	13,13	15,32	17,69	3,48	AT2G34860.2 Symbols: EDA3 DnaJ/Hsp40 cysteine-rich domain superfamily protein
Eucgr.B02605.1	0,48	0,02	0,09	0,10	0,40	AT5G60900.1 Symbols: RLK1 receptor-like protein kinase 1 chr5:24498467-24501494 REVERSE LENGTH=748
Eucgr.C03960.1	1,22	0,04	0,31	0,52	1,02	AT3G24430.1 Symbols: HCF101 ATP binding chr3:8868731-8872154 REVERSE LENGTH=532
Eucgr.C02507.1	1,32	7,47	6,07	6,63	1,84	AT4G30720.1 Symbols: FAD/NAD(P)-binding oxidoreductase family protein
Eucgr.K00883.2	607,60	16,06	29,74	37,61	288,85	AT4G03210.1 Symbols: XTH9 xyloglucan endotransglucosylase/hydrolase 9
Eucgr.K02468.1	66,99	551,20	399,69	392,53	97,85	AT1G18800.1 Symbols: NRP2 NAP1-related protein 2 chr1:6481466-6483463 REVERSE LENGTH=256
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.B02479.5	8,67	34,19	31,92	29,91	11,70	AT3G45300.1 Symbols: IVD, ATIVD, IVDH isovaleryl-CoA-dehydrogenase chr3:16621659-16624848 REVERSE LENGTH=409
Eucgr.C02516.1	1,62	0,13	0,00	0,08	1,40	AT4G30845.1 Symbols: unknown protein;
Eucgr.F02642.1	0,27	0,00	0,02	0,04	0,23	AT5G09360.1 Symbols: LAC14 laccase 14 chr5:2906426-2908658 REVERSE LENGTH=569
Eucgr.A01352.1	0,57	0,03	0,02	0,01	0,60	AT4G08850.1 Symbols: Leucine-rich repeat receptor-like protein kinase family protein
Eucgr.A01447.2	14,35	4,98	5,05	8,17	13,76	AT4G27800.1 Symbols: TAP38, PPH1 thylakoid-associated phosphatase 38
Eucgr.D00397.1	0,52	0,06	0,04	0,06	0,44	AT5G45470.1 Symbols: Protein of unknown function (DUF594) chr5:18422164-18424764 REVERSE LENGTH=866
Eucgr.B03360.1	12,37	0,93	1,94	2,24	11,59	AT1G70230.1 Symbols: TBL27 TRICHOME BIREFRINGENCE-LIKE 27 chr1:26450389-26451724 FORWARD LENGTH=416

Eucgr.G03419.1	1,67	0,10	0,21	0,20	1,46	AT2G31900.1 Symbols: XIF, ATXIF, ATMYO5 myosin-like protein XIF chr2:13560760-13569623 REVERSE LENGTH=1556
Eucgr.H01685.1	60,58	12,89	16,34	18,83	50,93	AT4G02630.1 Symbols: Protein kinase superfamily protein chr4:1151683-1153161 FORWARD LENGTH=492
Eucgr.I00651.1	23,67	74,34	61,77	52,40	30,88	AT5G65720.1 Symbols: ATNIF51, NIF51, NFS1, ATNFS1 nitrogen fixation S (NIFS)-like 1
Eucgr.K02098.1	1,29	0,00	0,22	0,47	1,85	AT4G12970.1 Symbols: STOMAGEN, EPFL9 stomagen chr4:7586244-7586798 REVERSE LENGTH=102
Eucgr.K02948.1	2,31	0,10	0,03	0,31	2,14	AT3G51660.1 Symbols: Tautomerase/MIF superfamily protein chr3:19164229-19165076 REVERSE LENGTH=112
Eucgr.F02749.1	0,61	53,80	22,67	6,72	0,82	AT1G21460.1 Symbols: SWEET1, AtSWEET1 Nodulin MtN3 family protein chr1:7512030-7513281 REVERSE LENGTH=247
Eucgr.C02594.1	28,57	4,87	4,63	5,60	25,81	AT2G24270.4 Symbols: ALDH11A3 aldehyde dehydrogenase 11A3 chr2:10327109-10329601 REVERSE LENGTH=503
Eucgr.H00494.1	6,14	20,25	26,24	27,22	7,22	AT5G26570.1 Symbols: PWD, OK1, ATGWD3 catalytics;carbohydrate kinases;phosphoglucan, water dikinases
Eucgr.A01640.1	4,07	0,08	0,22	0,57	3,60	AT4G20990.1 Symbols: ATACA4, ACA4 alpha carbonic anhydrase 4 chr4:11219772-11221126 FORWARD LENGTH=267
Eucgr.L01886.1	0,66	20,69	7,89	2,80	0,82	AT5G51440.1 Symbols: HSP20-like chaperones superfamily protein chr5:20891242-20892013 FORWARD LENGTH=210
Eucgr.B02634.1	32,97	2,94	6,54	9,05	31,72	AT1G14630.1 Symbols: unknown protein;
Eucgr.H02330.1	22,76	4,23	3,01	3,00	20,90	AT4G37730.1 Symbols: AtbZIP7, bZIP7 basic leucine-zipper 7 chr4:17723800-17724717 FORWARD LENGTH=305
Eucgr.F00464.1	28,18	85,70	85,04	82,49	36,89	AT1G09960.1 Symbols: SUT4, ATSUT4, SUC4, ATSUC4 sucrose transporter 4
Eucgr.L00920.1	0,57	0,03	0,01	0,04	0,56	AT3G49340.1 Symbols: Cysteine proteinases superfamily protein chr3:18293347-18294577 REVERSE LENGTH=341
Eucgr.L02477.1	3,59	0,66	0,34	0,83	5,03	AT2G31900.1 Symbols: XIF, ATXIF, ATMYO5 myosin-like protein XIF chr2:13560760-13569623 REVERSE LENGTH=1556
Eucgr.K00240.1	0,61	15,71	20,30	32,17	0,72	No hits found
Eucgr.B02498.1	113,87	14,29	40,05	58,07	104,95	AT2G28410.1 Symbols: unknown protein;
Eucgr.J00254.2	102,60	8,85	18,16	28,91	87,50	AT5G05830.1 Symbols: RING/FYVE/PHD zinc finger superfamily protein chr5:1755910-1756825 FORWARD LENGTH=204
Eucgr.C00760.1	6,07	0,20	0,60	1,33	7,75	AT5G20860.1 Symbols: Plant invertase/pectin methylesterase inhibitor superfamily
Eucgr.K01270.1	13,04	0,28	0,23	0,37	15,01	AT2G40435.1 Symbols: unknown protein
Eucgr.I02212.1	0,39	4,15	7,02	12,24	0,39	AT2G17480.1 Symbols: MLO8, ATMLO8 Seven transmembrane MLO family protein chr2:7590559-7593768 REVERSE LENGTH=593
Eucgr.K02666.1	11,07	56,40	52,31	43,86	10,44	AT5G50860.1 Symbols: Protein kinase superfamily protein chr5:20693778-20696983 REVERSE LENGTH=580
Eucgr.B03599.2	17,72	3,00	3,77	4,78	16,45	No hits found
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.J00341.1	27,89	0,69	3,03	5,81	25,77	No hits found
Eucgr.H01338.1	10,78	0,18	0,38	0,41	13,22	AT3G05910.1 Symbols: Pectinacylesterase family protein chr3:1764509-1767240 REVERSE LENGTH=415
Eucgr.G01458.1	14,20	0,30	0,49	0,58	13,53	AT5G42700.1 Symbols: AP2/B3-like transcriptional factor family protein
Eucgr.I00788.1	4,85	0,51	0,76	1,50	4,57	AT5G67280.1 Symbols: RLK receptor-like kinase chr5:26842430-26845126 REVERSE LENGTH=751
Eucgr.B03185.1	13,19	0,54	0,83	2,82	13,56	AT2G03350.1 Symbols: Protein of unknown function, DUF538 chr2:1019733-1021071 REVERSE LENGTH=179
Eucgr.G01214.1	2,95	0,21	0,45	0,67	2,72	AT4G33820.1 Symbols: Glycosyl hydrolase superfamily protein chr4:16217010-16219515 REVERSE LENGTH=570

Eucgr.K02352.1	2,14	0,00	0,25	0,41	1,96	AT1G05680.1 Symbols: UGT74E2 Uridine diphosphate glycosyltransferase 74E2
Eucgr.H00801.5	7,59	38,25	42,75	37,11	8,29	AT2G32700.6 Symbols: LUH LEUNIG_homolog chr2:13867235-13871844 FORWARD LENGTH=785
Eucgr.K02353.1	2,12	0,05	0,15	0,36	2,06	AT1G05680.1 Symbols: UGT74E2 Uridine diphosphate glycosyltransferase 74E2
Eucgr.B02381.1	10,55	0,00	0,33	0,85	10,43	No hits found
Eucgr.B03602.2	193,14	7,72	16,05	23,58	137,13	AT3G13510.1 Symbols: Protein of Unknown Function (DUF239) chr3:4403776-4405741 FORWARD LENGTH=419
Eucgr.K00961.1	1,99	25,38	20,89	15,29	1,94	AT2G20880.1 Symbols: Integrase-type DNA-binding superfamily protein chr2:8986053-8987063 FORWARD LENGTH=336
Eucgr.I01833.1	1,88	0,00	0,13	0,14	1,77	AT3G28180.1 Symbols: ATCSLC04, CSLC04, ATCSLC4, CSLC4 Cellulose-synthase-like C4
Eucgr.E01689.1	1,55	0,04	0,05	0,11	1,70	AT5G08640.2 Symbols: FLS1 flavonol synthase 1 chr5:2804009-2805175 FORWARD LENGTH=336
Eucgr.H02604.1	91,88	8,08	9,71	9,37	75,19	AT2G30520.1 Symbols: RPT2 Phototropic-responsive NPH3 family protein
Eucgr.H02774.1	0,43	0,02	0,16	0,01	0,41	AT3G07130.1 Symbols: ATPAP15, PAP15 purple acid phosphatase 15 chr3:2255763-2257981 REVERSE LENGTH=532
Eucgr.A00492.1	10,69	0,31	1,97	3,86	10,28	AT1G19330.2 Symbols: unknown protein
Eucgr.J01875.1	11,96	0,36	0,78	1,55	11,50	AT2G30140.2 Symbols: UDP-Glycosyltransferase superfamily protein chr2:12872200-12873691 FORWARD LENGTH=454
Eucgr.I01733.1	14,44	84,36	69,78	70,48	14,06	AT3G27540.1 Symbols: beta-1,4-N-acetylglucosaminyltransferase family protein
Eucgr.H05065.1	0,32	0,04	0,02	0,01	0,34	AT2G38290.1 Symbols: ATAMT2, AMT2;1, AMT2 ammonium transporter 2
Eucgr.A02126.1	1369,19	134,99	259,54	284,25	467,15	AT5G17920.2 Symbols: ATCIMS Cobalamin-independent synthase family protein
Eucgr.J00502.1	28,27	4,09	5,06	7,33	27,45	AT3G44380.1 Symbols: Late embryogenesis abundant (LEA) hydroxyproline-rich glycoprotein family
Eucgr.B03026.1	1,02	4,48	4,62	3,74	1,32	No hits found
Eucgr.C02733.1	12,63	67,36	80,38	96,78	15,27	AT5G25280.2 Symbols: serine-rich protein-related chr5:8773882-8774544 FORWARD LENGTH=220
Eucgr.C03707.1	6,11	1,45	1,73	2,28	5,86	AT4G29310.1 Symbols: Protein of unknown function (DUF1005) chr4:14437892-14439609 REVERSE LENGTH=424
Eucgr.F03859.2	33,36	101,32	71,97	78,01	40,09	AT4G29850.1 Symbols: Eukaryotic protein of unknown function (DUF872)
Eucgr.D00477.1	3,33	0,37	0,12	0,22	4,20	AT4G18740.1 Symbols: Rho termination factor chr4:10303543-10304479 REVERSE LENGTH=245
Eucgr.B03753.1	12,70	1,38	2,68	6,03	12,37	No hits found
Eucgr.I01191.1	5,14	1,11	0,77	0,65	6,35	AT3G19000.1 Symbols: 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein
Eucgr.D02033.1	50,03	7,66	9,50	11,64	51,25	AT4G22570.1 Symbols: APT3 adenine phosphoribosyl transferase 3 chr4:11882310-11885250 REVERSE LENGTH=183
Eucgr.G00854.1	8,35	31,91	33,46	32,78	10,10	AT4G33510.1 Symbols: DHS2 3-deoxy-d-arabino-heptulosonate 7-phosphate synthase
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.A00534.1	14,19	54,90	54,03	46,35	17,02	AT2G38280.2 Symbols: FAC1, ATAMPD AMP deaminase, putative / myoadenylate deaminase, putative
Eucgr.B02014.1	16,39	1,97	4,29	6,06	16,18	AT1G54730.2 Symbols: Major facilitator superfamily protein chr1:20424471-20429978 FORWARD LENGTH=470
Eucgr.D01737.1	1,62	13,99	12,83	9,19	1,70	AT5G53110.1 Symbols: RING/U-box superfamily protein chr5:21529022-21533008 FORWARD LENGTH=382
Eucgr.H01397.1	2,80	0,16	0,23	0,14	2,81	AT4G12910.1 Symbols: scpl20 serine carboxypeptidase-like 20 chr4:7550576-7553051 REVERSE LENGTH=497

Eucgr.J02154.1	3,05	0,05	0,35	0,84	3,81	No hits found
Eucgr.B00830.1	2,26	9,06	10,46	12,27	2,71	AT1G73750.1 Symbols: Uncharacterised conserved protein UCP031088, alpha/beta hydrolase
Eucgr.F01551.1	25,89	4,48	6,22	7,93	27,99	AT1G75130.1 Symbols: CYP721A1 cytochrome P450, family 721, subfamily A, polypeptide 1
Eucgr.H00646.1	0,70	0,04	0,20	0,21	0,74	AT5G64740.1 Symbols: CESA6, IXR2, E112, PRC1 cellulose synthase 6 chr5:25881555-25886333 FORWARD LENGTH=1084
Eucgr.F02183.1	1716,98	105,00	291,10	447,72	519,92	AT4G14960.2 Symbols: TUA6 Tubulin/FtsZ family protein chr4:8548769-8550319 REVERSE LENGTH=450
Eucgr.K00551.3	9,90	2,75	1,74	0,95	10,13	AT5G66460.1 Symbols: MAN7, AtMAN7 Glycosyl hydrolase superfamily protein
Eucgr.E00821.1	3,07	0,38	0,35	0,78	3,03	AT4G24000.1 Symbols: ATCSLG2, CSLG2 cellulose synthase like G2 chr4:12462142-12465471 FORWARD LENGTH=722
Eucgr.B02486.1	2814,83	239,55	537,13	845,92	821,98	AT5G60490.1 Symbols: FLA12 FASCICLIN-like arabinogalactan-protein 12 chr5:24325916-24326665 REVERSE LENGTH=249
Eucgr.I01189.1	27,02	7,50	8,61	10,57	31,80	AT3G19000.1 Symbols: 2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase superfamily protein
Eucgr.F04235.1	28,30	2,07	3,77	5,89	28,69	AT1G21090.1 Symbols: Cupredoxin superfamily protein chr1:7384854-7386199 FORWARD LENGTH=242
Eucgr.A01371.1	10,65	28,80	34,26	35,50	11,87	AT3G07650.4 Symbols: COL9 CONSTANS-like 9 chr3:2442494-2443901 FORWARD LENGTH=372
Eucgr.A01444.1	6,26	0,45	1,14	1,70	6,81	AT5G24320.1 Symbols: Transducin/WD40 repeat-like superfamily protein chr5:8284858-8287651 REVERSE LENGTH=694
Eucgr.B03946.1	7,30	0,88	1,40	1,96	8,05	AT5G40240.2 Symbols: nodulin MtN21 /EamA-like transporter family protein
Eucgr.H00308.1	505,18	47,41	110,30	194,67	268,51	AT3G59690.1 Symbols: IQD13 IQ-domain 13 chr3:22049572-22051479 FORWARD LENGTH=517
Eucgr.D02037.1	10,34	42,33	30,56	18,36	12,06	AT1G12110.1 Symbols: NRT1.1, CHL1-1, NRT1, B-1, ATNRT1, CHL1 nitrate transporter 1.1
Eucgr.I02691.1	87,53	15,36	20,83	22,59	75,88	AT5G64550.1 Symbols: Iorocrin-related chr5:25801794-25803698 REVERSE LENGTH=634
Eucgr.C01809.1	29,77	9,97	10,66	15,35	31,65	AT4G32570.1 Symbols: TIFY8 TIFY domain protein 8 chr4:15716433-15718702 REVERSE LENGTH=361
Eucgr.J00751.1	12,23	1,25	2,83	4,05	13,42	AT3G21070.2 Symbols: NADK1 NAD kinase 1 chr3:7380434-7383874 REVERSE LENGTH=524
Eucgr.F01752.1	21,08	62,97	56,81	55,11	24,28	AT4G35230.1 Symbols: BSK1 BR-signaling kinase 1 chr4:16755325-16758041 REVERSE LENGTH=512
Eucgr.F03305.1	96,42	2,13	6,32	16,93	98,09	No hits found
Eucgr.H00896.1	76,24	2,72	10,01	15,98	69,72	AT2G47160.1 Symbols: BOR1 HCO3- transporter family chr2:19357740-19360787 REVERSE LENGTH=704
Eucgr.F00581.1	5,38	23,68	22,14	20,45	5,64	AT1G30755.1 Symbols: Protein of unknown function (DUF668) chr1:10905991-10908773 REVERSE LENGTH=615
Eucgr.J00385.2	66,16	2,03	4,51	7,69	68,84	AT5G22930.1 Symbols: Protein of unknown function (DUF1635) chr5:7668229-7669315 REVERSE LENGTH=238
Eucgr.I01383.1	5,06	0,54	2,27	2,71	5,31	AT4G34880.1 Symbols: Amidase family protein chr4:16615549-16617424 FORWARD LENGTH=466
Eucgr.C00546.1	8,12	0,82	1,83	1,98	8,69	AT5G11590.1 Symbols: TINY2 Integrase-type DNA-binding superfamily protein
Eucgr.J02264.2	8,00	20,31	19,28	21,47	8,49	No hits found
Eucgr.H00495.1	144,56	1,81	5,52	12,64	176,15	AT3G07880.1 Symbols: SCN1 Immunoglobulin E-set superfamily protein chr3:2514175-2515544 FORWARD LENGTH=240
GENE ID	FPKM					Annotation
	CT	CH	NAR	CHSTOP	NARSTOP	TAIR
Eucgr.G03246.4	100,42	12,92	15,05	15,54	141,43	AT3G54500.1 Symbols: unknown protein
Eucgr.J01177.3	8,06	1,76	2,79	3,96	8,69	AT5G13500.3 Symbols: unknown protein;

Eucgr.H05081.4	4,84	0,93	1,11	1,60	5,34	AT4G34240.1 Symbols: ALDH3I1, ALDH3 aldehyde dehydrogenase 3I1 chr4:16389801-16392633 FORWARD LENGTH=550
Eucgr.H02297.1	8,31	32,34	24,25	19,28	8,92	No hits found
Eucgr.F02039.1	14,22	41,88	33,89	24,33	16,13	AT5G55860.1 Symbols: Plant protein of unknown function (DUF827) chr5:22610146-22612166 FORWARD LENGTH=649
Eucgr.D00506.1	257,68	0,26	2,84	6,70	239,70	AT5G45890.1 Symbols: SAG12 senescence-associated gene 12 chr5:18613300-18614759 FORWARD LENGTH=346
Eucgr.H04094.1	89,16	1,50	9,20	14,90	104,05	AT3G62270.1 Symbols: HCO3- transporter family chr3:23042528-23045633 REVERSE LENGTH=703
Eucgr.H02634.1	386,73	21,78	47,14	72,69	396,09	AT3G23730.1 Symbols: XTH16 xyloglucan endotransglucosylase/hydrolase 16
Eucgr.B00824.1	2280,86	132,19	233,56	407,73	1834,47	AT3G18280.1 Symbols: Bifunctional inhibitor/lipid-transfer protein/seed storage 2S albumin superfamily protein