



UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<https://jnnp.bmj.com/content/83/10/970>

DOI: 10.1136/jnnp-2012-302770

Direitos autorais / Publisher's copyright statement:

©2012 by BMJ. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

Published in final edited form as:

J Neurol Neurosurg Psychiatry. 2012 October ; 83(10): 970–974. doi:10.1136/jnnp-2012-302770.

Olfactory Impairment in Familial Ataxias

Mariana Moscovich, MD^{2,3,6}, Renato Puppi Munhoz, MD, MSc^{2,6}, Helio A Teive, MD², Salmo Raskin, MD^{2,7}, Margaret J Carvalho, MD⁴, Egberto R Barbosa, MD, PhD⁴, Ronald Ranvaud, PhD^{4,5}, Jilin Liu, Ms³, Karen McFarland, PhD³, Tetsuo Ashizawa, MD³, Andrew J Lees, MD¹, and Laura Silveira-Moriyama, MD, PhD^{1,8,*}

¹Reta Lila Weston Institute of Neurological Studies, UCL Institute of Neurology, London, UK

²Department of Neurology, Hospital de Clínicas, Federal University of Paraná, Curitiba-PR, Brazil

³Department of Neurology, University of Florida, Center for Movement Disorders & Neurorestoration, McKnight Brain Institute, Gainesville, FL, USA

⁴Department of Neurology, University of Sao Paulo School of Medicine, Sao Paulo, Brazil

⁵Department of Physiology and Biophysics, Institute of Biomedical Sciences, University of Sao Paulo, Brazil

⁶Department of Neurology, Pontifical Catholic University of Parana, Curitiba, Parana, Brazil

⁷Core for Advanced Molecular Investigation, Graduate Program in Health Sciences, Center for Biological and Health Sciences, Pontifical Catholic University of Paraná Curitiba, Parana, Brazil

⁸Department of Neurology, University of Campinas, UNICAMP, Brazil

Abstract

The main clinical manifestations of the spinocerebellar ataxias (SCAs) result from the involvement of the cerebellum and its connections. Cerebellar activity has been consistently observed in functional imaging studies of olfaction, but the anatomical pathways responsible for this connection have not yet been elucidated. Previous studies have demonstrated olfactory deficit

Corresponding Author: Laura Silveira-Moriyama, Reta Lila Weston Institute of Neurological Studies, UCL Institute of Neurology, 1 Wakefield St, London, WC1N 1PJ, United Kingdom. Phone: + 44 20 76794025, fax: + 44 20 7278 4993, l.moriyama@ion.ucl.ac.uk.

COMPETING INTERESTS, FUNDING

Dr. Moscovich reports no disclosures. Dr. Munhoz reports no disclosures.

Dr. Ashizawa is funded by NINDS grant RC1NS068897, National Ataxia Foundation, Marigold Foundation. Dr. Teive received personal compensation for educational activities with Allergan, Boehringer, Ingelheim, Ipsen, Novartis, and Roche and serves as an editorial board member of Parkinsonism and Related Disorders, Journal of Neurology Research, Parkinson's Disease, Current Neurology and Neuroscience Reports, and Arquivos de Neuropsiquiatria. Dr. Carvalho reports no disclosures. Dr. Raskin reports no disclosure. Liu reports no disclosure. Dr. Lees is funded by PSP Association, Weston Trust - The Reta Lila Howard Foundation grant, is a member consultant of Genus, advisory board of Novartis, Teva, Meda, Boehringer Ingelheim, GSK, Ipsen, Lundbeck, Allergan, Orion, BIAL, Noscira, Roche and received honoraria from Novartis, Teva, Meda, Boehringer Ingelheim, GSK, Ipsen, Lundbeck, Allergan, Orion, BIAL, Noscira, Roche. Dr. Barbosa received personal compensation for educational activities from Boehringer-Ingelheim. Dr. Ranvaud reports no disclosure. Dr. McFarland reports no disclosure.

COPYRIGHT LICENCE STATEMENT

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive license (or non-exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd, and its Licensees to permit this article (if accepted) to be published in the Journal of Neurology, Neurosurgery and Psychiatry and any other BMJPLG products and to exploit all subsidiary rights, as set out in our license.

AUTHORS' CONTRIBUTIONS

M Moscovich: conceptualized study, analyzed data in study, drafted manuscript, revised manuscript R P Munhoz: drafted manuscript, revised manuscript H A Teive: revised manuscript S Raskin: revised manuscript M J Carvalho: revised manuscript E R Barbosa: revised manuscript R Ranvaud: revised manuscript J Liu: revised manuscript K McFarland: revised manuscript T Ashizawa: revised manuscript A J Lees: revised manuscript L Silveira-Moriyama: designed study, interpreted data in study, analyzed data in study, drafted manuscript, revised manuscript.

in SCA2, Friedreich's ataxia (FA) and in small groups of ataxia of diverse etiology. We used a validated version of the 16 item smell identification test from Sniffin' Sticks (SS-16) was used to evaluate 37 patients with genetically determined autosomal dominant ataxia, and 31 with familial ataxia of unknown genetic basis. This data was also compared to results in 106 Parkinson's disease (PD) patients and 218 healthy controls. The SS-16 score was significantly lower in ataxia than in the control group ($p < 0.001$, 95% CI for $\beta = 0.55$ to 1.90) and significantly higher in ataxia than in PD ($p < 0.001$, 95% CI for $\beta = -4.58$ to -3.00) when adjusted for age ($p = 0.001$, 95% CI for $\beta = -0.05$ to -0.01), gender ($p = 0.19$) and history of tobacco use ($p = 0.41$). When adjusted for general cognitive function we found no significant difference between the ataxia and control group. Our study confirms previous findings of mild hyposmia in ataxia, and further suggests this may be due to general cognitive deficits rather than specific olfactory problems.

Search Terms

Movement disorders; Smell; Cerebellar ataxia; Cerebellar degeneration; Cognition

Introduction

Olfactory dysfunction is documented in a number of neurological disorders with marked neurodegenerative changes found in the brain at post-mortem, including Alzheimer's disease (AD), Parkinson's disease (PD), progressive supranuclear palsy, multiple system atrophy (MSA), corticobasal degeneration, and parkinsonism-dementia complex of Guam.[1–6] Ataxias result from the involvement of cerebellar structures and extra-cerebellar lesions, especially those due to brainstem involvement.[7] Spinocerebellar ataxias (SCA) are a genetically and clinically heterogeneous group of autosomal dominantly inherited progressive ataxia disorders. Up to now, almost 31 different gene *loci* have been found.[8–10]

A few reports have investigated olfactory function in cerebellar ataxias, showing a mild but significant olfactory impairment in patients with varied forms of ataxias.[11–14] Functional magnetic resonance imaging (MRI) studies show odor-induced cerebellar activity that is independent of sniffing.[15] The functional role of the cerebellum in olfaction and pathways through which olfactory information reach the cerebellum are unknown.[16]

We aimed at investigating if smell was impaired in a large sample of patients with familial ataxias, including fourteen patients with SCA10, a condition in which smell testing has never been described before.

Methods

All subjects were recruited in Brazil between 2004 and 2011. Subjects were recruited at the University of Sao Paulo and the Federal University of Parana (between 2009 and 2011). The protocol used to test the sense of smell and evaluate cognitive function was identical and the researchers are part of a collaborative network.

Patients with a diagnosis of hereditary episodic ataxia and X linked ataxias, participants with an active upper respiratory tract infection, MMSE scores below 18 and those with a previous history of head trauma were excluded. Consent was obtained from all participants and the protocol was approved by the local ethics committees.

Subjects

Smell testing was performed in sixty-eight patients including those with SCA diagnoses confirmed by genetic test (one SCA 1, two SCA 2, seventeen SCA 3, one SCA 6, one SCA 7, fourteen SCA 10 and one SCA 13) and also thirty-one familial ataxia patients without genetic confirmation.

Genetically confirmed SCA

Of the 37 patients with confirmed SCA diagnoses, seventeen (45.9%) were female and seventeen (45.9%) were smokers. Mean age was 48.4 years [standard deviation (SD) 11 years], mean age of onset was 35.3 years (SD 10.7 years), and mean disease duration was 13 years (SD 8.5 years). In the SCA subgroup, the mean disease duration was 11 years (SD 5 years) on the SCA 3 group and 17.6 years (SD 10.6 years) for the SCA 10 group. The mean SARA score for the SCA 3 group was 14.4 (SD 6.2) compared with 10.68 (SD 5.7) on the SCA 10 group. Genomic DNA was isolated from peripheral blood using standard protocols and mutations were screened and confirmed using previously established methods.[17–19]

Familial ataxia without genetic diagnosis

A total of thirty-one patients with familial ataxia of unknown etiology were tested. Fifteen (48.5%) were female and nine (27.3%) were smokers. All patients had a clinical diagnosis of familial ataxia, progressive ataxia with gait and stance impairment, speech disturbance and oculomotor abnormalities with no apparent medical causes such as vitamin deficiencies, infections, or exposure to toxins. Genetic testing was performed in 20 patients but none of the tested mutations was identified, and in others genetic testing was not performed due to socioeconomic reasons.

Mean age was 45 years (SD 10.9 years), mean age of onset was 34 years (SD 13 years), and mean disease duration was 10.9 years (SD 8.8 years).

Control Groups

For the comparison groups we used data from 106 PD patients and 218 control subjects tested for a previous study.[4] In the PD group, the mean age was 61.2 years (SD 11.0 years), 35 (33%) of subjects were female and 41 (38%) were smokers. Mean age of onset of the PD patients was 48.9 years (SD 13.4 years) and mean disease duration was 12.3 years (SD 9.0 years). In the control group, mean age was 50.9 years (SD 17.1 years), 92 (42.2%) were female and 103 (47.2%) were smokers. (Table 1)

Clinical studies

Smell testing

A previously validated Brazilian-Portuguese translation of the 16 item smell identification test from Sniffin' Sticks (SS-16) [20] was used. Clinical assessment was conducted using the Scale for the Assessment and Rates of Ataxia (SARA) [21, 22] (scores ranging from 3 to 30) and a validated translation of the Mini-Mental Status Examination (MMSE) [23]

Statistical Analyses

In the ataxia group the association between the score in the SS-16 and other clinical variables (disease duration, SARA, MMSE) was done using multiple linear regressions (MLR) adjusting for age, gender and smoking as covariates. To investigate the difference in the average SS-16 score between groups when adjusting for other possible influencing factors such as age, gender, history of smoking and score in the MMSE, MLR analyses were used. When the three groups of patients were compared, indicator variables were used to

compare the ataxia group with control and PD groups. Because there was a significant effect of the MMSE on the SS-16 scores and a significant interaction between MMSE and the variable comparing ataxia and PD subjects, subsequent analyses were performed to investigate the relationship between the SS-16 and MMSE in the three groups of subjects using MLR and also partial correlations. Partial correlation coefficients between MMSE and SS-16 when adjusting for age and gender were compared between the groups ataxia and controls using the Z distribution using the online software Stattools (<http://www.stattools.net/>). Except where otherwise stated, all analyses were performed using the statistics software SPSS v19. Assumptions for the MLR analyses were checked by visual inspection of the residuals.

Results

In the ataxia group (n=67), the SS-16 score was significantly associated with the MMSE score ($p=0.001$, 95% CI for $\beta = 0.18$ to 0.63) when adjusted for age ($p=0.95$), gender ($p=0.03$, 95% CI for $\beta = -2.55$ to -0.15), smoking ($p=0.11$), SARA score ($p=0.37$), and disease duration ($p=0.91$). The variance inflation factor (VIF) for the covariates ranged between 1.11 to 1.38, indicating an absence of significant multicollinearity.

A multiple linear regression (MLR) including all 391 subjects demonstrated that the SS-16 score was significantly lower in the ataxia group than in the control group ($p=0.001$, 95% CI for $\beta = 0.52$ to 1.88) and significantly higher in the ataxia group than in PD group ($p<0.001$, 95% CI for $\beta = -4.60$ to -3.02) when adjusted for age ($p<0.001$, 95% CI for $\beta = -0.05$ to -0.01), gender ($p=0.19$) and history of tobacco use ($p=0.31$) (see figure 1); the VIF ranged between 1.08 and 2.91. When also adjusting for the MMSE ($p<0.001$, 95% CI for $\beta = 0.17$ to 0.37) in addition to age ($p=0.042$) and gender ($p=0.04$) the MLR showed the SS-16 score in the ataxia group to be higher than in the PD group ($p<0.001$, 95% CI for $\beta = -5.45$ to -3.80) and not different from that of controls ($p=0.31$); the VIF ranged between 1.08 and 2.41).

The scatter-plot between MMSE and SS-16 in the three groups and subtypes SCA3 and SCA10 (see figure 2 and supplemental figure 1) shows a moderate correlation between the SS-16 and the MMSE. The partial correlation coefficient between MMSE and SS-16 when adjusting for age, gender and smoking was $r=0.514$ [$p<0.001$, degrees of freedom (df) = 62] in the ataxia group, $r= 0.292$ ($p<0.001$, df =213) in the control group, and $r=0.028$ ($p=0.78$) in the PD group, indicating that variations in the MMSE explain roughly 26% of the variation of the SS-16 in the ataxia group, 9% of the variation in the control group, and not a significant amount of the SS-16 variation in the PD group. Using the Z distribution there was a significant difference between the partial correlation coefficient in the ataxia and control groups ($p=0.03$, $Z=1.8404$) indicating that the variation in the MMSE explains a larger amount of the variation of the SS-16 in the ataxia than in the control group.

Although it is a possibility that olfaction only develops in later stages of disease, our current study fails to demonstrate such association, because there was no association between the SS-16 and disease duration (Pearson's correlation p value = 0.11). Regarding disease severity, there was an association between SS-16 and SARA scores (Pearson's correlation -0.38 , $p=0.005$) showing that more affected subjects have worse olfactory performance, but when adjusting for MMSE and age - and other factors that could potentially affect the olfactory abilities - there was no independent association between the SS-16 and SARA ($p=0.37$), although the MMSE remained an independent predictor ($p=0.001$).

Discussion

Our study provides independent confirmation of the smell deficit found in a large number of heterogeneous ataxia patients, as none of the patients in the current study were included in any of the previous reports. These findings add to a large picture of olfactory deficits in different neurodegenerative diseases. [2, 8–12] (Listed in Table 2)

The main clinical manifestations of spinocerebellar ataxias result from the involvement of cerebellar structures but they often present with extra-cerebellar features.[7, 8] The decrement of olfactory function observed in previous studies was always small and far less marked than that reported in PD or AD, for instance.[2, 11, 12, 24]

Abele et al [12] demonstrated a moderate impairment of olfaction in 8 patients with MSA and 11 patients with sporadic ataxia with unknown etiology, although, when controlled for age the authors did not adjust the results for MMSE scores. Connely et al [11] studied a group of 35 non demented patients with assorted degenerative ataxias (including SCA 3, 7, 2 and FA), showing a deficit in olfaction when compared to age and gender matched control subjects. Fernandez-Ruiz et al [13] reported olfactory impairments in 29 patients with autosomal dominant, recessive and sporadic ataxia in comparison to 29 age and gender matched controls, however, they did not correlate the UPSIT with the MMSE scores. Velazquez-Perez et al [25] found that UPSIT scores were lower in a group of 53 SCA 2 patients when compared to 53 controls but when demented subjects were excluded, there was no significant difference in UPSIT scores between groups. Recently, Braga-Neto et al [14] demonstrated olfactory dysfunction in 41 subjects with SCA 3 even when matched for age, gender and MMSE scores. They excluded all patients whose MMSE score was below 24 or individuals with less than 5 years of education.

The role of the cerebellum in olfaction has been proposed but never fully clarified and the anatomical pathways that lead to this connection have not yet been demonstrated, however, in 1997 Yousen et al [15] reported functional MRI findings in five adult man with normal sense of smell, showing that olfactory nerve mediated stimulation activated the orbitofrontal and cerebellar areas, suggesting that the cerebellum is involved in sensory discrimination and attention to tasks. Sobel et al [26] using a functional magnetic resonance imaging (fMRI) demonstrated that the cerebellum shows olfactory-related activation which was dependent on the odorant concentration, reinforcing the idea of a role for the cerebellum in olfaction. Nevertheless, the same study showed cerebellar activation during the sniffing of nonodorized air, which was hypothesized as part of a cerebellar role in the maintenance of a feedback mechanism regulating the sniffing magnitude. In another functional study, Qureshy et al [27] mapped the human brain during olfactory processing and reported a cerebellar activation during olfactory naming, suggesting the cerebellum may have a role in cognitive olfactory processing. Savic reported [28–31] in 4 different papers using fMRI or PET scan that the cerebellum is activated during odor discrimination, odor recognition memory and during passive smelling. Finally, Ferdon and Murphy [32] demonstrated in an fMRI study with ten young and ten elderly adults, that olfactory tasks caused activation of the cerebellar lobes, especially in superior semilunar lobule, inferior semilunar lobule and posterior quadrangular lobule.

Besides the motor control and olfaction, the cerebellum has been well recognized to be involved in cognitive processing [33–36] These findings were consistent with a number of studies evidencing a cerebellar role during attention[37], memory [33, 38], language[39], and verbal fluency. [40]

Stoodley and Schmahmann [41] showed similar findings in a fMRI study with healthy subjects, showing cerebellar activation during cognitive functioning in the posterior lobe

areas, these findings were supported in different studies[35, 41, 42] and the posterior lobes were recognized as cognition regions.

Burk [33] studied cognition in SCA and FA patients reporting mild cognitive deficits in SCA1, SCA2 and SCA3 present on verbal memory tasks, supported by the hypothesis that these deficits are due to a disruption of cerebrotocerebellar circuitries and not from the cerebellar degeneration per se.

The relationship between cognitive and smell test scores has previously been reported in control subjects [43], subjects with mild cognitive impairment and in patients with Alzheimer's patients, and in other neurodegenerative diseases such as PD[44] and Progressive Supranuclear Palsy [6]. Ours is the first study to show an association between olfaction and cognitive function in ataxias, but we suggest in future studies cognitive performance - measured by the crude score on the MMSE or more sophisticated neuropsychometric testing - should be taken into consideration when studying olfaction in neurodegenerative diseases, and ataxia in particular.

Quantitative correlation of clinical findings with MRI data was not possible in this study, future studies including voxel-based morphometry, diffusion tensor imaging, and functional MRI might help clarify this subject further. In studies with homogeneous groups of genetic ataxia patients, correlation with repeat length might also provide additional insights.

Our data, derived from the largest series of patients with a heterogeneous group of ataxia tested for smell deficit so far, confirm the presence of hyposmia in cerebellar ataxia, and further demonstrates that cognitive deficits in the ataxia group might be a confounding factor, and should be taken into consideration in future studies. Although the two deficits might be independent, it is also possible that cognitive difficulties partly explain the smell deficit found in the ataxia group when compared to controls. .

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was funded by the Reta Lila Weston Trust for Medical Research. Dr. Laura Silveira-Moriyama is beneficiary of a Reta Lila Weston Fellowship, receives a grant from Parkinson's UK and travel grant from UCB and Teva.

References

1. Ahlskog JE, Waring SC, Petersen RC, et al. Olfactory dysfunction in Guamanian ALS, parkinsonism, and dementia. *Neurology*. 1998 Dec; 51(6):1672–1677. [PubMed: 9855521]
2. Meshulam RI, Moberg PJ, Mahr RN, et al. Olfaction in neurodegenerative disease: a meta-analysis of olfactory functioning in Alzheimer's and Parkinson's diseases. *Arch Neurol*. 1998 Jan; 55(1):84–90. [PubMed: 9443714]
3. Silveira-Moriyama L, Schwingenschuh P, O'Donnell A, et al. Olfaction in patients with suspected parkinsonism and scans without evidence of dopaminergic deficit (SWEDDs). *J Neurol Neurosurg Psychiatry*. 2009 Jul; 80(7):744–748. [PubMed: 19276101]
4. Silveira-Moriyama L, Carvalho Mde J, Katzenschlager R, et al. The use of smell identification tests in the diagnosis of Parkinson's disease in Brazil. *Mov Disord*. 2008 Dec 15; 23(16):2328–2334. [PubMed: 18785265]
5. Silveira-Moriyama L, Guedes LC, Kingsbury A, et al. Hyposmia in G2019S LRRK2-related parkinsonism: clinical and pathologic data. *Neurology*. 2008 Sep 23; 71(13):1021–1026. [PubMed: 18809839]

6. Silveira-Moriyama L, Hughes G, Church A, et al. Hyposmia in progressive supranuclear palsy. *Mov Disord*. 2010 Apr 15; 25(5):570–577. [PubMed: 20209627]
7. Manto M, Marmolino D. Cerebellar ataxias. *Curr Opin Neurol*. 2009 Aug; 22(4):419–429. [PubMed: 19421057]
8. Schols L, Bauer P, Schmidt T, et al. Autosomal dominant cerebellar ataxias: clinical features, genetics, and pathogenesis. *Lancet Neurol*. 2004 May; 3(5):291–304. [PubMed: 15099544]
9. Klockgether T, Paulson H. Milestones in ataxia. *Mov Disord*. 2011 May; 26(6):1134–1141. [PubMed: 21626557]
10. Klockgether T. Update on degenerative ataxias. *Curr Opin Neurol*. 2011 Aug; 24(4):339–345. [PubMed: 21734495]
11. Connelly T, Farmer JM, Lynch DR, et al. Olfactory dysfunction in degenerative ataxias. *J Neurol Neurosurg Psychiatry*. 2003 Oct; 74(10):1435–1437. [PubMed: 14570842]
12. Abele M, Riet A, Hummel T, et al. Olfactory dysfunction in cerebellar ataxia and multiple system atrophy. *J Neurol*. 2003 Dec; 250(12):1453–1455. [PubMed: 14673578]
13. Fernandez-Ruiz J, Diaz R, Hall-Haro C, et al. Olfactory dysfunction in hereditary ataxia and basal ganglia disorders. *Neuroreport*. 2003 Jul 18; 14(10):1339–1341. [PubMed: 12876469]
14. Braga-Neto P, Felicio AC, Pedroso JL, et al. Clinical correlates of olfactory dysfunction in spinocerebellar ataxia type 3. *Parkinsonism Relat Disord*. 2011 Jun; 17(5):353–356. [PubMed: 21367642]
15. Yousem DM, Williams SC, Howard RO, et al. Functional MR imaging during odor stimulation: preliminary data. *Radiology*. 1997 Sep; 204(3):833–838. [PubMed: 9280268]
16. Mainland JD, Johnson BN, Khan R, et al. Olfactory impairments in patients with unilateral cerebellar lesions are selective to inputs from the contralesional nostril. *J Neurosci*. 2005 Jul 6; 25(27):6362–6371. [PubMed: 16000626]
17. Teive HA, Munhoz RP, Raskin S, et al. Spinocerebellar ataxia type 10: Frequency of epilepsy in a large sample of Brazilian patients. *Mov Disord*. 2010 Dec 15; 25(16):2875–2878. [PubMed: 20818609]
18. Teive HA, Munhoz RP, Arruda WO, et al. Spinocerebellar ataxia type 10 - A review. *Parkinsonism Relat Disord*. 2011 Nov; 17(9):655–661. [PubMed: 21531163]
19. Teive HA, Munhoz RP, Raskin S, et al. Spinocerebellar ataxia type 6 in Brazil. *Arq Neuropsiquiatr*. 2008 Sep; 66(3B):691–694. [PubMed: 18949263]
20. Kobal G, Hummel T, Sekinger B, et al. "Sniffin' sticks": screening of olfactory performance. *Rhinology*. 1996 Dec; 34(4):222–226. [PubMed: 9050101]
21. Subramony SH. SARA--a new clinical scale for the assessment and rating of ataxia. *Nat Clin Pract Neurol*. 2007 Mar; 3(3):136–137. [PubMed: 17290240]
22. Schmitz-Hubsch T, du Montcel ST, Baliko L, et al. Scale for the assessment and rating of ataxia: development of a new clinical scale. *Neurology*. 2006 Jun 13; 66(11):1717–1720. [PubMed: 16769946]
23. Brucki SM, Nitri R, Caramelli P, et al. Suggestions for utilization of the mini-mental state examination in Brazil. *Arq Neuropsiquiatr*. 2003 Sep; 61(3B):777–781. [PubMed: 14595482]
24. Hawkes C. Olfaction in neurodegenerative disorder. *Mov Disord*. 2003 Apr; 18(4):364–372. [PubMed: 12671941]
25. Velazquez-Perez L, Fernandez-Ruiz J, Diaz R, et al. Spinocerebellar ataxia type 2 olfactory impairment shows a pattern similar to other major neurodegenerative diseases. *J Neurol*. 2006 Sep; 253(9):1165–1169. [PubMed: 16609806]
26. Sobel N, Prabhakaran V, Hartley CA, et al. Odorant-induced and sniff-induced activation in the cerebellum of the human. *J Neurosci*. 1998 Nov 1; 18(21):8990–9001. [PubMed: 9787004]
27. Qureshy A, Kawashima R, Imran MB, et al. Functional mapping of human brain in olfactory processing: a PET study. *J Neurophysiol*. 2000 Sep; 84(3):1656–1666. [PubMed: 10980035]
28. Savic I. Imaging of brain activation by odorants in humans. *Curr Opin Neurobiol*. 2002 Aug; 12(4):455–461. [PubMed: 12139995]
29. Savic I. Brain imaging studies of the functional organization of human olfaction. *Neuroscientist*. 2002 Jun; 8(3):204–211. [PubMed: 12061500]

30. Savic I, Gulyas B, Berglund H. Odorant differentiated pattern of cerebral activation: comparison of acetone and vanillin. *Hum Brain Mapp.* 2002 Sep; 17(1):17–27. [PubMed: 12203685]
31. Savic I. Processing of odorous signals in humans. *Brain Res Bull.* 2001 Feb; 54(3):307–312. [PubMed: 11287135]
32. Ferdon S, Murphy C. The cerebellum and olfaction in the aging brain: a functional magnetic resonance imaging study. *Neuroimage.* 2003 Sep; 20(1):12–21. [PubMed: 14527566]
33. Burk K. Cognition in hereditary ataxia. *Cerebellum.* 2007; 6(3):280–286. [PubMed: 17786824]
34. Molinari M, Chiricozzi FR, Clausi S, et al. Cerebellum and detection of sequences, from perception to cognition. *Cerebellum.* 2008; 7(4):611–615. [PubMed: 18941861]
35. Tedesco AM, Chiricozzi FR, Clausi S, et al. The cerebellar cognitive profile. *Brain.* 2011 Dec; 134(Pt 12):3669–3683.
36. Schmahmann JD, Sherman JC. The cerebellar cognitive affective syndrome. *Brain.* 1998 Apr; 121(Pt 4):561–579. [PubMed: 9577385]
37. Gottwald B, Mihajlovic Z, Wilde B, et al. Does the cerebellum contribute to specific aspects of attention? *Neuropsychologia.* 2003; 41(11):1452–1460. [PubMed: 12849763]
38. Chiricozzi FR, Clausi S, Molinari M, et al. Phonological short-term store impairment after cerebellar lesion: a single case study. *Neuropsychologia.* 2008; 46(7):1940–1953. [PubMed: 18342342]
39. Klein D, Milner B, Zatorre RJ, et al. The neural substrates underlying word generation: a bilingual functional-imaging study. *Proc Natl Acad Sci U S A.* 1995 Mar 28; 92(7):2899–2903. [PubMed: 7708745]
40. Peterburs J, Bellebaum C, Koch B, et al. Working memory and verbal fluency deficits following cerebellar lesions: relation to interindividual differences in patient variables. *Cerebellum.* 2010 Sep; 9(3):375–383. [PubMed: 20387024]
41. Stoodley CJ, Schmahmann JD. Functional topography in the human cerebellum: a meta-analysis of neuroimaging studies. *Neuroimage.* 2009 Jan 15; 44(2):489–501. [PubMed: 18835452]
42. Timmann D, Brandauer B, Hermsdorfer J, et al. Lesion-symptom mapping of the human cerebellum. *Cerebellum.* 2008; 7(4):602–606. [PubMed: 18949530]
43. Silveira-Moriyama L, Azevedo AM, Ranvaud R, Barbosa ER, Doty RL, Lees AJ. Applying a new version of the Brazilian-Portuguese UPSIT smell test in Brazil. *Arq Neuropsiquiatr.* 2010 Oct; 68(5):700–705. [PubMed: 21049178]
44. Morley JF, Weintraub D, Mamikonyan E, Moberg PJ, Siderowf AD, Duda JE. Olfactory dysfunction is associated with neuropsychiatric manifestations in Parkinson's disease. *Mov Disord.* 2011 Sep; 26(11):2051–2057. [PubMed: 21611985]




Figure 1.

Box plot of the Sniffin Sticks score in the three patient groups and subtypes SCA3 and SCA10. The median (the horizontal line) is within the box containing the central 50% of the observations and the error bar contains the central 95% of the ordered observations. PD, Parkinson's disease.




Figure 2. Scatter plot of Sniffin Sticks and MMSE scores in the subjects who underwent both tests. Fit line showing the association between MMSE and Sniffin Sticks in each group. MMSE, Mini Mental State Examination

Table 1

Summary of clinical variables

	SCA	Familial but not genetically confirmed ataxias	PD	CONTROL
N	37	31	106	218
Age in years	Mean \pm SD, range 48.4 \pm 11	45 \pm 10.9	61.2 \pm 11	50.9 \pm 17.1
Age of onset in years	Mean \pm SD, range 35.3 \pm 10.7	34 \pm 13	48.9 \pm 13.4	N/A
Disease duration in years	Mean \pm SD, range 13 \pm 8.5	10.9 \pm 8.8	12.3 \pm 9	N/A
Disease severity	Mean \pm SD, range SARA 12.3 \pm 5.5	SARA 12 \pm 7	UPDRS 25.5 \pm 14	N/A
History of smoking	N (%) positive 17 (45.9)	9 (27.3)	41 (38.7)	103 (47.2)
Gender	N (%) woman 17 (45.9)	15 (48.3)	35 (33)	92 (42.2)
MMSE	Mean \pm SD, range 25 \pm 3.2	24 \pm 4	27.1 \pm 2.5	27.5 \pm 2.1
SS-16	Mean \pm SD, range 11 \pm 2.6	10 \pm 4	6.5 \pm 2.7	11.9 \pm 2.3

SCA = Spinocerebellar ataxia SD = standard deviation; N = number of subjects; SS-16 = 16 item identification test from Sniffin Sticks (possible range 0 to 16). MMSE = Mini-Mental State Examination; PD = Parkinson's disease; SARA = Scale for the Assessment and Rating of Ataxia; UPDRS = Unified Parkinson's Disease Rating Scale

Table 2

Previous Studies

Paper	Number of cases	Test used	Method of analysis	Outcomes
Satya-Murti et al., 1988 ⁴³	7 FA 9 other neurological disorder	BAEP + non standard test	Studied olfactory function	FA patients were significantly lower than the normal controls.
Fernandez-Ruiz et al., 2003 ¹³	12 SCA2 5 SCA3 1 SCA 10 5 Sporadic 5 Recessive 1 FA 25 PD 27 HD	UPSIT	Compare UPSIT scores of the different groups.	Smell deficit in SCA 2, autosomal recessive ataxia and sporadic ataxia, but no in patients with SCA3
Abele et al., 2003 ¹²	8 MSA 1 late onset sporadic ataxia of unknown etiology	SS + (3 tests of olfactory function)	Studied olfactory function	No significant differences in olfactory function between patients with sporadic ataxia = MSA-C
Connely et al., 2003 ¹¹	2 SCA2 5 SCA3 1 SCA7 4 unidentified forms of cerebral degeneration) 23 FA	UPSIT	Study olfactory function in patients with cerebellar disorder.	FA group were significantly lower than controls, Group with ataxia was also lower than control
Mainland et al., 2005 ¹⁶	7 focal unilateral cerebellar lesions	UPSIT	Examined the olfactory function in patients with unilateral cerebellar lesions.	Patients with unilateral cerebellar lesions were impaired at olfactory identification.
Velazquez-Perez et al., 2006 ²⁵	53 SCA2	UPSIT + non standard test (detection threshold, discrimination threshold)	Analyzed olfactory threshold and their relation to other features.	Significant impairment in SCA 2:in olfactory threshold, quality, identification and discrimination
Braga Neto et al., 2011 ¹⁴	41 SCA 3 46 control	SS-16	Analyzed olfactory identification and correlate with MMSE and non-cerebellar symptoms	Significantly reduced in patients with SCA3 and that sex, MMSE scores and RLS also influence the SS-16 scores.

FA= Friedreich Ataxia; SCA= Spinocerebellar Ataxia; BAEP= Brainstem auditory evoked potentials; UPSIT= University of Pennsylvania Smell Test; MSA= Multiple system atrophy; MSA-C= Multiple system atrophy type cerebellar; SS= Sniffin'Sticks; RLS= restless leg syndrome; MMSE = Mini-Mental State Examination PD=Parkinson Disease HD=Huntington Disease