

UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<https://oaj.fupress.net/index.php/ah/article/view/1841>

DOI: 10.13128/Acta_Herpetol-23582

Direitos autorais / Publisher's copyright statement:

© by Firenze University Press. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

A lizard acting as carrier of the amphibian-killing chytrid *Batrachochytrium dendrobatidis* in southern Brazil

MARIANA R. PONTES^{1,2,*}, GUILHERME AUGUSTO-ALVES^{1,3}, CAROLINA LAMBERTINI^{1,3}, LUÍS FELIPE TOLEDO¹

¹ Laboratório de História Natural de Anfíbios Brasileiros (LaHNAB), Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, São Paulo, 13083-862, Brazil. *Corresponding author. E-mail: maah.retuci@gmail.com

² Programa de Pós-Graduação em Ecologia, Instituto de Biologia, Universidade Estadual de Campinas (Unicamp), Campinas, São Paulo, Brazil

³ Programa de Pós-Graduação em Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas (Unicamp), Campinas, São Paulo, Brazil

Submitted on: 2018, 3th March; revised on: 2018, 10th August; accepted on: 2018, 29th September
Editor: Dario Ottonello

Abstract. Fungal infections are causing widespread population declines and extinctions in all vertebrate classes. Among them, an important fungal disease chytridiomycosis, caused by the pathogenic chytrid *Batrachochytrium dendrobatidis* (*Bd*). With an aquatic infectious phase, *Bd* does not survive desiccation for long, but may be transported by non-amphibian carriers. Such mechanism is key to understand amphibian-chytrid dynamics and may contribute to local amphibian conservation action plans. Therefore, we surveyed *Bd* in reptiles from two different Brazilian rainforests, looking for possible *Bd* carriers. We sampled 35 individuals belonging to 11 squamate families, five from the Atlantic Forest and 30 from the Amazon. We detected *Bd* in one adult lizard, *Placosoma glabellum*. This lizard feeds, shelters, and breeds in the leaf-litter, and moves between Atlantic Forest streams. Hence, it may be carrying *Bd* from stream to stream, and also spreading the pathogen to direct-developing amphibians, which have no contact with water bodies and are more susceptible to chytridiomycosis than aquatic species. This is the first record of a non-amphibian chytrid carrier in South America. We suggest that additional field and museum samplings will contribute to understand whether *Bd* can actually infect reptiles, and how reptile carriers can affect chytrid dynamics in the wild.

Keywords. Amphibia, Reptilia, Squamata, Amazon, Atlantic Forest, Chytridiomycosis, Infection dynamics, Pathogen vector.

Infectious diseases, caused by a diversity of pathogenic agents, have become one of the most important threats to wildlife (Daszak et al., 2000; Daszak et al., 2013). For example, fungal infections are responsible for worldwide vertebrate population declines (Bleher et al., 2009; Turner et al., 2011; Fisher et al., 2012; Lorch et al., 2016), especially jeopardizing amphibians (Berger et al., 1998; Lips et al., 2006). Chytridiomycosis is an amphibian infectious disease caused by the water-borne fungus *Batrachochytrium dendrobatidis* (hereafter *Bd*) (Longcore et al., 1999). *Bd* infects epidermal tissues

of amphibian hosts, affecting essential physiological processes as osmoregulation (Berger et al., 2005; Van Rooij et al., 2015), which may cause sub-lethal effects (e.g., Bovo et al., 2016; Salla et al., 2015), or death of infected hosts through asystolic cardiac arrest, as a consequence of reductions on blood plasma sodium and potassium concentrations (Voyles, 2009). It has also been noticed that *Bd* produces toxic factors that inhibits lymphocyte proliferation on hosts (Fites et al., 2013).

Chytridiomycosis is associated with massive losses of amphibian populations all over the globe (Olson and

Ronnenberg, 2014; James et al., 2015; Carvalho et al., 2017). This pathogen has a biphasic life-cycle, with flagellated free-swimming zoospores, often discharged into water bodies as streams and ponds, and sessile zoosporangia with intracellular development (Rosenblum et al., 2010; Greenspan et al., 2012). *Bd* does not survive desiccation due to its aquatic life-cycle (Johnson et al., 2003), but may be carried by non-amphibian species (McMahon et al., 2013), which might contribute to pathogen transmission (Kilburn et al., 2011). Past studies have shown that non-amphibian organisms such as algae (Johnson and Speare, 2003), nematodes (Shapard et al., 2012), crayfishes (Brannelly et al., 2015), reptiles (Kilburn et al., 2011), and waterfowls (Garmyn et al., 2012) also transport *Bd* zoospores. In this sense, besides the studies with amphibian hosts it is essential to increase knowledge about other organisms that may be acting as carriers of this pathogen.

Thus, since there are no *Bd* records for reptiles in South America, here in order to find additional *Bd* potential carriers we swab-sampled reptiles from Brazil's Atlantic Forest and Amazon, both tropical rainforests, but different in terms of *Bd* occurrence and infection intensity (Carvalho et al., 2017; Lambertini et al., 2017). We swabbed reptiles collected from the states of Paraná, São Paulo and Espírito Santo in Brazil's Atlantic Forest, and from the state of Pará in the Brazilian Amazon. We sampled a total of 35 individuals of 11 Squamata families, five from the Atlantic Forest and 30 from the Amazon (Table 1).

We swabbed live individuals throughout their body, ventral and dorsal surfaces, limbs (for lizards), and head (as in Hyatt et al., 2007) to test for the presence of *Bd* zoospores. We then extracted DNA from each swab using Prep-Man Ultra® (Life Technologies) and proceeded with TaqMan® qPCR assay for *Bd* detection and quantification (Boyle et al., 2004; Lambertini et al., 2013). We considered positive results greater than or equal to one zoospore genomic equivalent (GE) (Kriger et al., 2007).

We detected one *Bd* positive individual, *Placosoma glabellum* (Peters, 1870) (currently housed at Museu de Zoologia “prof. Adão J. Cardoso” Universidade Estadual de Campinas: ZUEC-REP 3735), with a load of 3.14 *Bd* zoospore GE. To confirm the positive result, we rinsed this individual in running water for 10 minutes (following the protocols of Rodriguez et al., 2014 and Becker et al., 2016), swabbed the lizard again and run an additional qPCR experiment. In the second run we detected a load of 1.67 zoospore GE.

This is the first report of *Bd* in a reptile from South America. This pathogen has already been detected in lizards and snakes from Panama, also by qPCR detection and without histological analysis, which would con-

Table 1. Sampled species for *Batrachochytrium dendrobatidis* presence. Number of sampled individuals is in parentheses. Species in bold was positive for *Bd*. Taxonomy follows Costa and Bérnills (2018).

Species	Municipality, State	Biome
Squamata: “Lizards”		
Dactyloidae		
<i>Norops</i> sp.(1)	Belém, Pará	Amazon
Gekkonidae		
<i>Hemidactylus mabouia</i> (1)	Belém, Pará	Amazon
Gymnophthalmidae		
<i>Leposoma scincoides</i> (1)	Santa Teresa, Espírito Santo	Atlantic Forest
<i>Placosoma glabellum</i> (2)	Morretes, Paraná	Atlantic Forest
Phyllodactylidae		
<i>Thecadactylus rapicauda</i> (2)	Belém, Pará	Amazon
Polychrotidae		
<i>Polychrus</i> sp.(1)	Belém, Pará	Amazon
Mabuyidae		
<i>Copeoglossum nigropunctatum</i> (1)	Belém, Pará	Amazon
Sphaerodactylidae		
<i>Chatogekko amazonicus</i> (1)	Belém, Pará	Amazon
<i>Coleodactylus</i> sp. (1)	Belém, Pará	Amazon
<i>Gonatodes</i> sp.(1)	Belém, Pará	Amazon
<i>Lepidoblepharis</i> sp. (2)	Belém, Pará	Amazon
Tropiduridae		
<i>Plica umbra</i> (3)	Belém, Pará	Amazon
<i>Uranoscodon</i> sp. (3)	Belém, Pará	Amazon
Squamata: “Serpentes”		
Boidae		
<i>Boa constrictor</i> (1)	Belém, Pará	Amazon
Dipsadidae		
<i>Dipsas catesbyi</i> (1)	Belém, Pará	Amazon
<i>Erythrolamprus</i> sp. (1)	Belém, Pará	Amazon
<i>Helicops angulatus</i> (2)	Acará, Pará	Amazon
<i>Imantodes cenchoa</i> (2)	Belém, Pará	Amazon
<i>Imantodes lentiferus</i> (1)	Acará, Pará	Amazon
<i>Leptodeira annulata</i> (1)	Belém, Pará	Amazon
<i>Oxyrhopus clathratus</i> (1)	Peruíbe, São Paulo	Atlantic Forest
<i>Thamnodynastes</i> sp. (1)	Dores do Rio Preto, Espírito Santo	Atlantic Forest
Typhlopidae		
<i>Amerotyphlops reticulatus</i> (4)	Belém, Pará	Amazon

firm whether these reptiles are infected with *Bd* or only act as carriers (Kilburn et al., 2011). Anyway, despite the absence of histological analyses, it is clear that reptiles, including arboreal snakes and semiaquatic or terrestrial lizards, carry *Bd*, contributing to zoospore dispersion

across different habitats (Kilburn et al., 2011; present study).

The *Bd*-positive lizard was collected on 9 January 2013, at the Estrada da Graciosa road (PR-410), in the municipality of Morretes, state of Paraná. This site is within the same area where three divergent *Bd* lineages were isolated: *Bd*-Brazil/Asia-2, *Bd*-GPL and a hybrid between them (Schloegel et al., 2012; Jenkinson et al., 2016; O'Hanlon et al., 2018). Besides that, this individual lizard was collected on top of a rock inside a stream, syntopic to an endemic rheophilic frog species, *Cycloramphus rhyakonastes* (Anura: Cycloramphidae) (Nunes-de-Almeida et al., 2016). This is the only known population of this species (Silvano, 2004), and given that *Bd* infected individuals were already recorded [with a prevalence of 55% (10 out of 18 sampled), L.F.T., unpublished data], continuous exchange of *Bd* in this area can facilitate hybridization of the pathogen (Jenkinson et al., 2016), producing some strains that could potentially be hypervirulent (Greenspan et al., 2018), posing a real short-term threat to the endemic *C. rhyakonastes*. Furthermore, *Placosoma glabellum* is a terrestrial small-size lizard with distribution restricted to south and southeast of Brazil (Uetz, 2017). Studies have shown that *Bd* can survive up to seven weeks out of amphibian hosts, specifically in lake water (Johnson and Speare, 2003). Also, it can survive up to three hours on dry bird feathers (Johnson and Speare, 2005). This ability to survive may lead to a greater range of *Bd* zoospore dissemination by other organisms. In this specific case, the *Bd*-positive lizard species could carry *Bd* zoospores from the stream into the forest, potentially transmitting this pathogen to other water bodies, or to direct-developing anurans that lives in leaf-litter inside forested areas, and can be less tolerant to chytridiomycosis than indirect-developing aquatic anurans (Mesquita et al., 2017).

Given that reptiles are more vagile than amphibians, and less restricted to water bodies, they may play an important role in *Bd* dissemination in the wild. In this sense, we highlight the need for more studies attempting to understand how reptiles and other non-amphibian species may contribute to *Bd* transmission dynamics in wildlife.

ACKNOWLEDGMENTS

We thank Alexandre F. R. Missassi, Anat Belasen, Anyelet Valencia-Aguilar, Carlos Henrique L. Nunes-de-Almeida, Clarisse Betancourt-Roman, David Rodriguez, Kelly R. Zamudio, Timothy Y. James, and Thomas S. Jenkinson for field assistance. This work was funded by São Paulo Research Foundation (FAPESP #2016/25358-3) and

National Council for Scientific and Technological Development (CNPq #300896/2016-6). MRP and CL thank the Coordination for the Improvement of Higher Education Personnel (CAPES 001) for fellowships. Sampling permit was approved by Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio 17242-3) and is in accordance with Sistema Nacional de Gestão do Patrimônio Genético e do Conhecimento Tradicional Associado (Sis-Gen A60B4D9).

REFERENCES

- Becker, C.G., Rodriguez, D., Lambertini, C., Toledo, L.F., Haddad, C.F.B. (2016): Historical dynamics of *Batrachochytrium dendrobatidis* in Amazonia. *Ecography* **39**: 954-960.
- Berger, L., Speare, R., Daszak, P., Green, D.E., Cunningham, A.A., Goggin, C.L., Slocombe, R., Ragan, M.A., Hyatt, A.D., McDonald, K.R., Hines, H.B., Lips, K.R., Marantelli, G., Parkes, H. (1998): Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *P. Natl. Acad. Sci. USA*. **95**: 9031-9036.
- Berger, L., Speare, R., Skerratt, L.F. (2005): Distribution of *Batrachochytrium dendrobatidis* and pathology in the skin of green tree frogs *Litoria caerulea* with severe chytridiomycosis. *Dis. Aquat. Organ.* **68**: 65-70.
- Bleher, B.S., Hicks, A.C., Behr, M., Meteyer, C.U., Berlowski-Zier, B.M., Buckles, E.L., Coleman, J.T.H., Darling, S.R., Gargas, A., Niver, R., Okoniewski, J.C., Rudd, R.J., Stone, W.B. (2009): Bat white-nose syndrome: an emerging fungal pathogen? *Science* **323**: 227.
- Bovo, R., Andrade, D.V., Toledo, L.F., Longo, A.V., Rodriguez, D., Haddad, C.F.B., Zamudio, K.R., Becker, C.G. (2016): Physiological responses of Brazilian amphibians to an enzootic infection of the chytrid fungus *Batrachochytrium dendrobatidis*. *Dis. Aquat. Organ.* **11**: 245-252.
- Boyle, D.G., Boyle, D.B., Olsen, V., Morgan, J.A.T., Hyatt, A.D. (2004): Rapid quantitative detection of chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples using real-time Taqman PCR Assay. *Dis. Aquat. Organ.* **60**: 141-148.
- Brannelly, L.A., McMahon, T.A., Hinton, M., Lenger, D., Richards-Zawacki, C.L. (2015): *Batrachochytrium dendrobatidis* in natural and farmed Louisiana crayfish populations: prevalence and implications. *Dis. Aquat. Organ.* **112**: 229-235.
- Carvalho, T., Becker, C.G., Toledo, L.F. (2017): Historical amphibian declines and extinctions in Brazil linked to chytridiomycosis. *P. R. Soc. B*. **284**: 20162254.

- Costa, H., Bérnils, R.S. (2018): Répteis do Brasil e suas unidades federais: Lista de espécies. *Herpetologia Brasileira* **7**: 11-57.
- Daszak, P., Cunningham, A.A., Hyatt, D. (2000): Emerging infectious diseases of wildlife - Threats to biodiversity and human health. *Science* **287**: 443-449.
- Daszak, P., Zambrana-Torrelli, C., Bogich, T.L., Fernandez, M., Epstein, J.H., Murray, K.A., Hamilton, H. (2013): Interdisciplinary approaches to understanding disease emergence: the past, present, and future drivers of Nipah virus emergence. *P. Natl. Acad. Sci. USA* **110**: 3681-3688.
- Fisher, M.C., Henk, D.A., Briggs, C.F., Brownstein, J.S., Madoff, L.C., McCraw, S.L., Gurr, S.J. (2012): Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484**: 186-194.
- Fites, J.S.M., Ramsey, J.P., Holden, W.M., Collier, S.P., Sutherland, D.M., Reinert, L.K., Gayek, A.S., Dermody, T.S., Aune, T.M., Oswald-Richter, K., Rollins-Smith, L.A. (2013): The invasive Chytrid Fungus of amphibians paralyzes lymphocyte responses. *Science* **342**: 366-369.
- Garmyn, A., Van Rooij, P., Pasmans, F., Hellebuyck, T., Van Den Broeck, W., Haesebrouck, F., Martel, A. (2012): Waterfowl: Potential environmental reservoirs of the chytrid fungus *Batrachochytrium dendrobatidis*. *Plos One* **7**: e35038.
- Greenspan, S.E., Longcore, J.E., Calhoun, A.J. (2012): Host invasion by *Batrachochytrium dendrobatidis*: fungal and epidermal ultrastructure in model anurans. *Dis. Aquat. Organ.* **100**: 201-210.
- Greenspan, S.E., Lambertini, C., Carvalho, T., James, T.Y., Toledo, L.F., Haddad, C.F.B., Becker, C.G. (2018): Hybrids of amphibian chytrid show high virulence in native hosts. *Sci. Rep.* **8**: 9600.
- Hyatt, A.D., Boyle, D.G., Olsen, V., Boyle, D.B., Berger, L., Obendorf, D., Dalton, A., Kriger, K., Hero, M., Hines, H., Phillott, R., Campbell, R., Marantelli, G., Gleason, F., Colling, A. (2007): Diagnostic assays and sampling protocols for the detection of *Batrachochytrium dendrobatidis*. *Dis. Aquat. Organ.* **73**: 175-192.
- James, T.Y., Toledo, L.F., Röder, D., Leite, D.S., Belasen, A.M., Betancourt-Román, C.M., Jenkinson, T.S., Soto-Azat, C., Lambertini, C., Longo, A.V., Ruggeri, J., Collins, J.P., Burrowes, P.A., Lips, K.R., Zumudio, K.R., Longcore, J.E. (2015): Disentangling host, pathogen, and environmental determinants of a recently emerged wildlife disease: lessons from the first 15 years of amphibian chytridiomycosis research. *Ecol. Evol.* **5**: 4079-4097.
- Jenkinson, T.S., Betancourt Román, C.M., Lambertini, C., Valencia-Aguilar, A., Rodriguez, D., Nunes-de-Almeida, C.H.L., Ruggeri, J., Belasen, A.M., Silva Leite, D., Zumudio, K.R., Longcore, J.E., Toledo, L.F., James, T.Y. (2016): Amphibian-killing chytrid in Brazil comprises both locally endemic and globally expanding populations. *Mol. Ecol.* **25**: 2978-2996.
- Johnson, M.L., Berger, L., Philips, L., Speare, R. (2003): Fungicidal effects of chemical disinfectants, UV light, desiccation and heat on the amphibian chytrid *Batrachochytrium dendrobatidis*. *Dis. Aquat. Organ.* **57**: 255-260.
- Johnson, M.L., Speare, R. (2003): Survival of *Batrachochytrium dendrobatidis* in water: quarantine and disease control implications. *Emerg. Infect. Dis.* **9**: 922-925.
- Johnson, M.L., Speare, R. (2005): Possible modes of dissemination of the amphibian chytrid *Batrachochytrium dendrobatidis* in the environment. *Dis. Aquat. Organ.* **65**: 181-186.
- Kilburn, V.L., Ibáñez, R., Green, D.M. (2011): Reptiles as potential vectors and hosts of the amphibian pathogen *Batrachochytrium dendrobatidis* in Panama. *Dis. Aquat. Organ.* **97**: 127-134.
- Kruger, K.M., Ashton, K.J., Hines, H.B., Hero, J.M. (2007): On the biological relevance of a single *Batrachochytrium dendrobatidis* zoospore: a reply to Smith (2007). *Dis. Aquat. Organ.* **73**: 257-260.
- Lambertini, C., Rodriguez, D., Brito, F.B., Leite, D.S., Toledo, L.F. (2013): Diagnóstico do fungo quitridio: *Batrachochytrium dendrobatidis*. *Herpetologia Brasileira* **2**: 12-17.
- Lambertini, C., Becker, C.G., Bardier, C., Leite, D.D., Toledo, L.F. (2017): Spatial distribution of *Batrachochytrium dendrobatidis* in South American caecilians. *Dis. Aquat. Organ.* **124**: 109-116.
- Lips, K.R., Brem, F., Brenes, R., Reeve, J.D., Alford, R.A., Voyles, J., Carey, C., Livo, L., Pessier, A.P., Collins, J.P. (2006): Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *P. Natl. Acad. Sci. USA* **103**: 3165-3170.
- Longcore, J.E., Pessier, A.P., Nichols, D.K. (1999): *Batrachochytrium dendrobatidis* gen. et sp. nov., a chytrid pathogenic to amphibians. *Mycologia* **91**: 219-227.
- Lorch, J.M., Knowles, S., Lankton, J.S., Michell, K., Edwards, J.L., Kapter, J.M., Staffen, R. A., Wild, E.R., Schmidt, K.Z., Ballmann, A.E., Blodgett, D., Farrell, T.M., Glorioso, B. M., Last, L.A., Price, S.J., Schuler, K.L., Smith, C.E., Wellehan, J.F.X., Blehert, D.S. (2016): Snake fungal disease: an emerging threat to wild snakes. *Philos. T. Roy. Soc. B* **371**: 20150457.
- McMahon, T.A., Brannelly, L.A., Chatfield, M.W.H., Johnson, P.T.J., Joseph, M.B., McKenzie, V.J., Richards-Zawacki, C.L., Venesky, M.D., Rohr, J.R. (2013):

- Chytrid fungus *Batrachochytrium dendrobatidis* has nonamphibian hosts and releases chemicals that cause pathology in the absence of infection. *P. Natl. Acad. Sci. USA*. **110**: 210-215.
- Mesquita, A.F.C., Lambertini, C., Lyra, M., Malagoli, L.R., James, T.Y., Toledo, L.F., Haddad, C.F.B., Becker, C.G. (2017): Low resistance to chytridiomycosis in direct-developing amphibians. *Sci. Rep.* **7**: 16605.
- Nunes-de-Almeida, C.H.L., Zamudio, K.R., Toledo, L.F. (2016): The semiterrestrial tadpole of *Cycloramphus rhyakonastes* Heyer, 1983 (Cycloramphidae). *J. Herpetol.* **50**: 289-294.
- Olson, D.H., Ronnenberg, K.L. (2014): Global *Bd* mapping project: 2014 update. *FrogLog* **111**: 17-21.
- O'Hanlon, S. J., Rieux, A., Farrer, R. A., Rosa, G.M., Waldman, B., Bataille, A., Kosch, T.A., Murray, K.A., Brankovics, B., Fumagalli, M., Martin, M.D., Wales, N., Alvarado-Rybak, M., Bates, K.A., Berger, L., Böll, S., Brookes, L., Clare, F., Courtois, E.A., Cunningham, A.A., Doherty-Bone, T.M., Ghosh, P., Gower, D.J., Hintz, W.E., Höglund, J., Jenkinson, T.S., Lin, C.F., Laurila, A., Loyau, A., Martel, A., Meurling, S., Miaud, C., Minting, P., Pasmans, F., Schmeller, D., Schmidt, B.R., Shelton, J.M.G., Skerratt, L.F., Smith, F., Soto-Azat, C., Spagnoletti, M., Tessa, G., Toledo, L.F., Valenzuela-Sánchez, A., Verster, R., Vörös, J., Webb, R.J., Wierzbicki, C., Wombwell, E., Zamudio, K.R., Aanensen, D.M., James, T.Y., Gilbert, M.T.P., Weldon, C., Bosch, J., Balloux, F., Garner, T.W.J., Fisher, M.C. (2018): Recent Asian origin of chytrid fungi causing global amphibian declines. *Science* **360**: 621-627.
- Rodriguez, D., Becker, C.G., Pupin, N.C., Haddad, C.F.B., Zamudio, K.R. (2014): Long-term endemism of two highly divergent lineages of the amphibian-killing fungus in the Atlantic Forest of Brazil. *Mol. Ecol.* **23**: 774-787.
- Rosenblum, E.B., Voyles, J., Poorten, T.J., Stajich, J.E. (2010): The deadly chytrid fungus: a story of an emerging pathogen. *Plos Pathog.* **6**: e1000550.
- Salla, R.F., Gamero, F.U., Ribeiro, L.R., Rizzi, G.M., Medico, S.E.D., Rissoli, R.Z., Vieira, C.A., Silva-Zacarin, E.C.M., Leite, D.S., Addalla, F.C., Toledo, L.F.; Costa, M.J. (2015): Cardiac adaptations of Bullfrog tadpoles in response to chytrid infection. *J. Exp. Zool. Part A* **323**: 487-496.
- Schloegel, L.M., Toledo, L.F., Longcore, J.E., Greenspan, S.E., Vieira, C.A., Lee, M., Zhao, S., Wangen, C., Ferreira, C.M., Hipolito, M., Davies, A.J., Cuomo, C.A., Daszak, P., James, T.Y. (2012): Novel, panzootic and hybrid genotypes of amphibian chytridiomycosis associated with the bullfrog trade. *Mol. Ecol.* **21**: 5162-5177.
- Silvano, D. (2004): *Cycloramphus rhyakonastes*. The IUCN Red List of Threatened Species Available at: <http://www.iucnredlist.org/details/56378/0> [accessed on 29 June 2018].
- Shapard, E.J., Moss, A.S., San Francisco, M.J. (2012): *Batrachochytrium dendrobatidis* can infect and cause mortality in the nematode *Caenorhabditis elegans*. *Mycopathologia* **173**: 121-126.
- Turner, G.G., Reeder, D.M., Coleman, J.T.H. (2011): A five-year assessment of mortality and geographic spread of white-nose syndrome in North American bats and a look to the future. *Bat Research News* **52**: 13-27.
- Uetz, P. (2017): *Placosoma glabellum*, The Reptile Database. Available at: <http://reptile-database.reptarium.cz/species?genus=Placosoma&species=glabellum> [accessed on 5 February 2018].
- Van Rooij, P., Freddy Haesebrouck, A.M.F., Pasmans, F. (2015): Amphibian chytridiomycosis: a review with focus on fungus-host interactions. *Vet. Res.* **46**: 137.
- Voyles, J., Young, S., Berger, I., Campbell, C., Voyles, W.F., Dinudom, A., Cook, D., Webb, R., Alford, R.A., Skerratt, L.F., Speare, R. (2009): Pathogenesis of chytridiomycosis, a cause of catastrophic amphibian decline. *Science* **326**: 582-585.