

UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<http://www.medicinaoral.com/medoralfree01/aop/25874.pdf>

DOI: <https://doi.org/10.4317/medoral.25874>

Direitos autorais / Publisher's copyright statement:

©2023 by Medicina Oral. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

***Fusobacterium nucleatum* and *Prevotella* in women with periodontitis and preterm birth**

Karyne Martins Lima ¹, Claudia Maria Coelho Alves ¹, Flávia Castelo Branco Vidal ², Isaac Suzart Gomes-Filho ³, Jéssica Costa e Costa ¹, Ricardo Della Coletta ⁴, Vandilson Pinheiro Rodrigues ^{1,2}, Fernanda Ferreira Lopes ¹

¹ Postgraduate Program in Dentistry (PPGO), Federal University of Maranhão, São Luís, Brazil

² Department of Morphology, Federal University of Maranhão, São Luís, Brazil

³ Department of Health, State University of Feira de Santana, Feira de Santana, Bahia, Brazil

⁴ Department of Oral Diagnosis, School of Dentistry, University of Campinas, Piracicaba, Brazil

Correspondence:

Faculdade de Odontologia, Universidade Federal do Maranhão
Avenida dos Portugueses, 1966, Cidade Universitária Dom Delgado
São Luís, MA 65085-580, Brazil
karyne.martins@discente.ufma.br

Received: 15/12/2022

Accepted: 12/06/2023

Lima KM, Alves CM, Vidal FC, Gomes-Filho IS, Costa JC, Coletta RD, et al. *Fusobacterium nucleatum* and *Prevotella* in women with periodontitis and preterm birth. Med Oral Patol Oral Cir Bucal. 2023 Sep 1;28 (5):e450-6.

Article Number: 25874 <http://www.medicinaoral.com/>
© Medicina Oral S. L. C.I.F. B 96689336 - pISSN 1698-4447 - eISSN: 1698-6946
eMail: medicina@medicinaoral.com

Indexed in:

Science Citation Index Expanded
Journal Citation Reports
Index Medicus, MEDLINE, PubMed
Scopus, Embase and Emcare
Indice Médico Español

Abstract

Background: Studies try to explain the hypothesis that maternal periodontitis may be associated with preterm birth. **Material and Methods:** This is a case-control study with 120, 40 cases (gestational age <37 weeks) and 80 controls (gestational age ≥37 weeks), that were submitted to the clinical periodontal examination and subgingival biofilm collection. Bacterial DNA of subgingival biofilm was performed and processed by qPCR.

Results: Periodontitis was statistically significant in the Case group (35%) when compared to the Control group (11.2%) and Gingival Bleeding Index (GBI), sites with PS ≥ 4mm and sites with CAL ≥ 5mm were statistically higher in the Case group ($p < 0.05$). The proportions of *Pi* ($p = 0.026$) and *Fn* ($p = 0.041$) of subgingival biofilm were higher in the Case group. A greater number of sites with PS ≥ 4mm ($r = -0.202$; $p = 0.026$) and CAL ≥ 5mm ($r = -0.322$; $p < 0.001$) were correlated to lower gestational age.

Conclusions: Periodontitis, preterm delivery, and/or low birth weight may have a possible relationship based on clinical parameters and the ratio of *Pi* and *Fn* at periodontal sites.

Key words: Periodontal diseases, premature birth, *fusobacterium nucleatum*, *prevotella intermedia*.

Introduction

Defined as a multifactorial chronic inflammatory disease, periodontitis is associated with the presence of dysbiotic biofilm with a predominance of gram-negative anaerobic microorganisms and is mediated by the

inflammatory response of the host (1). Periodontopathogens and their products can penetrate the bloodstream, reaching tissues of distant oral cavity structures. As a result, it can induce premature birth, preeclampsia, and cause or worsen cardiovascular diseases, low birth

weight, aspiration pneumonia, and kidney diseases (2). Many researchers have investigated the association between maternal periodontitis and premature birth and/or low birth weight babies (3,4) while there is research that shows that it is not possible to prove an association between the oral condition and adverse effects on pregnancy (5). Although the exact mechanisms that clearly explain the participation of periodontitis in the adverse effects of pregnancy have not yet been established, some theories try to seek an explanation through two mechanisms: by the action of periodontal pathogens or by the effect of inflammatory mediators, such as interleukin-1 (IL-1), IL-6, IL-8, tumor necrosis factor- α (TNF- α) or prostaglandin E2 (PGE2), in the fetal-placental unit (6). Among the bacteria associated with periodontitis are the anaerobic gram-negative, *Prevotella intermedia*, and *Fusobacterium nucleatum* (7).

The presence of DNA from periodontopathogens has already been detected in the genital tract, amniotic fluid, and placental tissues, which supports how bacterial species of oral origin can participate in placental and intrauterine infections (8,9). Thus, the present study aimed to analyze whether there is a relationship between the clinical and microbiological parameters of periodontitis and the birth of premature babies, quantifying the pathogens *Prevotella intermedia* (Pi) and *Fusobacterium nucleatum* (Fn) in the subgingival biofilm of postpartum women to verify if there is a relationship between the clinical and microbiological parameters of periodontitis and the birth of premature babies.

Material and Methods

- Ethical Considerations

This research was approved by the Research Ethics Committee of the University Hospital of the Federal University of Maranhão under protocol number 002673/2011-60.

- Study design

A case-control study was carried out at the Maternal and Child Unit of the University Hospital of the Federal University of Maranhão, with puerperal women at the bedside attended up to 48 hours postpartum. Information about the mother's and newborn's health conditions were obtained from the medical records that were made available to the researchers at the time of collection.

The Case Group (A1) was formed by mothers of children born with a gestational age < 37 weeks and the Control Group (A2) by mothers of newborns with a gestational age \geq 37 weeks.

Women smokers were excluded; alcoholics; those who used fixed orthodontic appliances; mouth breathers; chronic use of antibiotics or non-steroidal or steroidal anti-inflammatory drugs in the 6 months preceding the survey; the presence of a systemic condition that could influence the assessment (diabetes mellitus, systemic

arterial hypertension, obesity, viral and bacterial infections); use of antihypertensive, anticonvulsant or immunosuppressant drugs or any other drug that is known to present the possibility of resulting in gingival tissue hyperplasia, twin pregnancy.

- Clinical Stage

A previously trained professional applied the questionnaire data collected on oral hygiene habits (performing or not brushing after meals and its frequency, using or not flossing and its frequency, and whether you visited the dentist in the last year). The clinical examination was performed with the aid of a frontal light attached to the examiner's head. The following parameters were evaluated: 1- Probing Depth (PS) recorded at 6 sites for each tooth (mesiobuccal and mesiolingual, distobuccal and distolingual and in the mid-buccal and mid-lingual region), 2 - Clinical Attachment Level (CAL) distance in millimeters between the cemento-enamel junction and the bottom of the sulcus or periodontal pocket (6 sites), 3- Visible Plaque Index (VPI) (buccal, mesial, distal and lingual), 4- Gingival Bleeding Index (6 sites), recorded 10 seconds after probe removal North Carolina PC-PUNC 156 (Hu-Friedy, Chicago, USA) (Hu-Friedy, Chicago, USA) of the periodontal pocket or sulcus (10,11). Each postpartum woman was classified according to the presence or absence of periodontitis according to the following criteria: at least 4 or more teeth with one or more sites with PS \geq 4 mm, with CAL \geq 3 mm in the same site, and the presence of bleeding on probing (12).

- Biofilm Collection

The collection of the subgingival biofilm was performed after the removal of the supragingival biofilm with sterile gauze. Samples were taken with Gracey 5-6 / 7-8 periodontal curettes (Hu Friedy, Chicago, USA) from four different sites (those with the greatest probing depth of each quadrant) throughout the same depth. The biofilms were placed in sterile tubes containing a Tris-EDTA solution (150 μ L Tris-EDTA + 100 μ L of NaOH solution) for freezing in a -80°C freezer (Indrel Ultra Freezer -80°C).

- DNA Extraction

DNA extraction was performed as described by the QIAamp DNA User Manual (Qiagen), with modifications. 200 μ L of the sample, 20 μ L of proteinase K and 200 μ L of Buffer AL were added to a 2ml tube and incubated at 56°C for 10 minutes in a dry bath. Then 200 μ L of ethanol (96-100%) (Vetec Química, Brazil) was added and the entire solution was added to a silica column for DNA capture.

500 μ L of washing solutions were added followed by centrifugation and finally, the column was incubated in 200 μ L of buffer AE for 5 minutes. DNA was collected after centrifugation at 8000rpm for 1 minute. The purified DNA was submitted to quantification in the Nanovue device (GE, USA) to evaluate the concentration and integrity of the DNA. All material was frozen in

-80°C refrigerator Indrel Ultra Freezer -86°C).

- Detection and Quantification by Real-Time Polymerase Chain Reaction (qPCR)

For the detection and quantification of the bacteria under study, the DNA of the bacteria needs positive controls, as these are necessary to generate the standard curves of the bacterial species used in the biofilm model. Concentrations of 1-0.001ng of DNA were defined for the standard curve generation. The logarithm of the corresponding values from the quantification cycle was used to obtain a linear regression. For the quantification of bacteria in the biofilm samples, the concentration of the extracted DNA mixture was determined using the NanoDrop ND-1000.

After defining the standard curve, qPCR of the DNA of the samples was performed with the 16S ribosomal RNA gene primer of all the bacteria to control the quality of the sample's DNA. The specific PCR primers *Prevotella intermedia* (Pi) (5'-TTTGTGGGGAGTA-AAGCGGG-3') and *Fusobacterium nucleatum* (Fn) (5'-ATT GTC GCT AAA AAT T-3') were used (13).

All reactions were quantified individually in separate wells for each bacterial species and performed in triplicate. The qPCR was performed in a total reaction volume of 20 µL, containing 10 µL of SYBR® Green PCR Master Mix (Life Technologies, Zug, Switzerland), 5 µL of the sample (diluted to contain 1 ng of DNA), and 5 µL of solution of primer from the respective bacterium (10µM, a mixture of 3' and 5' primers).

Amplification took place in a StepOne thermocycler (Applied Biosystems) by initial incubation of 2 min at 50 °C and 10 min at 95 °C, followed by 40 cycles of 15 s at 95 °C and 1 min at 60 °C. Ct obtained, the DNA concentration of the sample was calculated for each or-

ganism using the theoretical genomic weight of each of the bacteria (13).

- Statistical analysis

Data were analyzed using GraphPad Prism version 8 software (GraphPad Software, San Diego, USA). Descriptive statistics were performed using measures of frequency, mean and standard deviation. The results were presented through tables, bar graphs, and heat maps. Categorical variables were compared between case and control groups using the chi-square or Fisher's exact test. Normality analysis of continuous variables was performed using the Shapiro-Wilk test. After this procedure, the independent student t-test was selected for the comparative analysis between the groups. In addition, Pearson's coefficient (r) was used to analyze the correlation between the variables gestational age (in weeks), the proportion of pathogens, and periodontal parameters. The significance level adopted for all analyzes was 5% ($p < 0.05$).

Results

The birth weight of the Case group was statistically lower compared to the Control group ($p < 0.001$). The Case group also observed a lower mean number of prenatal consultations ($p = 0.045$). There was a concentration of younger women in the Control group ($p = 0.014$). Furthermore, arterial hypertension was more prevalent in the Case group ($p < 0.001$) (Table 1).

The evaluation of the periodontal condition, expressed in Table 2, revealed that the prevalence of periodontitis was statistically significant in the Case group (35%) when compared to the Control group (11.2%). It was also observed that the parameters of Gingival Bleeding Index (GSI), sites with PS ≥ 4 mm, and sites with CAL ≥ 5 mm were statistically higher in the Case group ($p < 0.05$).

Table 1: Distribution of general characterization variables between groups.

Variables			Groups		p value
			Case <37 weeks	Control ≥ 37 weeks	
Newborn data	Gender	Male	20 (50,0)	31 (38,8)	0,239
		Female	20 (50,0)	49 (61,2)	
	Race/Color skin	White	13 (32,5)	41 (51,2)	0,102
		Black	3 (7,5)	3 (3,8)	
		Brown	18 (45,0)	32 (40,0)	
		Indigenous	6 (15,0)	4 (5,0)	
	Birth weight (kg)		2,1 \pm 0,6	3,2 \pm 0,7	<0,001*
Maternal data	Prenatal consultation number		4,7 \pm 1,9	5,6 \pm 2,3	0,045*
	Age group	Up to 20 years	4 (10,0)	26 (32,5)	0,014*
		21 to 30 years	22 (55,0)	35 (43,8)	
		31 to 35 years	12 (30,0)	11 (13,7)	
		More than 35 years	2 (5,0)	8 (10,0)	
	Arterial hypertension	No	20 (50,0)	67 (83,8)	<0,001*
		Yes	20 (50,0)	13 (16,2)	

Data were presented as n (%) or mean \pm standard deviation. *Statistically significant difference ($p < 0.05$).

Table 2: Comparative analysis of oral hygiene variables and periodontal condition between groups.


Variables		Age gestational		p value
		Case <37 semanas	Control ≥37 semanas	
Daily tooth brushing	1 time a day	6 (15,0)	7 (8,7)	0,287
	2 times a day	9 (22,5)	28 (35,0)	
	3 times a day or more	25 (62,5)	45 (56,3)	
Brushing after meals	No	0 (0)	2 (2,5)	0,549
	Yes	40 (100)	77 (97,5)	
Dental floss / tape	do not use	29 (72,5)	51 (63,8)	0,337
	Uses	11 (27,5)	29 (36,2)	
Dental consultation during pregnancy	No	32 (80,0)	67 (83,8)	0,610
	Yes	8 (20,0)	13 (16,2)	
Periodontitis	Absent	26 (65,0)	71 (88,8)	0,001*
	Gift	14 (35,0)	9 (11,2)	
Periodontal parameters	VPI	41,2 ±22,9	39,7 ±20,8	0,720
	GSI	14,7 ±18,6	8,2 ±12,8	0,026*
	Sites with PS ≥4mm	3,1 ±4,1	1,1 ±1,8	<0,001*
	Sites with CAL ≥5mm	0,9 ±1,8	0,2 ±0,9	0,008*

Data were presented as n (%) or mean±standard deviation. VPI = Visible Plaque Index. GSI = gingival bleeding index. CAL = clinical attachment level. PS = probing depth. *Statistically significant difference ($p < 0.05$).

Fig. 1 illustrates the distribution of pathogens in the samples evaluated. There were no significant differences between the groups when comparing the total amount of *Pi* and *Fn*. On the other hand, the analysis of the proportion of pathogens in the samples revealed that the proportions of *Pi* ($p = 0.026$) and *Fn* ($p = 0.041$) were higher in the Case group.


Linear correlation analysis is illustrated in the heat map

(Fig. 2). The results showed that the greater the number of sites with PS ≥ 4mm ($r = -0.202$; $p = 0.026$) and CAL ≥ 5mm ($r = -0.322$; $p < 0.001$), to lower the gestational age. The proportions of *Pi* and *Fn* showed a strong direct correlation ($r = 0.941$; $p < 0.001$). The VPI showed an inverse correlation with the proportion of *Fn* ($r = -0.206$; $p = 0.023$). In addition, all periodontal parameters showed significant direct correlations with each other.



*Indicates statistically significant differences ($P < 0.05$)

Fig. 1: Comparative analysis of the amount and proportion in the biofilm of the pathogens *Prevotella intermedia* (a and b) and *Fusobacterium nucleatum* (c and d).



GA = gestational age in weeks. % Pi = Proportion of *Prevotella intermedia*. % Fn = Proportion of *Fusobacterium nucleatum*. VPI = visible plaque index. GBI = gingival bleeding index. CAL 5mm = number of sites with clinical attachment level ≥ 5 mm. PPD 4mm = number of sites with probing depth ≥ 4 mm. PPD 4mm, CAL 5mm

Fig. 2: Pearson correlation analysis between gestational age (in weeks), pathogens and periodontal variables.

Discussion

The study showed the relationship between periodontitis and preterm delivery and/or low birth weight, through clinical and microbiological parameters of periodontal disease. It is noteworthy that the analysis of the proportion of pathogens in the samples revealed higher proportions of *Pi* and *Fn* in postpartum women with preterm delivery.

It is noteworthy that in both groups there were no significant differences in the total amount of *Pi* and *Fn* between groups, however, the proportion of *Pi* ($p=0.026$) and *Fn* ($p=0.041$) within the universal DNA community of the subgingival biofilm of postpartum women were higher in the case group. Data partially corroborate another case-control study with pregnant women in which the results found an association in the clinical presentation of periodontitis, but in the microbial analysis was no significant association between preterm delivery and low birth weight with most periodontal pathogens (14). A possible explanation for this divergence may lie in the method used for microbial analysis by PCR (15) that detects only presence or absence and by qPCR, as in the present study.

Another relevant aspect is the methodology presented here by the use of the 16S universal primer, which allowed the identification of bacterial phenotypic sequence present in most bacteria, with subsequent use of a bacterial primer specific to the *Fn* and *Pi* pathogens (16). This fact clarifies that the total amount of *Pi* and *Fn* is not significant between the two groups (case-control), however, the analysis of the proportion of pathogens in the samples revealed higher proportions of *Pi* and *Fn* in the group with preterm and/or low delivery birth weight.

Case-control studies have associated the presence of oral pathogens, including *Fn* and *Pi*, with hypertensive disorders during pregnancy, as they have been shown to have higher levels in hypertensive women than in normotensive patient groups (9,17). In the present study, it was found that there were more hypertensive mothers among the case group participants, a situation that corroborates the literature on the birth of premature and/or low birth weight babies (8). In addition, high levels of bacteria and viruses in the subgingival and placental samples occur more in women

with preeclampsia and periodontitis compared to women with preeclampsia and without periodontitis (17). The number of prenatal consultations was lower in the case group and this data is similar in part to the findings of Martínez-Martínez *et al* (18) who carried out a study to identify pathogens in biofilms by PCR. They concluded that the main periodontal bacteria are not related to premature birth, but that mothers in the case group show less care for hygiene and health, such as lower number of prenatal consultations, higher level of supragingival biofilm, lower toothbrushing frequency, higher levels of active caries and more need for dental treatment. Thus, they suggest that preterm birth would be multifactorial and the only presence of periodontal bacteria would not be able to trigger it (18).

Fn is one of the most prevalent oral species related to adverse effects in pregnancy (19). It has been detected in a wide variety of placental and fetal tissues, including amniotic fluid, fetal membranes, umbilical cord blood, neonatal gastric aspirate, fetal lung, and stomach, associated with chorioamnionitis, preeclampsia, preterm birth, stillbirth, and early neonatal sepsis (19,20).

Periodontitis is attributed to interactions between host immunity and multiple microbial species, including *P. gingivalis*, *F. nucleatum*, *P. intermedia*, *A. actinomycetemcomitans*, *T. denticola*, in the subgingival environment, and the DNA of periodontal pathogens is highly detectable, in patients with periodontitis compared to controls (16). To date, the study by Doyle *et al* (21) can be considered the largest in the number of participants with more than 1000 individuals in the placental microbiota analysis, which found bacterial DNA in more than 50% of placental tissues, revealing differences between the oral and vaginal microbiota being the vaginal microbiota with the highest association with adverse effects on pregnancy in their findings. However, it is noteworthy that periodontal pathogens were detected simultaneously in pregnant women's subgingival biofilm and the placenta (22).

The periodontal clinical and microbiological parameters of postpartum women showed periodontitis as a factor associated with preterm birth, and these mothers had babies with lower weights compared to the control group. Some studies (23-26) reported an association between periodontitis and preterm birth or low birth weight, while others found no association. The different results may be explained by the use of different parameters for the diagnosis of periodontitis as well as the use of different terms: premature newborn, low birth weight, or premature and low birth weight (27).

It is important to highlight that a systematic review of 10 case-control studies concluded that there is a relationship between periodontal disease and increased risk of preterm birth and low birth weight or low birth weight with Odds Ratios ranging between 2.04 and 4.19, and

in this work, there was only one study that found no association (28). Thus, the results presented here such as the GSI parameters, sites with PS \geq 4mm, and sites with CAL \geq 5mm, which were statistically higher in the Case group ($p < 0.05$) support the relationship between periodontitis and adverse effects of pregnancy.

For Fischer *et al* (29), it is premature to rule out the association between periodontitis and adverse pregnancy outcomes by studies that showed failure in periodontal treatment to reduce adverse pregnancy outcomes. One possible hypothesis is that periodontal pathogens spread to the placental tissue before periodontal treatment. It is important to consider that the association must be related to the presence of oral pathogens in the placenta and the amount, prevalence, and exposure time.

Other studies also sought answers to possible failures in the work to explain the non-association between periodontitis and adverse effects on pregnancy, such as heterogeneity in the clinical definition of the severity and extent of periodontitis used to distinguish case and control groups. Studies fail to control for shared risk factors or confounding factors between periodontitis and adverse effects on pregnancy, and finally, most studies did not consider the spread and survival of periodontal pathogens to the placenta as a mechanism that could induce these effects regardless of ongoing disease in the oral cavity (30).

This research presents strong bridges such as the criteria used for the diagnosis of periodontitis, the sample size respecting the case and control groups, and the robust technique for detecting periodontal pathogens through qPCR. The findings contribute to the hypothesis of an association between *Pi* and *Fn* bacteria and the prematurity outcome, however, we were unable to show a causal relationship.

Conclusions

Periodontitis and preterm delivery may have a possible relationship based on clinical parameters and the ratio of *Pi* and *Fn* at periodontal sites, contributing to the multifactorial nature of preterm delivery. However, we cannot assert forcefully the cause-and-effect relationship between the objects studied in the research. Thus, more studies are needed to corroborate the preliminary data of the present study.

References

1. Caton JG, Armitage G, Berglundh T, Chapple ILC, Jepsen S, Kornman KS, *et al*. A new classification scheme for periodontal and peri-implant diseases and conditions - Introduction and key changes from the 1999 classification. *J Clin Periodontol*. 2018;45:S1-8.
2. Bobetsis YA, Graziani F, Gürsoy M, Madianos PN. Periodontal disease and adverse pregnancy outcomes. *Periodontol* 2000. 2020;83:154-74.
3. Offenbacher S, Katz V, Fertik G, Collins J, Boyd D, Maynor G, *et al*. Periodontal Infection as a Possible Risk Factor for Preterm Low Birth Weight. *J Periodontol*. 1996;67:1103-13.

4. Figuero E, Han YW, Furuichi Y. Periodontal diseases and adverse pregnancy outcomes: Mechanisms. *Periodontol* 2000. 2020;83:175-88.
 5. Chaparro A, Blanlot C, Ramirez V, Sanz A, Quintero A, Inostroza C, *et al*. Porphyromonas gingivalis, Treponema denticola and toll-like receptor 2 are associated with hypertensive disorders in placental tissue: a case-control study. *J Periodontol Res*. 2013;48:802-9.
 6. Vander Haar EL, So J, Gyamfi-Bannerman C, Han YW. *Fusobacterium nucleatum* and adverse pregnancy outcomes: Epidemiological and mechanistic evidence. *Anaerobe*. 2018;50:55-9.
 7. Saadaoui M, Singh P, Al Khodor S. Oral microbiome and pregnancy: A bidirectional relationship. *J Reprod Immunol*. 2021;145:103293.
 8. Kayar NA, Alptekin NÖ, Erdal ME. Interleukin-1 receptor antagonist gene polymorphism, adverse pregnancy outcome and periodontitis in Turkish women. *Arch Oral Biol*. 2015;60:1777-83.
 9. Barak S, Oettinger-Barak O, Machtei EE, Sprecher H, Ohel G. Evidence of Periopathogenic Microorganisms in Placentas of Women With Preeclampsia. *J Periodontol*. 2007;78:670-6.
 10. Ainamo J, Bay I. Problems and proposals for recording gingivitis and plaque. *Int Dent J*. 1975;25:229-35.
 11. Pihlstrom BL. Measurement of Attachment Level in Clinical Trials: Probing Methods. *J Periodontol*. 1992;63:1072-7.
 12. Gomes-Filho IS, Cruz SS, Rezende EJC, dos Santos CAST, Soledade KR, Magalhães MA, *et al*. Exposure measurement in the association between periodontal disease and prematurity/low birth weight. *J Clin Periodontol*. 2007;34:957-63.
 13. Kirakodu SS, Govindaswami M, Novak MJ, Ebersole JL, Novak KF. Optimizing qPCR for the Quantification of Periodontal Pathogens in a Complex Plaque Biofilm. *Open Dent J*. 2008;2:49-55.
 14. Calixto NR, Alves C, Abreu L, Thomaz EB, Vidal F, Gomes-Filho IS, *et al*. Detection of periodontal pathogens in mothers of preterm birth and/or low weight. *Med Oral Patol Oral Cir Bucal*. 2019;24:e776-81.
 15. Choi BK, Park SH, Yoo YJ, Choi SH, Chai JK, Cho KS, *et al*. Detection of Major Putative Periodontopathogens in Korean Advanced Adult Periodontitis Patients Using a Nucleic Acid-Based Approach. *J Periodontol*. 2000;71:1387-94.
 16. Swati P, Thomas B, Vahab SA, Kapaettu S, Kushtagi P. Simultaneous detection of periodontal pathogens in subgingival plaque and placenta of women with hypertension in pregnancy. *Arch Gynecol Obstet*. 2012;285:613-9.
 17. Tanneeru S, Mahendra J, Shaik M. Evaluation of Microflora (Viral and Bacterial) in Subgingival and Placental Samples of Pregnant Women with Preeclampsia with and without Periodontal Disease: A Cross-Sectional Study. *J Int Soc Prev Community Dent*. 2020;10:171.
 18. Martínez-Martínez RE, Moreno-Castillo DF, Loyola-Rodríguez JP, Sánchez-Medrano AG, Miguel-Hernández JHS, Olvera-Delgado JH, *et al*. Association between periodontitis, periodontopathogens and preterm birth: is it real?. *Arch Gynecol Obstet*. 2016;294:47-54.
 19. Casarin RC V., Barbagallo A, Meulman T, Santos VR, Sallum EA, Nociti FH, *et al*. Subgingival biodiversity in subjects with uncontrolled type-2 diabetes and chronic periodontitis. *J Periodontol Res*. 2013;48:30-6.
 20. Wang X, Buhimschi CS, Temoin S, Bhandari V, Han YW, Buhimschi IA. Comparative Microbial Analysis of Paired Amniotic Fluid and Cord Blood from Pregnancies Complicated by Preterm Birth and Early-Onset Neonatal Sepsis. *PLoS One*. 2013;8:e56131.
 21. Doyle RM, Harris K, Kamiza S, Harjunmaa U, Ashorn U, Nkhoma M, *et al*. Bacterial communities found in placental tissues are associated with severe chorioamnionitis and adverse birth outcomes. *PLoS One*. 2017;12:e0180167.
 22. DiGiulio DB. Diversity of microbes in amniotic fluid. *Semin Fetal Neonatal Med*. 2012;17:2-11.
 23. Leal ASM, de Oliveira AEF, Brito LMO, Lopes FF, Rodrigues VP, Lima KF, *et al*. Association between Chronic Apical Periodontitis and Low-birth-weight Preterm Births. *J Endod*. 2015;41:353-7.
 24. Reza Karimi M, Hamissi JH, Naeini SR, Karimi M. The Relationship Between Maternal Periodontal Status of and Preterm and Low Birth Weight Infants in Iran: A Case Control Study. *Glob J Health Sci*. 2015;8:184.
 25. Pozo E, Mesa F, Ikram MH, Puertas A, Torrecillas-Martínez L, Ortega-Oller I, *et al*. Preterm birth and/or low birth weight are associated with periodontal disease and the increased placental immunohistochemical expression of inflammatory markers. *Histol Histopathol*. 2016;31:231-7.
 26. Sugita N, Kobayashi T, Kikuchi A, Shimada Y, Hirano E, Sasahara J, *et al*. Immunoregulatory gene polymorphisms in Japanese women with preterm births and periodontitis. *J Reprod Immunol*. 2012;93:94-101.
 27. Abati S, Villa A, Cetin I, Dessole S, Lugliè PF, Strohmenger L, *et al*. Lack of association between maternal periodontal status and adverse pregnancy outcomes: a multicentric epidemiologic study. *J Matern Fetal Neonatal Med*. 2013;26:369-72.
 28. Teshome A, Yitayeh A. Relationship between periodontal disease and preterm low birth weight: systematic review. *Pan African Medical Journal*. 2016;24:215.
 29. Fischer LA, Demerath E, Bittner-Eddy P, Costalonga M. Placental colonization with periodontal pathogens: the potential missing link. *Am J Obstet Gynecol*. 2019;221:383-392.e3.
 30. Fogacci MF, Cardoso E de OC, Barbirato D da S, de Carvalho DP, Sansone C. No association between periodontitis and preterm low birth weight: a case-control study. *Arch Gynecol Obstet*. 2018;297:71-6.
- Funding**
This work was carried out with the support of the Brazilian Coordination for the Improvement of Higher Education Personnel (CAPES, Finance Code 001 and PROCAD 552434/2011-7) and the Foundation for Support to Research and Scientific and Technological Development of Maranhão (FAPEMA, Finance Code UNIVERSAL-00707/14).
- Conflict of interest**
The authors declare that they have no conflict of interest.
- Ethics**
This research was approved by the Research Ethics Committee of the University Hospital of the Federal University of Maranhão under protocol number 002673/2011-60.
- Authors contributions**
Study conception and design: ISGF, CMCA, FFL
Data Collection: JCC, KML
Statistical Analysis and interpretation of results: FCBV, RDC, VPR
Draft manuscript preparation: KML, FFL