

1.2799

RT-IMECC
IM/4080

COMPACTNESS OF TOPOLOGICAL
SPACES OF MODELS


Xavier Caicedo

Dezembro

RP 64/92

Relatório de Pesquisa

**Instituto de Matemática
Estatística e Ciência da Computação**



**UNIVERSIDADE ESTADUAL DE CAMPINAS
Campinas - São Paulo - Brasil**

R.P.
IM/64/92



2000870685

IMECC

Periódico

ABSTRACT – We study $[k, \lambda]$ -compactness of topological spaces in general and spaces of structures associated to abstract logics in particular. A characterization in terms of ultrafilter convergence of preservation of $[k, \lambda]$ -compactness by products of general topological spaces is given. The Abstract Compactness Theorem of Makowsky and Shelah becomes then a simple corollary from a general topological result. Reciprocally, several results on compactness shown first for logics are seen to hold true in all topological spaces.

IMECC – UNICAMP
Universidade Estadual de Campinas
CP 6065
13081 Campinas SP
Brasil

O conteúdo do presente Relatório de Pesquisa é de única responsabilidade do autor.

Dezembro – 1992

I. M. E. C. C.
BIBLIOTECA

Compactness of topological spaces of models

Xavier Caicedo

Campinas, November 1992

We discuss various forms of compactness in model theoretical logics, their inter-relations, and consequences, trying to separate clearly purely topological facts from genuine model theoretical facts. We discuss in this context results due to Shelah, Makowsky, and Mundici. Some of them are just special cases of general topological results, others reveal an interesting topological behaviour of the spaces of structures generated by a logic. In particular, inspired by the "Abstract Compactness Theorem" of Makowsky and Shelah [M-Sh], we prove a characterization in terms of ultrafilter convergence of preservation of $[\kappa, \lambda]$ -compactness by products of general topological spaces. The Abstract Compactness Theorem becomes then a simple corollary from a general topological result. Reciprocally, several results on compactness shown first for logics are seen to hold true in all topological spaces.

This work was realized with the support of FAPESP (Fundacao de Amparo a Pesquisa do Estado de Sao Paulo) while a Visiting Professor at the "Universidade Estadual de Campinas" the second semester of 1992. I received generous logistic support from the "Centro de Logica Epistemologia e Historia da Ciencia" and the "Instituto de Matematica, Estatistica e Ciencia da Computacao" of UNICAMP. Special thanks are due to A.M. Sette and C. Cifuentes for many stimulating ideas and discussions and their active participation in the seminary which originated this notes.

I. TOPOLOGICAL PRELIMINARIES, $[\kappa, \lambda]$ -COMPACTNESS

The following generalization of the notion of compactness of a topological space was considered first by Alexandroff and Urysohn [AU], and studied later by Smirnov [S], Gaal [G] and Vaughan [V1], see also Kunen-Vaughan [Ku-V]. The identity between these notions and those introduced by Makowsky and Shelah [Ma-Sh] in logic was noticed first by Mannila [M].

Definition 1.1. A topological space X is said to be $[\kappa, \lambda]$ -compact, for $\omega \leq \lambda \leq \kappa \leq \infty$, if any covering of X by a family of at most κ open sets has a covering subfamily of power less than λ . Equivalently, any family of at most κ closed subsets of X for which every subfamily of power less than λ has non-empty intersection (we will call this the λ -intersection property) has itself non-empty intersection.

$[\infty, \omega]$ -compactness is ordinary compactness of topological spaces, $[\omega, \omega]$ -compactness is usually called *countable compactness* in topology, and $[\infty, \omega_1]$ -compactness is the so called *Lindelof property*.

The reader should be aware that the notation utilized for $[\kappa, \lambda]$ -compactness in model theory, which we adopt here, reverses the more natural notation utilized in the topological literature.

LEMMA 1.2. X is $[\kappa, \lambda]$ -compact if and only if it is $[\mu, \mu]$ -compact for any $\mu \leq \kappa$.

Proof. $[\kappa, \lambda]$ -compactness implies trivially $[\mu, \mu]$ -compactness for $\lambda \leq \mu \leq \kappa$. Now, if X is not $[\kappa, \lambda]$ -compact, let $\{C_\alpha\}_{\alpha < \mu}$ be a family of closed sets giving a counterexample of minimal cardinality μ to $[\mu, \lambda]$ -compactness. It is in fact a counterexample to $[\mu, \mu]$ -compactness because an intersection $\bigcap_{i < \delta} C_{\alpha_i}$ with $\delta < \mu$ is non-empty: for $\delta < \lambda$ by hypothesis, and for $\lambda \leq \delta < \mu$ because otherwise we would have a counterexample to $[\lambda, \delta]$ -compactness contradicting minimality of μ . \square

Therefore, we may reduce to the study of $[\kappa, \kappa]$ -compactness. This takes a very simple and useful form if κ is regular cardinal. First we introduce a stronger notion of κ -compactness.

Definition 1.3. A space X is κ -chain compact if and only if any descending chain $\{C_\alpha\}_{\alpha < \kappa}$ of non-empty closed sets of X has non-empty intersection.

LEMMA 1.3. a) κ -chain compactness implies $[\kappa, \kappa]$ -compactness.

b) If $\text{cof}(\kappa) \geq \lambda$ and X is $[\kappa, \lambda]$ -compact then it is κ -chain compact.

c) (Alexandroff and Urysohn [AU]) If κ is regular, then X is $[\kappa, \kappa]$ -compact if and only if X is κ -chain compact.

Proof. a) If $\{C_\alpha\}_{\alpha < \kappa}$ is a counterexample to $[\kappa, \kappa]$ -compactness then the family $\{D_\alpha\}_{\alpha < \kappa}$ where $D_\alpha = \bigcap_{\beta < \alpha} C_\beta$ is a counterexample to κ -chain compactness.

b) Given a descending chain of closed non-empty closed sets $\{C_\alpha\}_{\alpha < \kappa}$, then the intersection of less than λ many C_α 's contains a C_β which is non-empty by hypothesis; hence, the hypothesis of $[\kappa, \lambda]$ -compactness applies and the full intersection is non-empty.

c) From (a) and (b). \square

COROLLARY 1.4. *If X is $[\text{cof}(\kappa), \text{cof}(\kappa)]$ -compact then it is $[\kappa, \kappa]$ -compact.*

Proof. If X is not $[\kappa, \kappa]$ -compact, then it is not κ -chain compact. Let $\{C_\alpha\}_{\alpha < \kappa}$ be a descending chain providing a counterexample to κ -chain compactness, then any cofinal descending subchain of length $\text{cof}(\kappa)$ provides a counterexample to $\text{cof}(\kappa)$ -chain-compactness, and so to $[\text{cof}(\kappa), \text{cof}(\kappa)]$ -compactness by Lemma 1.3(c), since $\text{cof}(\kappa)$ is regular. \square

COROLLARY 1.5. a) (Alexandroff and Urysohn [AU]). *X is $[\kappa, \omega]$ -compact if and only if it is $[\mu, \mu]$ -compact for any infinite regular cardinal $\mu \leq \kappa$.*

b) (Vaughan [V2]). *If $\text{cof}(\kappa) \geq \lambda$, then X is $[\kappa, \lambda]$ -compact if and only if it is $[\mu, \lambda]$ -compact for any infinite regular cardinal $\mu \leq \kappa$.*

Proof. a) One direction is trivial. For the other notice that by hypothesis, X is $[\text{cof}(\mu), \text{cof}(\mu)]$ -compact and so $[\mu, \mu]$ -compact by Corollary 1.4, for any $\mu \leq \kappa$. Now apply Lemma 1.2.

b) One direction is trivial. For the other notice that $[\text{cof}(\kappa), \text{cof}(\kappa)]$ -compactness holds by hypothesis; hence, $[\kappa, \kappa]$ -compactness holds by Corollary 1.4. Now, let $\lambda \leq \mu < \kappa$, then $[\mu^+, \lambda]$ -compactness holds by hypothesis and so $[\mu, \mu]$ -compactness holds, then apply Lemma 1.2 \square

Remark. It follows from the above corollary, part (a), that a space X is compact if and only if it is $[\mu, \mu]$ -compact for any regular $\mu \leq \text{Weigh}(X)$.

Definition 1.6. Given a subset S of a topological space X , a point $x \in X$ will be called a κ -accumulation point of S in X if and only if for any open neighborhood V of x we have $|V \cap S| \geq \kappa$.

THEOREM 1.7. (Alexandroff and Urysohn, [AU]). *Let κ be a regular cardinal. A space X is $[\kappa, \kappa]$ -compact if and only if any subset $S \subseteq X$ of power $\geq \kappa$ has a κ -accumulation point.*

Proof. Assume X is $[\kappa, \kappa]$ -compact. Given $x_\alpha \in S$, $x_\alpha \neq x_\beta$ for $\alpha < \beta < \kappa$, let $C_\alpha = \text{Cl}\{x_\beta : \beta \geq \alpha\}$; then the C_α form a descending chain of nonempty closed sets. By Corollary 1.3(c) there is a point x in the intersection of the C_α and so in the adherence of each $\{x_\beta : \beta \geq \alpha\}$ for any $\alpha < \kappa$. Any neighborhood V of x contains then a sequence $\{x_{\beta_\alpha} : \alpha < \kappa\}$ with $\beta_\alpha > \alpha$. By regularity of κ again the sequence $\{\beta_\alpha : \alpha < \kappa\}$ and so the family $\{x_{\beta_\alpha} : \alpha < \kappa\}$ has power κ .

Conversely, assume that $[\kappa, \kappa]$ -compactness fails, then by Corollary 1.3(c), there is a descending chain of non-empty closed sets $\{C_\alpha\}_{\alpha < \kappa}$ with empty intersection. We may pick

$x_\alpha \in C_\alpha$ all distinct, otherwise the sequence would be finally constant and the intersection would not be empty. Let x be a κ -accumulation point of the family $S = \{x_\alpha\}_{\alpha < \kappa}$, then any neighborhood V of x intersects a subsequence of S of power κ which evidently must be cofinal. Therefore, V intersects a cofinal sequence of the C_α and so all the C_α 's, which shows x is in the adherence of all C_α and so in their intersection because they are closed. This is a contradiction. \square

In some respects $[\kappa, \lambda]$ -compactness has the same good behaviour of full compactness. The following is obvious.

LEMMA 1.8. a) *If $f: X \rightarrow Y$ is continuous and X is $[\kappa, \lambda]$ -compact, then $f(X)$ is $[\kappa, \lambda]$ -compact.*
 b) *A closed subspace of a $[\kappa, \lambda]$ -compact space is $[\kappa, \lambda]$ -compact.*

In others, it behaves badly. Thichonoff's theorem fails. Product of $[\kappa, \lambda]$ -compact spaces is rarely $[\kappa, \lambda]$ -compact. For example, a product of countably compact spaces is not necessarily countably compact, neither a product of Lindelof spaces needs to be Lindelof. See Vaughan [V1] for a discussion of this matter.

II. COMPACTNESS OF SPACES OF STRUCTURES

For the definition of a logic for first order structures see Lindstrom [L] or Ebbinghaus [E]. We will always assume that logics are *regular*, that is they contain first order logic, $L_{\omega\omega}$, and are closed at least under negations, finite conjunctions, substitutions, and relativizations. For a logic L , the *domain of L* , $\text{Dom}(L)$, will be the class of vocabularies σ where $L(\sigma)$ is defined. $L(\sigma)$ will be always assumed to be a set, that is we consider only *small* logics.

A logic L induces a topology in the class E_σ of structures of type σ , having for basis of open classes the L -elementary classes, for each $\sigma \in \text{Dom}(L)$. The closed classes have the form $\text{Mod}(T)$ for some theory $T \subseteq L(\sigma)$. We call $E_\sigma(L)$ to the corresponding large topological space. This space is a proper class and the same happens to the open and closed classes of the topology, but the basis is parametrized by the set $L(\sigma)$ and so the topology is also (parametrized by) a set; hence we may apply the ordinary concepts and result of topology without missgivings. These spaces are uniform with the following canonical uniformity: for each finite theory $F \subseteq L(\sigma)$ a basic of the uniformity is given by

$$U_F = \{ (A, B) \subseteq E_\sigma : A \models \phi \text{ if and only if } B \models \phi, \text{ for any } \phi \in F \}.$$

See [C1] where the interrelation between compactness and normality of spaces of structures is studied, or [C2] where we discuss the uniform continuity of operations between spaces of structures, and its model theoretical meaning.

Definition 2.1. A logic L is said to be $[\kappa, \lambda]$ -compact if the space $E_\tau(L)$ is $[\kappa, \lambda]$ -compact for any τ in the domain of L . In terms of theories this is expressed: L is $[\kappa, \lambda]$ -compact if whenever $\{T_\alpha\}_{\alpha < \kappa}$ is a family of theories such that $\bigcup_{\beta < \delta} T_{\alpha_\beta}$ is satisfiable for any $\delta < \kappa$, then $\bigcup_{\alpha < \kappa} T_\alpha$ is satisfiable.

Of course, all the results about compactness of the previous sections apply to spaces of models.

Notice that the definition of $[\kappa, \lambda]$ -compactness does not put cardinality condition in the theories T_α . On the contrary, the following equivalent characterization makes reference to the cardinality of theories. We do not know if some form of it makes sense and holds for topological spaces in general. The proof of the equivalence seems to use essential model theoretic facts (although in a very simple fashion), see [M] or [Ma] for a proof.

LEMMA 2.2. L is $[\kappa, \lambda]$ -compact if whenever T, T' are theories in L such that $|T'| = \kappa$ and $T \cup S$ is satisfiable for a subset S of T' of power less than λ , then the theory $T \cup T'$ is satisfiable (moreover, we need this to hold for first order T' only!).

A weaker notion of compactness has been very much utilized in abstract model theory, L is (κ, λ) -compact if whenever any subset of less than λ sentences of a set of κ sentences has a model then the full set has a model. This amounts to state compactness for families of basic closed classes only. (∞, λ) -compactness of a logic L is equivalent to $[\infty, \lambda]$ -compactness, because the λ -intersection property for a family $\{F_i\}$ of closed classes implies the λ -intersection property for the associated family of closed basics generating the F_i . However, for $\kappa < \infty$, (κ, λ) -compactness does not imply $[\kappa, \lambda]$ -compactness, since whenever $|L(\tau)| > \kappa$ there may be closed classes which are the intersection of more than κ basics. In particular, what is called countable compactness in logic, this is (ω, ω) -compactness, does not correspond to topological countable compactness. For example, the logic $L_{\omega\omega}(Q_1)$ is (ω, ω) -compact but not $[\omega, \omega]$ -compact. Of course, both notions coincide for $|L(\tau)| = \kappa$.

Notice a curious connection between both notions: L is $[\kappa, \lambda]$ -compact if and only if any closed subclass of $E_\tau(L)$ is (κ, λ) -compact with respect to the topology inherited from $E_\tau(L_{\omega\omega})$. This is just a reformulation of Lemma 2.1 above.

The following is a very useful characterization of $[\kappa, \kappa]$ -compactness for regular logics, noticed first by Väänänen [V] and implicit in Lindström [L] for the countable case (see Th. 1.2.2 in [Ma]).

THEOREM 2.3. *The following are equivalent for a regular logic L and a regular cardinal κ :*

- i) L is $[\kappa, \kappa]$ -compact.
- ii) Any structure $\mathcal{M} = (M, <, \dots)$ where $<$ is a linear order of M of cofinality κ , has an elementary extension $\mathcal{M}' = (M', <', \dots)$ such that M is not cofinal in $(M', <')$.

Proof. (i) \Rightarrow (ii). Assume L is $[\kappa, \kappa]$ -compact. Given $\mathcal{M} = (A, <, \dots)$ as in (ii), let $(a_\alpha)_{\alpha < \kappa}$ be a strictly increasing cofinal sequence in $(A, <)$ and

$$T = \text{Th}_L[(A, <, \dots, a_\alpha, a)_{a \in A, \alpha < \kappa}].$$

If c is a new constant, then each theory $T_\delta = T \cup \{c > a_\alpha \mid \alpha < \delta\}$, $\delta < \kappa$, has a model (interpret c by a_δ). The family $\{\text{Mod}(T_\alpha)\}_{\alpha < \mu}$ forms a descending chain of closed classes; by regularity of κ and Lemma 1.3(c), the intersection of the family is non-empty, yielding a model of $T \cup \{c > a_\beta \mid \beta < \kappa\}$ that is the desired extension.

(ii) \Rightarrow (i) if L is not $[\kappa, \kappa]$ -compact, it is not chain-compact. Let $\{T_\alpha \mid \alpha < \kappa\}$ be a family of theories in $L(\tau)$ such that each $\bigcup_{\alpha < \gamma} T_\alpha$ has a model A_γ for $\gamma < \kappa$, but $\bigcup_{\alpha < \kappa} T_\alpha$ does not. Choose an ordering $<_\gamma$ of each A_γ with first element a_γ and let

$$\mathcal{M} = (\bigcup_{\gamma < \kappa} A_\gamma, <, \bigcup_{\gamma < \kappa} R^A_\gamma, \dots, S)(R \in \tau)$$

where $<$ is the ordered union of the $<_\gamma$ and

$$S = \bigcup_{\gamma < \kappa} (A_\gamma \times A_\gamma).$$

Evidently $(a_\gamma)_{\gamma < \kappa}$ is cofinal in $(M, <)$. Moreover, if $x \geq a_\gamma$ then $\mathcal{M} \models \{y : S(y, x)\} \upharpoonright \tau \approx A_{\gamma'}$, with $\gamma' > \gamma$ and so:

$$\mathcal{M} \models \forall x (x \geq a_\gamma \rightarrow \varphi\{y : S(y, x)\}) \quad (1)$$

for any $\varphi \in T_\gamma$. If $\mathcal{M}' > \mathcal{M}$, \mathcal{M}' must satisfy these sentences. Hence, it can not contain b such that $b > a_\gamma$ for all $\gamma < \kappa$, because then we would have by (1):

$$\mathcal{M}' \models \varphi \{y \mid S(y,b)\}$$

and so $\mathcal{M}' \models \{y : S(y,b)\} \upharpoonright \tau$ would be a model of all $\varphi \in \bigcup_{\gamma < \kappa} T_\gamma$. We conclude that $(a_\gamma)_{\gamma < \kappa}$ and so M , is cofinal in $(M', <')$. \square

COROLLARY 2.4. *The following are equivalent for a regular logic L :*

- i) L is compact.
- ii) Any infinite structure $\mathcal{M} = (M, <, \dots)$ where $<$ is a linear order of M has an elementary extension $\mathcal{M}' = (M', <', \dots)$ such that M is not cofinal in $(M', <')$.

COROLLARY 2.5 L is $[\omega, \omega]$ compact if and only if $(\omega, <)$ is not RPC_δ characterizable in L .

Proof. " \Rightarrow " obvious. " \Leftarrow " If L is not $[\omega, \omega]$ -compact then there is $\mathcal{M} = (M, <, \dots)$ of cofinality ω with $(M, <)$ cofinal in all its L -elementary extensions. Take a strictly increasing cofinal sequence $(a_n)_{n \in \omega}$ of $(M, <)$. The theory:

$$\begin{aligned} \text{Th}_L[(M, <, \dots, a, a_n)_{a \in A, n \in \omega}] \cup \{ \forall x \exists y (P(y) \wedge x < y) \} \cup \{ \neg \exists x (P(x) \wedge x < c_0) \} \\ \cup \{ P(c_n) \wedge \neg \exists x (P(x) \wedge c_n < x < c_{n+1}) : n \in \omega \} \end{aligned}$$

gives a RPC_δ characterization of $(\omega, <)$, since the interpretation of P in any model must consist of the a_n alone. \square

It follows from the previous corollary that the failure of $[\omega, \omega]$ -compactness for L implies that $(\omega, <)$ is RPC_δ characterizable in L . In fact any structure (of power below the first measurable cardinal if there is any) is RPC_δ -characterizable in L under failure of countable compactness, as we will see next. This may be shown via Theorem 2.3, proving first that any ordinal below the first measurable is RPC_δ , but the following reformulation of the Rabin-Keisler theorem allows a more direct and elegant proof.

Definition 2.6. A structure A will be said to be L -full (also complete) if any relation $R \subseteq A^n$ is the interpretation of a predicate in the vocabulary of A .

Given an full structure and an extension $A <_L A^*$, we will denote by R^* the canonical extension in A^* of any relation R in A (that is, if R is the interpretation of P_R in A , then R^* is the interpretation of P_R in A^*). Notice that if $a \in A^m$ and $R = \{x \in A^n : A \models \phi(x, a)\}$ with $\phi \in L$, then $R^* = \{x \in (A^*)^n : A^* \models \phi(x, a)\}$, because the sentence $\forall x (P_R(x) \leftrightarrow \phi(x, a))$ holds in A if and only it holds in A^* . Hence, any relation holding between the R 's, expressible in L holding in A , holds for the corresponding R^* 's in A^* . The following is a reformulation of the Rabin-Keisler theorem [BS], [Ch-K].

LEMMA 2.7. *Let A be a L -full structure, $A <_{L_{\omega\omega}} A^*$, $P \subseteq A$, and $b \in P^* - P$. Define $U_b = \{S \subseteq P : b \in S^*\}$.*

- i) U_b is a non principal ultrafilter over P .
- ii) If $|P| <$ first measurable cardinal then any infinite $Q \subseteq A$ has $Q^* - Q$ no empty.

Proof. i) To show that it is an ultrafilter, utilize that the extension is full and the previous remarks; for example, $S, T \in U_b \Rightarrow b \in S^*, T^* \Rightarrow b \in S^* \cap T^* = \{x \in A^* : P_S(x) \wedge P_T(x)\} = \{x \in A^* : P_{S \cap T}(x)\}^* = (S \cap T)^*$, where the last equation holds because the sentence $\forall x ((P_S(x) \wedge P_T(x) \Rightarrow P_{S \cap T}(x))$ holds in A and so in A^* . Now, if the ultrafilter were principal, say $U_b = (\{a\})$ with $a \in A$, then we would have $b \in \{a\}^* = \{a\}$ and so $b = a \in P$.

ii) It is enough to show that any countably infinite subset Q of A is extended. Suppose Q is not extended. Given a countable family of elements of U_b , utilize Q as subindex set: $\{S_q : q \in Q\}$. Let $S = \bigcap_q S_q$ and $F(q, x)$ be the binary relation $x \in S_q$ in A . The following sentences hold trivially in A where we use the names of the sets as predicates:

$$\begin{aligned} \forall x [S_q(x) \rightarrow F(q, x)] & \quad \text{for each } q \in Q; \\ \forall x [\forall q (Q(q) \rightarrow F(q, x)) \rightarrow S(x)] \end{aligned}$$

hence, it holds for b in A^* that:

$$\begin{aligned} {}^*S_q(b) \rightarrow {}^*F(q, b) & \quad \text{for each } q \in Q \\ \forall q ({}^*Q(q) \rightarrow {}^*F(q, b)) \rightarrow {}^*S(b) \end{aligned}$$

Since ${}^*S_q(b)$ holds by hypothesis for $q \in Q^*$ then ${}^*F(q, b)$ holds in A^* for all $q \in Q$. As $Q = Q^*$ also by hypothesis then the antecedent of the last implication holds and so ${}^*S(b)$ holds, showing $S \in U_b$. We conclude that U_b is an ω -complete ultrafilter. But it is well known that as set carrying an ω -complete no principal ultrafilter must have power at least the first measurable cardinal [BS]. \square

COROLLARY 2.8. *If L is not $[\omega, \omega]$ -compact then any structure of power less than the first measurable is RPC_δ -characterizable in L .*

Proof. Let $T(Q, R)$ be a theory such that $B \models T \Leftrightarrow B|Q^B| \{R\} \approx (\omega, <)$, and fix a model B of T . Let C be the full expansion of $[B, A]$ where A has power smaller than the first measurable, and call P be the universe of A in C . We may assume the vocabularies of T and A are disjoint. Any L -extension of C contains an L -extension of B which must be a model of T (thanks to relativizations), and so $(Q^B, <) = (\omega, <)$ can not be extended in C by hypothesis. By the previous lemma P can not be extended either. This means that any model of $Th_L((C, c)_{c \in |C|})$ relativized to P is isomorphic to A , yielding a RPC_δ characterization of A in L . \square

The expressive power of theories in a non-countably compact logic is therefore quite strong.

COROLLARY 2.9. (Makowsky and Shelah, [Ma-Sh]). *If κ is a regular cardinal smaller than the first measurable cardinal (or arbitrary if there are not such cardinals), then $[\kappa, \kappa]$ -compactness of a logic implies $[\omega, \omega]$ -compactness.*

Proof. If L is not $[\omega, \omega]$ -compact, κ is RPC_δ -characterizable which contradicts $[\kappa, \kappa]$ -compactness. \square

Remark. If we had the stronger hypothesis in Corollary 2.8 that $(\omega, <)$ is PC_δ characterizable, then any structure (of power $<$ first measurable) would be PC_δ characterizable, by the L -theory of its full expansion (since in the infinite case it would contain a copy of a structure capturing ω). This is the case for example of $L_{\omega\omega}(Q_0)$. If $(\omega, <)$ has a characterization with models of power less than the first measurable, then any structure above certain cardinality is also PC_δ . Could it happen that any theory RPC_δ characterizing $(\omega, <)$ in L only has models of cardinality above the first measurable? In that case there would not be counterexample to $[\omega, \omega]$ -compactness in the form: if each finite subfamily of theories has models of power less than the first measurable, then it has models.

III. COMPACTNESS AND ROBINSON'S LEMMA

It is well known that in the presence of compactness interpolation implies Robinson's lemma for first order logic, this being true for any logic. Mundici and Makowsky discovered independently the remarkable result that in any logic with small dependence number, Robinson's theorem implies compactness, yielding the equations Robinson = Compactness + Interpolation.

Definition 3.1. A logic L satisfies the *Robinson property* if given any pair of vocabularies and structures $\sigma_1, A_1, i = 1, 2$, with $\mu = \sigma_1 \cap \sigma_2$ such that $A_1|_\mu \equiv_L A_2|_\mu$, there is a third structure B such that $A_1 \equiv_L B|_{\sigma_1}$ and $A_2 \equiv_L B|_{\sigma_2}$.

Of course, we may strength \equiv_L to $<_L$ above. In terms of theories the property may be expressed: if we have satisfiable theories $T_1 \subseteq L(\sigma_1)$, $T_1, T_2 \supseteq T$ with T complete in $L(\tau_1 \cap \tau_2)$ then $T_1 \cup T_2$ is satisfiable. The following is an equivalent characterization of Robinson's property which follows also by definition.

LEMMA 3.2. L satisfies the Robinson property if and only if any pair of PC_δ classes K_1, K_2 of L having structures $A_i \in K_i$ such that $A_1 \equiv_L A_2$, must have non-empty intersection.

THEOREM 3.3 (see Mundici, [Mu]). For logics with finite dependence, Robinson's property implies compactness. In general, for small logics with dependence number at most κ , Robinson's property implies $[\infty, \kappa]$ -compactness.

Proof. Suppose L is not $[\mu, \mu]$ compact for some regular $\mu \geq \kappa$ and let $\mathcal{A} = (A, < \dots)$ a expansion of a linear order of cofinality μ , having all its elementary extensions cofinal. Let $(a_\alpha)_{\alpha < \mu}$ be a well ordered cofinal sequence in $(A, <)$, introduce predicates $P_\alpha = \{x \in A \mid x \leq a_\alpha\}$ and let:

$$\mathcal{A}_1 = (A, <, \dots, a_\alpha, P_\alpha)_{\alpha < \mu}$$

$$\mathcal{A}_2 = (A \cup \{c\}, c, P_\alpha)_{\alpha < \mu}$$

where $c \notin A$. Let $\tau_1 = \tau(\mathcal{A}_1)$, $\tau_2 = \tau(\mathcal{A}_2)$ and $\tau = \tau_1 \cap \tau_2 = \{P_\alpha\}_\alpha$ then for $\delta < \kappa \leq \mu$ and $\tau_\delta = \{P_\alpha : \alpha < \delta\}$ we have

$$\mathcal{A}_1 \upharpoonright \tau_\delta = (A, P_\alpha)_{\alpha < \delta} \approx (A \cup \{c\}, P_\alpha)_{\alpha < \delta} = \mathcal{A}_2 \upharpoonright \tau_\delta.$$

As any sentence of L depends in less than κ symbols, then $\mathcal{M}_1 \upharpoonright \tau \equiv_L \mathcal{M}_2 \upharpoonright \tau$ by the isomorphism axiom of logics. By Robinson's theorem there is

$$\mathcal{M}' = (\mathcal{A}', <', \dots, c', a'_\alpha, P'_\alpha)_{\alpha < \mu}$$

of type $\tau_1 \cup \tau_2$ such that $\mathcal{M}' \upharpoonright \tau_1 \equiv_L \mathcal{M}_1$, $\mathcal{M}' \upharpoonright \tau_2 \equiv \mathcal{M}_2$. But

$$\mathcal{M}_1 \models \forall x [\neg P_\alpha(x) \rightarrow x > a_\alpha], \text{ for all } \alpha < \mu,$$

$$\mathcal{M}_2 \models \neg P_\alpha(c),$$

hence $\mathcal{M}' \models c' > a'_\alpha$ for all $\alpha < \mu$, and so $\mathcal{M}' \upharpoonright \tau(\mathcal{M})$ gives a non cofinal elementary extension of \mathcal{M} , a contradiction. \square

The finite dependence hypothesis of the first part of the above theorem may be weakened, thanks to a strong result due to Shelah [M-Sh] that we will not prove. It says that a $[\omega, \omega]$ -compact logic has finite dependence for each sentence (notice that this is easy if the logic is fully compact, but it is a deep theorem if we have only $[\omega, \omega]$ -compactness).

COROLLARY 3.4. *For small logics with dependence number smaller than the first measurable cardinal, we have: Robinson's lemma \Rightarrow compactness.*

Proof. Let μ be the dependence number of L and let κ be regular such that $\mu < \kappa < \text{first measurable}$. By Theorem 3.3, L is $[\kappa, \kappa]$ -compact. By Corollary 2.7 it must be $[\omega, \omega]$ -compact. By the finite dependence theorem of Shelah (Th. 2.2.1. in [Ma]), L has finite dependence. Hence, by Theorem 3.3 again we have full compactness. \square

For the next theorem we need a topological observation first. Define in a uniform space $x \equiv y$ if $(x, y) \in U$ for any element U of the uniformity.

LEMMA 3.5. *Let M and N be disjoint compact sets in a uniform space inseparable by finite unions of basic open sets (from any given basis), then there exists $x \in N, y \in M$ such that $x \equiv y$.*

Proof. Let $E = \{ (x, y) \in X \times X : x \equiv y \}$, then $E \cap N \times M$ is the intersection of the closed sets $C_{p,n} = \{ (x, y) \in N \times M : p(x, y) \leq 1/n \}$ of $M \times N$, where p runs through the system of pseudometrics defining the uniformity. Now, for each pseudometric p and number $n \in \omega$, the coverings $\{ V_{p, 1/2n}(x) : x \in N \}$ and $\{ V_{p, 1/2n}(y) : y \in M \}$, must have some $x \in N, y \in M$ with $V_{p, 1/2n}(x) \cap V_{p, 1/2n}(y)$ non empty. Otherwise, N and M would be separable by

opens, which could be reduced to finite unions of basics by compactness. Therefore $p(x,y) \leq 2(1/2n) = 1/n$. Since the pseudometrics are directed by \leq (see []), this shows that $E \cap N \times M$ has the f.i.p. By compactness of the product topology we have that E is non empty.

Remark. The above lemma holds for regular spaces.

COROLLARY 3.6 . *The following are equivalent if L is small and has dependence number less than the first measurable cardinal:*

- i) Robinson's property,
- ii) Int_δ (interpolation of disjoint PC_δ classes),
- iii) Interpolation + Compactness.

Proof. (i) \Rightarrow (ii, iii). Assume Robinson's property, then we have compactness by Theorem 3.3. Let K_1, K_2 be disjoint PC_δ classes in $E_\mu(L)$ and assume they are not separable by a sentence. Since the K_i are compact, the Lemma above implies the existence of $A_i \in K_i$ with $A_1 \equiv_L A_2$. Then Robinson's lemma provides a structure in the intersection of this two classes by Lemma 3.1.

(ii) \Rightarrow (i). Robinson's property is just a case of interpolation of PC_δ . If K_i are PC_δ classes of L as in Lemma 3.1, $i = 1, 2$, they must intersect, otherwise by Int_δ they would be separable by a sentence ϕ contradicting that $A_1 \equiv_L A_2$.

(iii) \Rightarrow (ii). Trivial, because disjoint PC_δ classes are reduced by compactness to disjoint PC classes. \square

Finally we relate Robinson's property to characterization of structures.

LEMMA 3.7. *Let L satisfy Robinson's property, if A and B are structures PC_δ -characterizable in L then $A \equiv_L B$ implies $A \approx B$.*

Proof. Assume $T_i \subseteq L(\mu_i)$ is the theory characterizing A_i , $\sigma \subseteq \mu_i$, and let (A_i, R_i) be model of T_i , $i = 1, 2$. Assume $A_1 \equiv_L A_2$, then by Robinson's property there is a model (A, R_1^*, R_2^*) with $(A, R_1^*) \equiv_L (A_1, R_1)$; hence, $A \approx A_1 \approx A_2$. \square

If the logic satisfies the pair preservation property, PPP (cf. [Ma]), then the Robinson's property implies the following stronger property (see [C2]):

Definition 3.8. A logic L satisfies the *relativized Robinson property* if given any pair of vocabularies $\sigma_1, P_1 \in \sigma_1$, and structures A_i , $i = 1, 2$, with $\mu = \sigma_1 \cap \sigma_2$ such that $A_1|P_1^{A_1}|_\mu \equiv_L A_2|P_2^{A_2}|_\mu$, there is a third structure B such that $A_1 \equiv_L B|_{\sigma_1}$ and $A_2 \equiv_L B|_{\sigma_2}$.

Therefore, under PPP, Lemma 3.7 holds for structures RPC_δ -characterizables in L . There is some confusion in the literature on relation to Robinson's property. What is usually called Robinson's property in the context of many sorted logics is in fact equivalent to the above stronger property.

COROLLARY 3.9. (compare with Mundici [Mu]). *If L satisfies PPP and Robinson's theorem but is not $[\omega, \omega]$ -compact then for structures of power less than the first non measurable cardinal, $A \equiv_L B$ implies $A \approx B$.*

Proof. If the logic is not $[\omega, \omega]$ -compact, then any structure of power less than the first measurable is RPC_δ in L by Corollary 2.10. Apply the previous lemma and observations. \square

IV. $[\kappa, \lambda]$ -COMPACTNESS AND ULTRAFILTER CONVERGENCE

It is well known full compactness of a topological space is equivalent to the convergence of any ultrafilter of subsets of the space (cf. [W]). This in turn may be expressed in terms of convergence of I-families of elements of the space with respect to ultrafilters over I. It is natural to ask if we have characterizations via ultrafilters of $[\kappa, \lambda]$ -compactness. In certain sense this is true as we show in this section. In fact, we get more than we sought: a characterization by ultrafilter convergence of families of $[\kappa, \lambda]$ -compact spaces closed under products, and of spaces with $[\kappa, \lambda]$ -compact powers.

Definition 4.1. Let U be an ultrafilter over a set I . We will say that a family $\{a_i : i \in I\}$ in a topological space X , U -converges to x (or that x is an U -limit of the family), if $\{i \in I : a_i \in V\} \in U$ for any open neighborhood V of x .

Evidently, $\{a_i : i \in I\}$ U -converges to x if and only if x is an adherence point in X of the ultrafilter $\mathfrak{a}(U) = \{S \subseteq X : \{i \in I : a_i \in S\} \in U\}$ in the ordinary sense of topology (cf. [W]). Therefore, X is compact if and only if any I-family U -converges in X for any set I and ultrafilter U over I . We show next that $[\kappa, \lambda]$ -compactness corresponds to U -convergency with respect to certain ultrafilters.

Definition 4.2. Let U be an ultrafilter over a set I , a space X will be called U -compact if and only if any I-family of X U -converges.

The following are obvious properties of U -convergence and U -compactness:

LEMMA 4.3. i) If $f: X \rightarrow Y$ is continuous and $\{a_i : i \in I\}$ U -converges in X to a , then $\{f(a_i) : i \in I\}$ U -converges in Y to $f(a)$.

ii) $\{(a_{i,\alpha})_\alpha : i \in I\}$ U -converges in $\prod_\alpha X_\alpha$ to $(a_\alpha)_\alpha$ if and only if $\{a_{i,\alpha} : i \in I\}$ U -converges in X_α to a_α , for each α .

iii) If each X_α is U -compact, then $\prod_\alpha X_\alpha$ is U -compact.

Proof. i) Given a neighborhood V of $f(a)$, then $\{i \in I : f(a_i) \in V\} \supseteq \{i \in I : a_i \in f^{-1}(V)\} \in U$.

ii) One direction follows from (i) by continuity of the projections. For the other, use that each basic neighborhood W of $(a_\alpha)_\alpha$ is a finite intersection of sets $\pi_\alpha^{-1}(V_\alpha)$ with V_α a neighborhood of a_α in X_α . For each such set $\{i \in I : a_{i,\alpha} \in \pi_\alpha^{-1}(V_\alpha)\} \in U$; hence, $\{i \in I : (a_{i,\alpha})_\alpha \in W\} = \{i \in I : a_{i,\alpha} \in \pi_\alpha^{-1}(V_\alpha)\} = \cap \{i \in I : a_{i,\alpha} \in \pi_\alpha^{-1}(V_\alpha)\} \in U$.

iii) Immediate from (ii). \square

Definition 4.4. An ultrafilter U over a set I is (κ, λ) -regular if and only if there is a family $\{I_\alpha\}_{\alpha < \kappa}$ such that $I_\alpha \in U$ and $\bigcap_{i < \lambda} I_{\alpha_i} = \emptyset$ for any family $\{\alpha_i : i < \lambda\} \subseteq \kappa$. U is κ -uniform if $|S| \geq \kappa$ for any $S \in U$, see [Ch-K] or [Ma].

PROPOSITION 4.5. If X is U -compact for a (κ, λ) -regular ultrafilter U , then X is $[\kappa, \lambda]$ -compact.

Proof. Let $\{I_\alpha\}_{\alpha < \kappa}$ be a family of elements of U such that the intersection of any λ of I_α 's is empty. We may assume $I = I_0$. Given a family $\{F_\alpha\}_{\alpha < \kappa}$ of closed sets in X with the λ -intersection property, define $F_t = \bigcap_{\alpha \in I_\alpha} F_\alpha$ for each $t \in I$. This set is non-empty, because t belongs to less than λ sets I_α by hypothesis, and by hypothesis the intersection of less than λ sets F_α is non-empty. Choose $a_t \in F_t$, then $J_\alpha = \{t \in I : a_t \in F_\alpha\} \in U$ because $t \in I_\alpha$ implies $a_t \in F_t \subseteq F_\alpha$ by construction, and so $J_\alpha \supseteq I_\alpha$. By hypothesis, $\{a_t\}_{t \in I}$ U -converges to some x of X ; and given an open neighborhood V of x , then $J = \{t \in I : a_t \in V\} \in U$. Therefore, $\{t : a_t \in V \cap F_\alpha\} = J \cap J_\alpha \in U$ for any α , and so this set is non-empty, showing that x belongs to adherence of any F_α , that is to any F_α . \square

Definition 4.6. Let $P(\kappa, \lambda) = P_\lambda(\kappa) = \{S \subseteq \kappa : |S| < \lambda\}$.

PROPOSITION 4.7. i) If X is $[\kappa, \lambda]$ -compact, then every I -family in X , with $I = P(\kappa, \lambda)$, U -converges for some (κ, λ) -regular ultrafilter U over I (which may depend on the family).

ii) Moreover, if $\kappa > \lambda$ then U may be chosen κ -uniform, and if $\kappa = \lambda$ then U may be chosen $\text{cof}(\kappa)$ -uniform.

Proof. Given $\{a_t : t \in I\}$, let $A_t = \{a_s : t \subseteq s\}$. The family of closed sets $\{\text{cl}(A_t) : t \in I, |t| < \omega\}$ has the $< \lambda$ -intersection intersection property, because

$$\bigcap_{i < \delta} \text{cl}(A_{t_i}) \supseteq \text{cl}(\bigcap_{i < \delta} A_{t_i}) = \text{cl}(A_{\bigcup_{i < \delta} t_i}),$$

and $t = \bigcup_{i < \delta} t_i \in I$ if $\delta < \lambda$ and the t_i are finite, so that A_t is non-empty. By $[\kappa, \lambda]$ -compactness there is an element $a \in \bigcap_{t \in I, |t| < \omega} \text{cl}(A_t)$. Hence, $V \cap A_t \neq \emptyset$ for any neighborhood V of a and any $t \in I, |t| < \omega$. This implies that the family $F = \{V \cap A_t : V \text{ open}, a \in V, t \in I, |t| < \omega\}$ has the finite intersection property. Then the family $F' = \{a^{-1}(W) : W \in F\} = \{s \in I : a_s \in W\} : W \in F\}$ has also the finite intersection property. Extend F' to an ultrafilter U over I . By construction, given an open neighborhood V of a , then $\{s \in I : a_s \in V\} \supseteq \{s \in I : a_s \in V \cap A_{\{\alpha\}}\} = a^{-1}(V \cap A_{\{\alpha\}}) \in F' \subseteq U$. Therefore $\{a_t : t \in I\}$ U -converges to a . Moreover, $I_\alpha = \{s \in I : \alpha \in s\} \supseteq \{s \in I : a_s \in V \cap A_{\{\alpha\}}\}$, and so

$I_\alpha \in U$. But the intersection of λ -many of these I_α 's is empty because no $s \in I$ may contain λ many ordinals. This shows that U is (κ, λ) regular.

It remains to show that U may be chosen to be κ -uniform (respectively $\text{cof}(\kappa)$ -uniform). Fix $t_0 \in P(\omega, \kappa)$. For each finite $t \subseteq \kappa$ with $t \supseteq t_0$ the set $\{s \supseteq t : a_s \in V\} = a^{-1}(V \cap A_t)$ is in U and so we may pick $s_t \in a^{-1}(V \cap A_t)$; hence $s_t \supseteq t$. This gives a function $f: B = \{t \in P(\omega, \kappa) : t \supseteq t_0\} \rightarrow I$ with the property that for any $r \in I$, $|f^{-1}(r)| = |\{t \in B : s_t = r\}| \leq |\{t \in B : t \subseteq r\}| < \lambda$, because $|r| < \lambda$ and so r has less than λ finite subsets. Since $|B| = \kappa$, the number of non-empty distinct and so disjoint $f^{-1}(r)$ must be at least $\text{cof}(\kappa)$ in case $\kappa = \lambda$, and κ in case $\kappa > \lambda$. Picking one s_t from each non-empty $f^{-1}(r)$ gives κ (respectively $\text{cof}(\kappa)$) elements of $a^{-1}(V \cap A_{t_0})$. Therefore, F' may be extended to a κ -uniform (respectively $\text{cof}(\kappa)$ -uniform) ultrafilter. \square

COROLLARY 4.8. i) If X is (κ, λ) -compact and $\kappa > \lambda$ then any $P(\kappa, \lambda)$ -family of X has a κ -accumulation point.

ii) If X is (κ, κ) -compact then any $P(\kappa, \kappa)$ -family of X has a $\text{cof}(\kappa)$ -accumulation point.

Proposition 4.7 does not guarantee a converse to Proposition 4.6, since the ultrafilter depends on the family. In fact, a converse is impossible because U -compactness corresponds to a strong form of (κ, λ) -compactness, as we will see next.

THEOREM 4.9. The following are equivalent for any class T of topological spaces:

- i) Any product of spaces in T is (κ, λ) -compact.
- ii) There exists a (κ, λ) -regular ultrafilter U (which may be taken over $I = P(\kappa, \lambda)$) such that all the spaces in T are U -compact.

Proof. Assume that any product of space in T is (κ, λ) -compact and there is no (κ, λ) -regular ultrafilter U over $I = P(\kappa, \lambda)$ such that all the elements of T are U -compact. Let Σ be the family of such ultrafilters and choose for each $U \in \Sigma$ a I -family $\{a_{U,i} : i \in I\}$ in some space $X_U \in T$ which does not U -converge. For each i , let $\sigma_i = (a_{U,i})_U \in \prod_{U \in \Sigma} X_U = X^*$. As this space is (κ, λ) -compact, then by Proposition 4.8, there is an ultrafilter $W \in \Sigma$ such that $\{\sigma_i : i \in I\}$ W -converges to some $\sigma = (a_U)_U \in X^*$. By continuity of the W -projection, then $\{a_{W,i} : i \in I\}$ W -converges to a_W in X_W , a contradiction.

Conversely, if each X_T is U -compact then $\prod_T X_T$ is U -compact by Lemma 4.3, and by (κ, λ) -regularity of U it follows from Proposition 4.6 that $\prod_T X_T$ is (κ, λ) -compact. \square

THEOREM 4.10. *The following are equivalent for any topological space X :*

- i) X^β is $[\kappa, \lambda]$ -compact for any cardinal β .
- ii) X is U -compact for some (κ, λ) -regular ultrafilter U (which may be taken over $I = P(\kappa, \lambda)$).

Proof. Take $T = \{ X \}$ in Theorem 1. \square

Definition 4.11. Call a space X *strongly $[\kappa, \lambda]$ -compact* (in short, *s- $[\kappa, \lambda]$ -compact*) if X^β is $[\kappa, \lambda]$ -compact for any cardinal β .

COROLLARY 4.12. X is s- $[\kappa, \lambda]$ -compact if and only if $X^{2^{2^{|P(\kappa, \lambda)|}}}$ is $[\kappa, \lambda]$ -compact.

Proof. In the proof of (i) \Rightarrow (ii) in Theorem 4.9 we utilized $[\kappa, \lambda]$ -compactness of X^β for $\beta = 2^{2^{|P(\kappa, \lambda)|}}$ only. \square

COROLLARY 4.13. If X is s- $[\kappa, \lambda]$ -compact, then $X \times Y$ is $[\kappa, \lambda]$ -compact for any compact space Y .

Proof. A compact space is U -compact for any U . Apply Theorem 4.10 and Lemma 4.3. \square

Example. Under the continuum hypothesis, plus $\omega_2 = 2^{\omega_1}$, if X^{ω_2} is countably compact, then $X^\beta \times Y$ will be countably compact for any β , and any compact space Y .

COROLLARY 4.14. Let κ be smaller than the first measurable cardinal (or arbitrary if no such cardinal exists). If X is s- $[\kappa, \kappa]$ -compact, then X is $[\omega, \omega]$ -compact.

Proof. Let U be the ultrafilter given by Theorem 4.10, such that X is U -compact. By (κ, κ) -regularity, U is non principal. If U is not (ω, ω) -regular, then it is ω -complete. But it is well known that the smallest ω -complete no principal ultrafilter is measurable (see [BS]). \square

Notice that the above fails strongly for plain compactness, $X = \omega$ with the discrete topology is $[\kappa, \kappa]$ compact for any $\kappa \geq \omega_1$ and however it is not $[\omega, \omega]$ -compact.

For $[\kappa, \kappa]$ -compactness with κ regular, theorems 4.9 and 4.10 take a more beautiful form. The condition on (κ, λ) -regularity of the ultrafilter U may be changed to κ -uniformity. We state the case of one single space.

THEOREM 4.15. Let κ be a regular cardinal then X is s- $[\kappa, \kappa]$ -compact if and only if X is U -compact for some κ -uniform ultrafilter U .

Proof. A κ -uniform ultrafilter is (κ, κ) -regular. On the other hand, Proposition 4.8 allows to choose the ultrafilter U being $\text{cof}(\kappa)$ -uniform. But for κ regular, $\text{cof}(\kappa) = \kappa$. \square

Of course, the ultrafilter U may be chosen over $P_\kappa(\kappa)$. Could it be chosen over κ ?

COROLLARY 4.16. *If X is $s\text{-}[\kappa^+, \kappa^+]$ -compact, then X is $s\text{-}[\kappa, \kappa]$ -compact.*

Proof. A κ^+ -uniform ultrafilter is (κ, κ) -regular by results of Kanamori [K] and Kunen-Pikry [K-P]. \square

COROLLARY 4.17. *Let κ be a regular cardinal or ∞ , then X is $s\text{-}[\kappa, \lambda]$ -compact if and only if X is $s\text{-}[\mu, \mu]$ -compact for any regular cardinal μ , $\lambda \leq \mu \leq \kappa$.*

Proof. Let μ , $\lambda \leq \mu \leq \kappa$, be singular, then $\lambda \leq \mu^+ \leq \kappa$. Apply then the previous corollary. \square

The previous results do not hold for plain $[\kappa, \lambda]$ -compactness, $X = \aleph_\omega$ discrete is $[\infty, \aleph_\omega^+]$ -compact but not $[\aleph_\omega, \aleph_\omega]$ -compact.

V. THE ABSTRACT COMPACTNESS THEOREM REVISITED

Let us apply the results of the previous section to spaces of structures. We start with the following simple observation which follows from the zero-dimensionality of spaces of structures.

FACT 5.1. $\{A_i : i \in I\}$ U -converges to A^* in $E_\sigma(L)$ if and only if for any sentence $\phi \in L(\sigma)$ we have: $A^* \models \phi \Leftrightarrow \{i \in P : A_i \models \phi\} \in U$.

If we use " \Rightarrow " instead of " \Leftrightarrow ", the last equivalence, this is just the definition of U -convergence expressed in terms of a basis, because the basic neighborhoods are the classes $\text{Mod}(\phi)$. The arrow becomes a double arrow since by applying it to the negation of ϕ and so: $A^* \not\models \phi \Rightarrow \{i \in P : A_i \models -\phi\} \in U \Rightarrow \{i \in P : A_i \models \phi\} \notin U$.

Let us recall the following definition from [Ma-Sh].

Definition 5.2. An ultrafilter U over a set I is related to a logic L if and only if for any structure A of type σ there is an extension A^* of the ultrapower A^I/U satisfying for any formula $\phi(x, \dots) \in L(\sigma)$ and functions $f, \dots \in A^I$:

Of course, the ultrafilter U may be chosen over $P_\kappa(\kappa)$. Could it be chosen over κ ?

COROLLARY 4.16. *If X is $s\text{-}[\kappa^+, \kappa^+]$ -compact, then X is $s\text{-}[\kappa, \kappa]$ -compact.*

Proof. A κ^+ -uniform ultrafilter is (κ, κ) -regular by results of Kanamori [K] and Kunen-Pikry [K-P]. \square

COROLLARY 4.17. *Let κ be a regular cardinal or ∞ , then X is $s\text{-}[\kappa, \lambda]$ -compact if and only if X is $s\text{-}[\mu, \mu]$ -compact for any regular cardinal μ , $\lambda \leq \mu \leq \kappa$.*

Proof. Let μ , $\lambda \leq \mu \leq \kappa$, be singular, then $\lambda \leq \mu^+ \leq \kappa$. Apply then the previous corollary. \square

The previous results do not hold for plain $[\kappa, \lambda]$ -compactness, $X = \aleph_\omega$ discrete is $[\infty, \aleph_\omega^+]$ -compact but not $[\aleph_\omega, \aleph_\omega]$ -compact.

V. THE ABSTRACT COMPACTNESS THEOREM REVISITED

Let us apply the results of the previous section to spaces of structures. We start with the following simple observation which follows from the zero-dimensionality of spaces of structures.

FACT 5.1. $\{A_i : i \in I\}$ U -converges to A^* in $E_\sigma(L)$ if and only if for any sentence $\phi \in L(\sigma)$ we have: $A^* \models \phi \Leftrightarrow \{i \in I : A_i \models \phi\} \in U$.

If we use " \Rightarrow " instead of " \Leftrightarrow ", the last equivalence, this is just the definition of U -convergence expressed in terms of a basis, because the basic neighborhoods are the classes $\text{Mod}(\phi)$. The arrow becomes a double arrow since by applying it to the negation of ϕ and so: $A^* \not\models \phi \Rightarrow \{i \in I : A_i \models -\phi\} \in U \Rightarrow \{i \in I : A_i \models \phi\} \notin U$.

Let us recall the following definition from [Ma-Sh].

Definition 5.2. An ultrafilter U over a set I is *related* to a logic L if and only if for any structure A of type σ there is an extension A^* of the ultrapower A^I/U satisfying for any formula $\phi(x, \dots) \in L(\sigma)$ and functions $f, \dots \in A^I$:

$$A^* \models \phi(f/U, \dots) \text{ iff } \{ i \in I : A \models \phi(f(i), \dots) \} \in U. \quad (1)$$

This definition says that A^* behaves as the ultrapower would if Loz theorem were true for L . However, A^* can not be chosen to be the ultrapower itself unless $L = L_{\omega\omega}$. The following fact reduces the relation (1) to pure topology.

Given a structure A , of type σ consider the expanded vocabulary $\sigma_{A,I} = \sigma + \{c_f : f \in A^I\}$, where each c_f is a constant symbol, and define for each fixed $j \in I$ the following expansion of A of type $\sigma_{A,I}$:

$$A^*_j = (A, f(j), \dots)_{f \in A^I}$$

where c_f is interpreted by $f(j)$.

LEMMA 5.3. A^* is an extension of A^I/U satisfying (1) if and only if A^* may be expanded to a U -limit of $\{A^*_j : j \in I\}$.

Proof. (1) implies that $(A^*, f/U, \dots)_{f \in A^I}$ is a U limit of the family by Fact 5.1. Conversely, a U -limit (A^*, a_f, \dots) of this family will have interpretations a_f for the constants c_f , $f \in A^I$. Moreover, for any formula $\phi(x, \dots)$ of type σ : $A^*|_\sigma \models \phi(a_f, \dots)$ if and only if $A^* \models \phi(c_f, \dots)$ if and only if $\{j \in I : A^*_j \models \phi(c_f, \dots)\} \in U$ if and only if $\{i \in I : A \models \phi(f(i), \dots)\} \in U$. Applying this to the atomic formulae, we get an isomorphism $a_f \mapsto f/U$ between the substructure of $A^*|_\sigma$ induced by $\{a_f : f \in A^I\}$ and the ultrapower A^I/U . Via this identification, (1) holds. \square

PROPOSITION 5.4. The following are equivalent for any logic L closed under relativizations, and any ultrafilter U :

- i) U is related to L .
- ii) The spaces $E_\sigma(L)$ are U -compact for any $\sigma \in \text{Dom}(L)$.

Proof. Assume U is related to L . Given a family $\{A_i : i \in I\} \subseteq E_\sigma$ code it in a single structure $A = (\cup_i A_i, \cup_i Q^{A_i}, \dots, I, R)$ where Q runs through the vocabulary σ , and $R = \cup_i \{i\} \times A_i$ so that $A_i = A|_{\{x : R(i,x)\}}$. If $A^I/U \subseteq A^*$ where A^* is given by Def. 5.2, let $g_0(i) = i$ be the identity function and $P^* = \{x \in A^* : A^* \models R(g_0/U, x)\}$. Utilizing that the logic has relativizations and property (1), we obtain for any sentence $\phi \in L(\sigma)$:

$$\begin{aligned} A^*|_{P^*} \models \phi & \text{ iff } A^* \models \phi^{(x : R(g_0/U, x))} \\ & \text{ iff } \{i \in I : A \models \phi^{(x : R(i, x))}\} \in U \end{aligned}$$

$$\text{iff } \{ i \in I : A_i \models \phi \} = \{ i \in I : A_i \models \phi \} \in U.$$

Hence, $\{A_i : i \in I\}$ U-converges to $A^*|P^*$.

Conversely, assume any I-family U-converges in the spaces of structures. Given a structure A, there is an U-limit (A^*, a_f, \dots) of the associated family $\{A^*_j\}$ which will satisfy (1) by Lemma 5.3 showing that U is related to L. \square

Remark. Notice that in the first part of the above proof, the existence of $A(\mu)^*$ satisfying (1) for the complete extension $A(\mu)$ of power $\mu \geq |I|$, will guarantee the U-convergence of all I-families of structures of power at most μ , because they may all be accommodated in A, via appropriate codings and renamings. Therefore, the U-convergence of a single I-family of structures of power μ in each cardinal $\mu \geq |I|$ yields U-compactness.

To apply our topological results, we need a further observation on the spaces of structures. Given vocabularies $\sigma_i, i \in I$, let $\oplus_i \sigma_i$ be the disjoint union of the vocabularies $\sigma_i + \{P_i\}$ where P_i is a monadic symbol not in σ_i . Then the cartesian product $\prod_i E_{\sigma_i}$ may be identified with the class of structures of type $\oplus_i \sigma_i$ of the form:

$$[A_i]_{i \in I} = (\cup_{i \in I} |A_i|, \Delta_i, \dots)_{i \in I}, \quad A_i \in E_{\sigma_i},$$

where the universe is the disjoint union of the universes of the A_i , and the disjoint renaming of $\sigma_i + \{P_i\}$ is interpreted by the disjoint copy Δ_i of A_i , with P_i interpreting the universe (cf. [C1]).

LEMMA 5.4. *If L is closed under relativizations then for any product space $\prod_i E_{\sigma_i}(L)$ there is a uniformly continuous onto operation $F: C \rightarrow \prod_i E_{\sigma_i}(L)$ where C is a closed subclass of $E_{\oplus_i \sigma_i}(L)$.*

Proof. Let C be the class of models of the sentences

$$\forall x \neg (P_i(x) \wedge P_j(x)), \quad i \neq j; \quad \forall x_1 \dots \forall x_n (R(x_1, \dots, x_n) \rightarrow P_i(x_1) \wedge \dots \wedge P_i(x_n)), \quad R \in \sigma_i.$$

The operation $A \mapsto A|_{\cup_i P_i}^A = [A|_{P_i}^A]$ from C to $\prod_i E_{\sigma_i}$ is evidently onto. The basic clopens of the product topology in $\prod_i E_{\sigma_i}(L)$ are given by finite conjunctions $\theta = \bigwedge_r \phi_r$ where $\phi_r \in L(\sigma_{i_r})$; hence, $A \models \theta \Leftrightarrow A|_{(\cup_r P_{i_r})}^A \models \theta$ for any A. As $(A|_{\cup_i P_i}^A)|_{(\cup_r P_{i_r})}^A = A|_{(\cup_r P_{i_r})}^A$, we have $A|_{\cup_i P_i}^A \models \theta \Leftrightarrow A \models \theta$, showing uniform continuity (cf. [C2]). \square

COROLLARY 5.5. *If L is closed under relativizations and $[\kappa, \lambda]$ -compact then any product space $\prod_1 E_{\sigma_1}(L)$ is $[\kappa, \lambda]$ -compact.*

Proof. Lemma 1.8, Sect I, and the previous Lemma. \square

It follows that a logic is $[\kappa, \lambda]$ -compact if and only if all the spaces $E_{\sigma}(L)$ are strongly $[\kappa, \lambda]$ -compact. Hence all the results of Section III apply. For example Corollary 2.7 becomes a case of Corollary 4.14. Similarly, the theorem of Makowsky and Shelah [Ma-Sh] that $[\kappa^+, \kappa^+]$ -compactness of a logic implies $[\kappa, \kappa]$ -compactness is a case of Corollary 4.16. Moreover, utilizing Theorem 4.9 and Proposition 5.4 we have:

THEOREM 5.6 (ABSTRACT COMPACTNESS THEOREM, [M-Sh]). *L is $[\kappa, \lambda]$ -compact if and only if there is a (κ, λ) -regular ultrafilter U related to L (which may be taken over $P_{\mu}(\kappa)$).*

Proof. L is $[\kappa, \lambda]$ -compact if and only if the spaces $\prod_1 E_{\sigma_1}(L)$ are all $[\kappa, \lambda]$ -compact by the previous Corollary. By Theorem 4.9., Sect. IV applied to the class $T = \{ E_{\sigma}(L) : \sigma \in \text{dom}(L) \}$ this is equivalent to the existence of a single (κ, λ) -regular ultrafilter U such that all $E_{\sigma}(L)$ are U -compact, which means U is related to L by Proposition 5.4. \square

BIBLIOGRAPHY

- [AU] P. Alexandroff and P. Urysohn, *Mémoire sur les espaces topologiques*, Verhandlungen dert Konink-lijke Akademie van Wetenschappen te Amsterdam, Afdeling Natuurkunde (Eerste Sectie) 14 (1929), 1-96.
- [BS] J.L.Bell and A.B. Slomson, *Models and Ultraproducts*. North Holland, 1971.
- [C1] X. Caicedo, *Compactness and normality in abstract model theory*. Ann of Pure and Applied Logic, to appear (1992).
- [C2] X. Caicedo, *Continuous operations on spaces of structures*. To appear in the Proceedings of the Symposium on "Quantifiers" celebrated at Sidlece, Poland, July, 1991.
- [Ch-K] C.C. Chang and J. Keisler, *Model Theory*, 2nd ed. North Holland 1973.
- [E] H.D. Ebbinghaus, *Extended logics; the general framework*. In Model theoretical logics. Springer Verlag, 1985.
- [K] Kanamori, *Weakly normal ultrafilters and irregular ultrafilters*. Trans. Amer. Math. Soc. 20 (1976)

- [K-P] K. Kunen and K.L. Pike, *On descendingly incomplete ultrafilters*. Jour of Symbolic Logic 36 (1971) 650-652.
- [K-V] K. Kunen and J.E. Vaughan, *Handbook of set theoretical topology*, North Holland, 1984.
- [L] P. Lindsdrom, *On extensions of elementary logic*. Theoria 35 (1969) 1-11.
- [Li] P. Lipparini, *Limit ultrapowers in abstract logic*. Journal of Symbolic Logic 52 (1987) 437-454
- [M] H. Mannila, *A topological characterization of $(\lambda, \nu)^*$ -compactness*. Ann. of Pure and Appl. Logic 25 (1983) 301-305.
- [Ma-Sh] J.A. Makowsky and S. Shelah, *Positive results in abstract model theory*. Ann. of Pure and Appl. Logic 24 (1983).
- [V1] J.E. Vaughan, *Some recent results in the theory of $[a, b]$ -compactness*. In TOPO 72, *General Topology and its Applications*. Lect. Notes in Math. 378. Springer Verlag 1974.
- [V2] J.E. Vaughan, *Some properties related to $[a, b]$ -compactness*. Fund. Math. 87 (3) (1975) 251-260.
- [W] S. Willard, *General Topology*. Addison Wesley, 1968
- [G] L. S. Gaal, *On the theory of (m, n) -compact spaces*, Pacific J. Math. 8 (1958) 721-734.
- [S] Y. M. Smirnov, *On topological spaces, compact in a given interval of powers*. Akad. Nauk SSSR Izvest. Ser. Mat. 14 (1950) 155-178.

RELATÓRIOS DE PESQUISA — 1992

- 01/92 Uniform Approximation the: Non-locally Convex Case — *João B. Prolla.*
- 02/92 Compactificação de $L^r_{\omega}(Q)$ com r Finito — *A. M. Sette and J. C. Cifuentes.*
- 03/92 Um Modelo para Aquisição da Especificação — *Cecilia Inés Sosa Arias and Ariadne Carvalho.*
- 04/92 Convergence Estimates for the Wavelet Galerkin Method — *Sônia M. Gomes and Elsa Cortina.*
- 05/92 Optimal Chemotherapy: A Case Study with Drug Resistance, Saturation Effect and Toxicity — *M. I. S. Costa, J. L. Boldrini and R. C. Bassanezi.*
- 06/92 On the Paper "Cauchy Completeness of Elementary Logic" of D. Mundici and A. M. Sette — *J. C. Cifuentes.*
- 07/92 What is the EM Algorithm for Maximum Likelihood Estimation in PET and How to Accelerate it — *Alvaro R. De Pierro.*
- 08/92 Bifurcation from infinity and multiple solutions for an elliptic system — *Raffaele Chiappinelli and Djairo G. de Figueiredo.*
- 09/92 Approximation Processes for Vector-Valued Continuous Functions — *João B. Prolla.*
- 10/92 Aplicação do Método de Fraissé à Compactificação de Lógicas com Quantificadores Co-filtro — *A. M. Sette and J. C. Cifuentes.*
- 11/92 Absolutely Summing Holomorphic Mappings — *Mário C. Matos.*
- 12/92 The Feynman-Dyson Proof of Maxwell Equations and Magnetic Monopoles — *Adolfo A. Jr. and Waldyr A. R. Jr.*
- 13/92 A Generalized Dirac's Quantization Condition for Phenomenological Non-abelian Magnetic Monopoles — *Adolfo M. Jr. and Waldyr A. R. Jr.*
- 14/92 Multiplicity Results for the 1-Dimensional Generalized p -Laplacian — *Pedro Ubilla.*
- 15/92 Nowhere Vanishing Torsion Closed Curves Always Hide Twice — *Sueli R. Costa. and Maria Del Carmen R. Fuster.*
- 16/92 Uniform Approximation of Continuous Convex-Cone-Valued Functions — *João B. Prolla.*
- 17/92 Monotonically Dominated Operators on Convex Cones — *A. O. Chiacchio, J. B. Prolla, M. L. B. Queiroz and M. S. M. Roversi.*
- 18/92 Testing the Concept of a Photon as an Extended Object in a Variation of Franson's Experiment — *V. Buonomano, A. J. R. Madureira and L. C. B. Ryff.*
- 19/92 A New Trust Region Algorithm for Boun Constrained Minimization — *Ana Friedlander, José Mario Martínez and Sandra A. Santos.*

- 20/92 A Priori Estimates for Positive Solutions of Semilinear Elliptic Systems Via Hardy-Sobolev Inequalities — *Ph. Clément, D.G. de Figueiredo and E. Mitidieri.*
- 21/92 On the Resolution of Linearly Constrained Convex Minimization Problems — *Ana Friedlander, José Mario Martínez and Sandra A. Santos.*
- 22/92 Convergência no Espectro Real de um Anel — *J. C. Cifuentes.*
- 23/92 Parallel Methods for Large Least Norm Problems — *Alvaro R. De Pierro. and Jose M. Lopes.*
- 24/92 A Generalization of Dirac Non-Linear Electrodynamics, and Spinning Charged Particles — *Waldyr A. Rodrigues Jr., Jayme Vaz Jr. and Erasmo Recami.*
- 25/92 Bifurcation of Certain Singularities of Sliding Vector Fields — *Marco Antonio Teizeira.*
- 26/92 Density of Infimum-Stable Convex Cone — *João B. Prolla.*
- 27/92 An Extended Decomposition Through Formalization in Product Spaces — *Alvaro R. De Pierro.*
- 28/92 Accelerating Iterative Algorithms for Symmetric Linear Complementarity Problems — *Alvaro R. De Pierro and José Marcos Lopes.*
- 29/92 Free Maxwell Equations, Dirac Equation, and Non-Dispersive de Broglie Wave-packets — *Waldyr A. Rodrigues Jr., Jayme Vaz Jr. and Erasmo Recami.*
- 30/92 New Results on the Equivalence of Regularization and Truncated Iteration for Ill-posed Problems — *Reginaldo J. Santos.*
- 31/92 Topological Defects: A Distribution Theory Approach — *Patricio S. Letelier and Anzhong Wang.*
- 32/92 Some Applications of the Generalized Khintchine's Inequality — *Mário C. Matos.*
- 33/92 A Technique for Finding The Center of A Polytope — *Antonio Carlos Moretti.*
- 34/92 On The Convergence of the Column-Updating Method — *José Mario Martínez.*
- 35/92 Post-Linear Formalism For Gravitating Strings: Crossed Straight Strings Collision — *D. V. Gal'tsov, Yu. V. Grats and P. S. Letelier.*
- 36/92 A Simple Model of Life Expectancy With Subjective Parameters — *R. C. Bassanezi and L. C. Barros.*
- 37/92 On the Growth of Caiman Crocodilus Yacare: A Strategy for Optimal Farming — *Alejandro B. Engel and R. C. Bassanezi.*
- 38/92 The Dirac Operator and the Structure of Riemann-Cartan-Weyl Spaces — *Quintino A. G. de Souza and Waldyr A. Rodrigues, Jr.*
- 39/92 On the Nature of the Gravitational Field — *Waldyr A. Rodrigues, Jr. and Quintino A. G. de Souza.*
- 40/92 Spinning Strings and Cosmic Dislocations — *D. V. Gal'tsov and P. S. Letelier.*
- 41/92 Natural Ultrabornological, Non-Complete, Normed Function Spaces — *Antonio Gilioli (edited by Klaus Floret and Chaim S. Hönig).*
- 42/92 Parallel and Semi-Parallel Immersions Into Space Forms — *Francesco Mercuri.*

- 43/92 Periodic Motions of Sucker Rod Pumping Systems — *Aloisio Freiria Neves and José Luiz Boldrini.*
- 44/92 Algorithms for Solving Nonlinear Systems of Equations — *José Mario Martínez.*
- 45/92 Spin and Electron Structure — *Matej Pavšič, Erasmo Recami, Waldyr A. Rodrigues Jr. and G. Daniele Maccarrone, Fabio Raciti, Giovanni Salesi.*
- 46/92 The Initial Value Problem For the Boussinesq Equations in a Time-Dependt Domain — *Carlos Conca Rozende and Marko Rojas Medar.*
- 47/92 A New Expansion for Jacobi Function — *E. Capelas de Oliveira.*
- 48/92 On the Equivalence of Convergences of Fuzzy Sets — *Marko Rojas Medar and Heriberto Román Flores.*
- 49/92 Monotonicity and Symmetry of Solutions of Elliptic Systems in General Domains — *Djairo G. de Figueiredo.*
- 50/92 Tendências em Programação Não Linear de Grande Porte — *José Mario Martínez.*
- 51/92 Teoría y Práctica en Optimización Numérica — *José Mario Martínez.*
- 52/92 The Clifford Bundle and the Nature of the Gravitational Field — *Waldyr A. Rodrigues Jr. and Quintino A.G. de Souza.*
- 53/92 Completamento e Compactificação de Espaços Zero-Dimensionais — *J. C. Cifuentes.*
- 54/92 Some Results on Polarized Partition Relations of Higher Dimension — *Walter Alexandre Carnielli and Carlos Augusto Di Prisco.*
- 55/92 The Generalized Wave Equation: Revisited — *G. C. Ducati and E. Capelas de Oliveira.*
- 56/92 The Four-Vertex Theorem for Constant Curvature Manifolds — *Sueli I. R. Costa and Marcelo Firer.*
- 57/92 The Riemann-Cartan-Weyl Geometry Generated by a Dirac-Hestenes Spinor Field — *Diego Lucio Rapoport-Campodónico, Waldyr A. Rodrigues, Jr., Quintino A. G. de Souza and Jayme Vaz Jr.*
- 58/92 On the Equivalence of Dirac and Maxwell Equations, and Quantum Mechanics — *Jayme Vaz Jr. and Waldyr A. Rodrigues.*
- 59/92 Convergência Fraca de Medidas em Espaços Métricos Separáveis e Completos — *Mario Antonio Gneri.*
- 60/92 Homoclinic Orbits for a Quasilinear Hamiltonian System — *Pedro Ubilla.*
- 61/92 Quasi - Newton Methods and the Solution of Some Fluid Mechanics Problems — *Márcia A. Gomes - Ruggiero, Daniel N. Kozakevich and José Mario Martínez.*
- 62/92 A New Construction of the Casimir Operators for the Fantappiè-de Sitter Group — *E. A. Notte Cuello and E. Capelas de Oliveira.*
- 63/92 Stationary Solutions for Generalized Boussinesq Models — *Sebastián A. Lorca and José Luiz Boldrini.*