

UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<https://www.scielo.br/j/spmj/a/VWMVmtTS798D4tP69NnSbQJ/>

DOI: <https://doi.org/10.1590/1516-3180.2022.0159.r1.19082022>

Direitos autorais / Publisher's copyright statement:

©2023 by Associação Paulista de Medicina. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

Prevalence risk of sarcopenia in older Brazilian adults during the pandemic: A cross-sectional analysis of the Remobilize Study

Patricia Parreira Batista^I, Monica Rodrigues Perracini^{II}, Juleimar Soares Coelho de Amorim^{III}, Maria do Carmo Correia de Lima^{IV}, Camila Astolphi Lima^V, Daniele Sirineu Pereira^{VI}, Renata Gonçalves Dantas^{VII}, Etienne Oliveira da Silva Fittipaldi^{VIII}, Aurélio Dias Santos^{IX}, Hércules Lázaro Moraes Campos^X, Leani Souza Máximo Pereira^{XI}

Universidade Federal de Minas Gerais (UFMG), Belo Horizonte (MG), Brazil

^IPT, MSc. Doctoral Student Postgraduate Program in Rehabilitation Sciences, Department of Physiotherapy, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte (MG), Brazil.
ID <https://orcid.org/0000-0002-8036-882X>

^{II}PT, PhD. Professor, Master's and Doctoral Programs in Physical Therapy, Universidade Cidade de São Paulo (UNICID), São Paulo (SP), Brazil; Master's and Doctoral Programs in Gerontology, Faculty of Medical Sciences, Universidade Estadual de Campinas (UNICAMP), Campinas (SP), Brazil.
ID <https://orcid.org/0000-0001-9331-3820>

^{III}PT, PhD. Professor, Physical Therapy Course, Instituto Federal do Rio de Janeiro, Rio de Janeiro (RJ), Brazil.
ID <https://orcid.org/0000-0003-3218-1769>

^{IV}PT, PhD. Faculty of Medical Sciences, Master's and Doctoral Programs in Gerontology, Universidade Estadual de Campinas (UNICAMP), Campinas (SP), Brazil.
ID <https://orcid.org/0000-0001-9018-5235>

^VPT, PhD. Postdoctoral Student of Master's and Doctoral Program in Physical Therapy, Universidade Cidade de São Paulo (UNICID), São Paulo (SP), Brazil.
ID <https://orcid.org/0000-0001-9882-6975>

^{VI}PT, PhD. Professor, Postgraduate Program in Rehabilitation Sciences, Department of Physiotherapy, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte (MG), Brazil.
ID <https://orcid.org/0000-0002-4868-9244>

^{VII}PT, MSc. Doctoral Student of Master's and Doctoral Program in Physical Therapy, Universidade Cidade de São Paulo (UNICID), São Paulo (SP), Brazil; and Professor of Physical Therapy, Universidade Estadual do Sudoeste da Bahia (UESB), Vitória da Conquista (BA), Brazil.
ID <https://orcid.org/0000-0002-1575-7204>

^{VIII}PT, PhD. Professor, Department of Physiotherapy, Universidade Federal de Pernambuco (UFPE), Recife (PE), Brazil.
ID <http://orcid.org/0000-0002-1524-6930>

^{IX}PT, MSc. Professor, Department of Physiotherapy, Centro Universitário Dr. Leão Sampaio (UNILEÃO), Juazeiro do Norte (CE), Brazil.
ID <https://orcid.org/0000-0003-3693-9864>

^XPT, MSc. Professor, Department of Physiotherapy, Universidade Federal do Amazonas/Instituto de Saúde e Biotecnologia (UFAM/ISB), Coari (AM), Brazil. Doctoral Student, Postgraduate Program in Public Health, Universidade Federal do Espírito Santo (UFES), Vitória (ES), Brazil.
ID <https://orcid.org/0000-0002-6919-8161>

^{XI}PT, PhD. Professor, Postgraduate Program in Rehabilitation Sciences, Department of Physiotherapy, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte (MG), Brazil.
ID <https://orcid.org/0000-0001-7253-4292>

KEY WORDS (MeSH terms):

Sarcopenia.
Mobility limitation.
Aged.
Physical distancing.
Pandemics.

AUTHORS' KEY WORDS:

SARC-F.
Risk of sarcopenia.
Screening.
Older adults.

ABSTRACT

BACKGROUND: Social distancing has led to lifestyle changes among older adults during the coronavirus disease 2019 (COVID-19) pandemic.

OBJECTIVES: This study aimed to estimate the prevalence risk of sarcopenia (RS) and investigate its associated factors during the COVID-19 pandemic in older Brazilian adults.

DESIGN AND SETTING: Cross-sectional observational analysis of baseline data as part of the Remobilize Study.

METHODS: Participants in the study were older adults (≥ 60 years), excluding those who were bedridden or institutionalized. The data collected consisted of answers about the RS (SARC-F), functional status, walking, sedentary behavior (SB), pain, comorbidity, and life space mobility.

RESULTS: A total of 1,482 older adults (70 ± 8.14 years, 74% women) participated in the study, and an RS prevalence of 17.1% was found. (95% confidence interval [CI] 15.25–19.15%). The adjusted multivariate model showed a significant association between RS and functional limitation (odds ratio [OR]: 19.05; CI 13.00–28.32), comorbidity (OR: 5.11; CI 3.44–7.81), pain (OR: 4.56; CI 3.33–6.28), total walking (OR: 0.99; CI 0.99–1.00), SB of 8–10 hours (OR: 1.85; CI 1.15–2.93), and SB of > 10 hours (OR: 3.93; CI 2.48–6.22). RS was associated with mobility during the pandemic (OR: 0.97; CI 0.96–0.98). $P < 0.05$.

CONCLUSIONS: During the pandemic, the prevalence of RS in older Brazilians was estimated at 17.1%. Moderate to severe functional limitation, comorbidities, presence of pain, walking, longer SB period, and reduced life space mobility significantly contributed to RS in older adults during the pandemic.

INTRODUCTION

Social restriction policies and lifestyle changes favor a reduction in mobility and the level of physical activity (PA), leading to a higher proportion of inactive people and an increase in sedentary behaviors (SB) during the pandemic.¹⁻⁴ A decline in life space mobility contributes to a reduction in intrinsic capacity, higher risk of sarcopenia (RS), and other adverse health consequences.^{5,6} After 7 days of total bed rest, there is already a significant deterioration in muscle function in community-dwelling older adults, and 2,000 steps per day are not enough to prevent these deleterious effects on the musculature.⁷ Coker et al. reported that a 15-day bed rest induces a significant reduction in fat-free muscle mass, poor performance, and increased fat in older individuals, which negatively impacts their mobility.⁸

A longer SB time observed during the pandemic is related to a worse prognosis in health conditions and a higher RS.^{1-3,7,9} These factors can alter the homeostasis between the pro- and anti-inflammatory systemic components and muscle anabolism and catabolism, leading to the reduction of physiological reserves in older adults. Consequences such as increased plasma pro-inflammatory cytokines, greater muscle catabolism drive, and anabolic and insulin resistance lead to a deleterious cycle of muscle function, explaining the higher incidence of RS in this population.¹⁻³

Sarcopenia is a generalized and progressive musculoskeletal disorder that is defined as a reduction in muscle mass and strength. It is a multifactorial disease with dynamic interrelationships and is commonly associated with a cascade of negative repercussions on health, functional limitation, and mortality.¹⁰⁻¹² Consequently, due to its considerable clinical impact on older individuals,

it increases health-related expenses and imposes a burden on the public health system, being more costly in socially unequal and/or developing countries, such as Brazil.^{10,12,13} Updates from the European Working Group on Sarcopenia in Older People (EWGSOP2) and the Asian Working Group for Sarcopenia proposed the practice of population screening for RS in older people through strength, assistance with walking, rising from a chair, climbing stairs, and falls (SARC-F) questionnaire, a self-reported screening questionnaire.^{10,14} Identifying sarcopenia in its early stages enables the control of its progression and/or reversal of the individual's clinical condition, thereby reducing the negative impacts caused by the disease.^{3,10,11,14,15}

OBJECTIVE

Due to the abovementioned reasons, this study aimed to verify the prevalence of RS and investigate the factors associated with the presence of RS during the coronavirus disease 2019 (COVID-19) pandemic.

METHODS

Design and sample

This study presents a cross-sectional analysis of baseline data collected from May to July 2020 through an online questionnaire as part of the Remobilize Study (www.remobilize.com.br).⁴ Using convenience snowball sampling, the online questionnaire (SurveyMonkey platform) was distributed throughout the Brazilian territory via social media (Facebook and Instagram), WhatsApp groups, social groups for older adults, and/or their friends and acquaintances. A pilot project for calibration and adjustments was conducted in advance. This study was approved by the University City of São Paulo Research Ethics Committee (May 18, 2020; CAAE 31592220.6.0000.0064) and is currently under progress.

The sample population consisted of community-dwelling older Brazilians (≥ 60 years) without distinction of sex, race, and/or social class. Following the exclusion criteria, those residing in long-term care facilities and/or bedridden were not eligible to participate in the study.⁴ Participants who presented with disabilities were allowed to have the questions be answered by a family member or caregiver.¹⁶ Participants without familiarity with the Internet were able to answer the survey via telephone.⁴

Measures

The sociodemographic, clinical, and lifestyle data are presented in **Table 1**. The self-reported functional comorbidity index questionnaire was used to detect the presence of comorbidities (two or more chronic diseases).¹⁷ All participants answered questions about the presence or absence of pain.

The SARC-F questionnaire is recommended by the EWGSOP2 and the Asian Working Group for Sarcopenia as a population screening tool for RS.^{10,14} The final score ranges from zero to ten points, and a score of ≥ 4 points identifies individuals with sarcopenia. SARC-F has a high specificity, but low to moderate sensitivity.^{10,14,15,18} Population screening for RS (SARC-F) allows the exclusion of older patients with preserved muscle function in primary health care and identification of changes in the early stages of muscle function, functionality, and RS in older adults.^{10,14,15,19}

Functional performance was assessed using the Older American Resources and Services questionnaire that has been translated and validated for the Brazilian population (BOMFAQ).^{20,21} It is a self-report questionnaire on the ability to perform 15 functional activities (eight basic and seven instrumental tasks). The scores for the activities performed with difficulty or requiring help were added, ranging from 0–15 points. Older adults were classified based on their scores: no (0), slight (1–3), moderate (4–6), and severe (> 7) functional limitation.²²

SB was assessed using one question about the duration of sitting activities in the prior week, referring to indoor activities (≤ 4 hours/day, 5–7 hours/day, 8–10 hours/day or ≥ 10 hours/day). Walking, including PA, utilitarian walking, and walking time, was assessed using the Incidental and Planned Exercise Questionnaire.²³ Validated for older adults, this is a simple, self-report questionnaire probing on walking activities during the prior week, specifically on the frequency and duration of the activity. The final score for walking as physical exercise and utilitarian walking was given by the product of frequency and duration for each item (minutes/week). The total walking time was calculated as the sum of walking as PA and utilitarian walking.

Life space mobility was measured using the Life-Space Assessment (LSA).²⁴ It estimates the individual perspective of mobility relative to the spatial area in five levels of life space in the prior week: mobility in the rooms at home, outside the bedroom (level 1), outside the home (level 2), a neighborhood close to home (level 3), circulation within the municipality where they reside (level 4), and inter-municipal areas (level 5). The answers were based on the frequency and need for mobility devices. The score was calculated as the product of frequency and performance skill, extracting a score based on level and the total by the sum of levels (0–120 points). Higher final scores indicated better mobility performance in the life space.

Statistical analysis

The prevalence of RS in participants was estimated using a 95% confidence interval (CI). Descriptive statistics were performed using absolute and relative frequencies for the total sample and RS, respectively. Continuous variables did not show a normal distribution in the Shapiro–Wilk test; therefore, the data are presented as medians and interquartile ranges. To compare the

Table 1. Total sample descriptive data and comparison between the groups with (strength, assistance with walking, rising from a chair, climbing stairs, and falls, SARC-F ≥ 4 points) and without risk of sarcopenia (RS) (SARC-F < 4 points)

		SARC-F		P value
		< 4 points (n = 1,228)	≥ 4 points (n = 254)	
Age, %	60–69 years	61.3%	31.5%	< 0.0001
	70–79 years	28.8%	26.0%	
	80 years and older	9.9%	42.5%	
Sex, %	Male	27.9%	16.9%	0.001
	Female	72.1%	83.1%	
Marital status, %	Single	10.3%	10.2%	< 0.0001
	Married	56.7%	39.0%	
	Divorced	12.7%	11.0%	
	Widowed	20.3%	39.8%	
Education, %	Illiterate	6.4%	14.9%	< 0.0001
	1–4 years	16.5%	31.5%	
	5–8 years	11.9%	13.8%	
	9 years or more	65.2%	39.8%	
Income ^a , %	Up to 1× minimum wage	32.6%	44.1%	< 0.0001
	2–3× minimum wage	27.9%	27.9%	
	4–7× minimum wage	19.4%	11%	
	8–10× minimum wage	7.6%	7.9%	
	More than 10× minimum wage	12.5%	9.1%	
Occupation, %	Active	39.2%	24.8%	< 0.0001
	Inactive	55.3%	61.8%	
	Unemployed	5.5%	13.4%	
Sitting time, %	< 4 hours	48.6%	28.8%	< 0.0001
	5–7 hour	31.0%	31.1%	
	8–10 hour	12.6%	16.9%	
	> 10 hours	7.8%	23.2%	
BOMFAQ (4 pts +), %		10.3%	73.6%	< 0.0001
Comorbidities (≥ 2), %		50.40%	87.40%	< 0.0001
Pain (yes), %		21.6%	55.5%	< 0.0001
Walking (exercise) Med (IQR)		0 (0–25.31)	0 (0–0)	< 0.0001
Walking (utilitarian) Med (IQR)		7.5 (0–33.75)	0 (0; 0)	< 0.0001
Walking (total) Med (IQR)		7.5 (0; 101.20)	0 (0; 7.5)	< 0.0001
LSA - During pandemic Med (IQR)	Total score	36 (24; 52)	24 (12; 32)	< 0.0001
	Level 1	8 (8; 8)	8 (6; 8)	< 0.0001
	Level 2	16 (12; 16)	12 (4; 16)	< 0.0001
	Level 3	6 (0; 12)	0 (0; 6)	< 0.0001
	Level 4	8 (0; 16)	0 (0; 4)	< 0.0001
	Level 5	0 (0; 0)	0 (0; 0)	0.0002

Med = median; IQR = interquartile range (1st and 3rd IQR); LSA = Life-Space Assessment; BOMFAQ = Brazilian OARS Multidimensional Functional Assessment Questionnaire; a score of four points or more refers to the presence of moderate to severe functional limitation; walking (as exercise, utilitarian, and total) in the previous week (min/week). ^aminimum wage in Brazil = R\$ 1,100.00 per month, corresponding to US\$ 194.01 (April 5, 2021).

groups with and without RS, Pearson's chi-square test was used for categorical variables and the Mann–Whitney test was used for continuous variables. The association between independent variables and outcome was based on odds ratios (ORs) estimates and their respective CIs through logistic regression without (crude model) and with adjustment (adjusted model). All analyses were

performed using Stata 14.0 (StataCorp LLC, College Station, Texas, United States), with a 5% statistical significance level.

RESULTS

A total of 1,482 participants were included in this study, and the study flowchart is shown in **Figure 1**. The prevalence of RS

during the pandemic was 17.1% (CI 15.25–19.15%). The distribution of SARC-F and total score items by age group and total scores are shown in **Figure 2**. Statistically significant differences were observed between the groups with and without RS in terms of age, sex, marital status, education, income, occupation, walking (exercise, utilitarian, and total), sitting time, functional limitation, presence of comorbidities, and pain. The RS group had a higher proportion of participants aged 80 years or older (42.5%), women (83.1%), lower income (44.1%), and 73.6% presented with moderate to severe functional limitation (**Table 1**). There were statistically significant differences in total LSA scores

during the pandemic. Lower LSA scores were observed in older patients with RS. During the pandemic, there was a difference between older patients with and without RS for all walking variables (exercise and total), with lower values in the RS group.

Crude logistic regression analysis showed a significant association for all analyzed variables. After adjusting for sociodemographic factors, the following variables remained statistically significant, as seen in **Table 2**: moderate to severe functional limitations, comorbidity, pain, walking (exercise and total), SB 8–10 hours, SB > 10 hours, and total LSA score during the pandemic.

DISCUSSION

The results showed a high prevalence of RS in older Brazilians at the beginning of the COVID-19 pandemic in Brazil and a substantial association between RS and moderate to severe functional limitation, comorbidities, pain, and a positive gradient with the number of hours in SB. The OR for RS increased from 1.85 in older patients who reported 8 to 10 hours of SB to 3.93 in those with 10 hours or more of SB. Older patients with moderate to severe functional limitation were 19.05 times more likely to be at RS. Furthermore, greater mobility in living spaces lowered the chances of RS during the pandemic.

The prevalence of RS (17.1%) in the present study was substantially higher than that found in studies before the COVID-19 pandemic.^{25–28} Dodds et al. reported a 4% prevalence of RS in 1,686 British older adults (aged ≥ 69 years),²⁵ while Kim and Won reported a rate of 7.5% among 2,123 Korean older adults (75.9 years).²⁶ Studies with a model of activity reduction (steps per day) in elders pointed to negative repercussions of greater catabolic drive on their musculature and metabolic and inflammatory markers during a short period of mobility restriction.^{29,30} With a




Figure 1. Study flowchart.




Figure 2. Prevalence of risk of sarcopenia in the total sample and by age group. Frequency of the items included in the strength, assistance with walking, rising from a chair, climbing stairs, and falls (SARC-F) questionnaire for older at risk for sarcopenia.

Table 2. Logistic regression analysis to verify the association of the factors contributing to the risk of sarcopenia

		Crude model			Adjusted model		
		OR	95% CI	P value	OR	95% CI	P value
BOMFAQ (4 pts +)		24.20	17.42; 33.40	< 0.001	19.05	13.00; 28.32	< 0.001
(ref.: 0–3 pts)							
Comorbidities (≤ 2)		6.82	4.70; 10.22	< 0.001	5.11	3.44; 7.81	< 0.001
(ref.: 0–1)							
Pain		4.54	3.42; 6.03	< 0.001	4.56	3.33; 6.28	< 0.001
(ref.: absence of)							
Walking (exercise)		0.99	0.986; 0.994	< 0.001	0.99	0.989; 0.997	0.001
Walking (utilitarian)		0.99	0.994; 0.999	0.041	0.99	0.997; 1.001	0.346
Walking (total)		0.99	0.993; 0.997	< 0.001	0.99	0.995; 0.999	0.008
Sedentary behavior (sitting time; ref. < 4 hours)	5–7 hour	1.69	1.20; 2.39	0.003	1.41	0.97; 2.04	0.072
	8–10 hour	2.26	1.48; 3.42	< 0.001	1.85	1.15; 2.93	0.01
	> 10 hours	5.02	3.34; 7.53	< 0.001	3.93	2.48; 6.22	< 0.001
LSA - During pandemic	Total score	0.95	0.94; 0.96	< 0.001	0.97	0.96; 0.98	< 0.001
	Level 1	0.79	0.74; 0.84	< 0.001	0.83	0.77; 0.89	< 0.001
	Level 2	0.90	0.88; 0.92	< 0.001	0.92	0.90; 0.95	< 0.001
	Level 3	0.92	0.91; 0.94	< 0.001	0.95	0.93; 0.97	< 0.001
	Level 4	0.93	0.91; 0.95	< 0.001	0.97	0.95; 0.98	< 0.001
	Level 5	0.92	0.88; 0.96	< 0.001	0.96	0.91; 0.99	0.047

Med = median; IQT = interquartile range (1st and 3rd IQR); LSA = Life-Space Assessment; BOMFAQ = Brazilian OARS Multidimensional Functional Assessment Questionnaire; a score of four points or more refers to the presence of moderate to severe functional limitation. Walking (as exercise, utilitarian, and total) in the previous week (minutes/week).

76% reduction in steps per day (< 1,500 steps/day) in 14 days, Breen et al. demonstrated a 3.9% reduction in fat-free lean mass; reduced insulin sensitivity (43%); and increased pro-inflammatory cytokines, TNF- α (12%), and C-reactive protein (25%) levels in 10 healthy older adults after the intervention (72.3 years).³⁰ These findings may support the higher RS prevalence in our study.

In Brazil, Barbosa-Silva et al. reported that sarcopenia had a prevalence of 8.4% (EWGSOP1) in 179 older adults.¹⁸ Sarcopenia (SARC-F \geq 6 points) and muscle function decline (SARC-F \geq 4 points) were 17.3% and 34.6%, respectively. The authors proposed the addition of calf circumference measurements to SARC-F to improve the instrument's measurement accuracy.¹⁸ The EWGSOP2 establishes an overlap of muscle strength in relation to muscle mass as a primary parameter in the diagnosis of sarcopenia, as muscle strength is the most reliable measure of muscle function.¹⁰ Furthermore, it is associated with adverse health outcomes and facilitates the use of the diagnostic algorithm in clinical practice.^{10,15,31} Thus, the present study considered values \geq 4 in the SARC-F as the cutoff point because of the improved accuracy in diagnosing muscle function in older Brazilian people and support from the scientific community.^{10,14,15,17} In addition, it is impossible to conduct anthropometric measurements due to pandemic-related restrictions.

Findings on sociodemographic differences between participants with and without RS were similar to those found in studies before the pandemic, whether in older patients with RS or with sarcopenia or on diagnostic parameters for sarcopenia.^{18,25–27,31–35}

The difference in the presence of moderate to severe functional limitation between the groups was significant. After adjusted logistic regression, those with moderate to severe functional limitation were 19.05 times more likely to be at RS. Similar findings were reported by Rolland et al, with a sample of 3,025 French older adults (80.5 years).³³ The authors found a lower functional performance in older adults at RS compared to the total sample and a significant association with reduced functional performance based on the gait speed and chair stand test results (OR: -0.04; CI 0.05–0.03 and OR: 13.1; CI 11.5–14.7). Longitudinal analyses with a 6-year follow-up confirmed the ability of SARC-F score \geq 4 points (RS) to predict reduced functional performance.³¹

Our logistic regression analysis, adjusted for sociodemographic factors, showed a significant association of the presence of comorbidity with RS, corroborating previous studies.^{25,33} Given the context of the pandemic, the combination of psychobehavioral factors, such as stress, worse sleep quality, food routine, and mood, as well as medical treatment and functional rehabilitation discontinuation, increased physical inactivity and SB, which triggered an accelerated progression of established chronic diseases due to the greater active systemic pro-inflammatory profile and higher muscle catabolism drive. Thus, monitoring these factors in older adults is necessary during and after the pandemic, including socio-demographic factors and their specifications.^{1,2,3,36}

Pain contributed to the highest RS among the participants in this study. Corroborating this study, Lustosa et al. investigated RS in 322 older Brazilian women complaining of non-specific acute

lower back pain, and the results showed an association between pain intensity and poor mobility and balance.³⁷ The authors pointed out that RS, if present in older women with lower back pain, can negatively influence functionality.³⁷ Pain is multifactorial and subjective. Moreover, psychosocial factors are known to interfere with pain and its pro-inflammatory process, and social isolation predisposes to the development of chronic pain.³⁸ Thus, pain in older people should not be neglected during and after the pandemic, and directions for non-pharmacological and pharmacological interventions should be considered.

A positive and significant association was observed between SB and RS, with a “dose-response” effect for a more extended period of SB, causing older adults with 10 h or more of SB per day to be 3.93 (CI 2.48–6.22; compared to < 4 hours) times more likely to be at RS. With a sample of 1,068 older adults (72.1 years), Tzeng et al. demonstrated that sitting for 7 hours or more per day was significantly associated with RS (OR: 1.98; CI 1.09–3.59).³⁹ Smith et al. also investigated the relationship between SB and sarcopenia in 14,585 older adults from six low- and middle-income countries.⁴⁰ The authors identified that regardless of the PA level and presence of comorbidities, 11 hours or more of SB increases RS by 2.14 times (CI 1.06–4.33; compared to < 4 hours), and each additional hour per day of SB was related to an increased risk of RS by 1.06 (CI 1.04–1.10).⁴⁰ Thus, our results confirm that the more sedentary the lifestyle during the pandemic, the greater the probability of RS and possibly the worse is the health condition and muscle function prognosis.

It is known that physical inactivity and PA levels below the recommendations proposed by the World Health Organization (WHO) are more frequent in older adults,^{3,41,42} and sarcopenic individuals have lower PA levels than non-sarcopenic individuals.^{25,34,43} In the present study, there was a difference in walking (exercise, utilitarian, and total) between the two groups, reflecting the low PA level in participants with RS during the pandemic. Saraiva et al. found a reduction in the practice of regular PA (≥ 3 times/week) during the pandemic in 557 older Brazilian (80 ± 8 years), ranging from 42% active (pre-pandemic) to 26% (during the pandemic).⁴⁴ Tzeng et al. showed that insufficiently active older adults had a 5.14 (CI 3.04–8.70) times higher RS.³⁹ Thus, physical inactivity is a modifiable risk factor for sarcopenia, and physical exercise is the first-line treatment for this muscle disease.^{2,15,32}

Our results showed lower life space mobility during the pandemic in the RS group. A similar and significant difference was found in a study published before the pandemic.⁴⁵ In this study, the group without RS had lower average age, was more active, and presented with a lower percentage of comorbidity than the group with RS. Higher mobility rates are associated with better muscle function, functional and cognitive performance, and social support.⁴⁶ This finding serves as a warning for this target

population, given the prolonged course of the pandemic and the deleterious relationship between restriction of outdoor mobility and skeletal musculature.

Some limitations of this study must be considered. Snowball sampling was carried out on an online platform, differentiating our sample from the general community. The participants could have had access to the Internet and a higher level of education or social support as opposed to the older Brazilian population in general. Our findings were extrapolated to older adults with characteristics similar to those of our sample. In addition, the study had a cross-sectional design, making it impossible to identify causality in the analyzed relationships. However, this cross-sectional analysis aimed to identify and verify RS and its contributing factors in the Rede Remobilize (Wave 1) cohort and establish a baseline for future longitudinal studies on the impacts of the pandemic and RS in older individuals. To our knowledge, this is the first study to assess RS in a consistent sample of community-dwelling older adults in Brazil during the pandemic. Finally, this study encourages the use of SARC-F in monitoring older patients because it is a viable tool in clinical practice for screening for muscle function decline and RS, as it allows for the adequacy of future health care actions in favor of healthy aging.^{19,47}

CONCLUSIONS

Moderate to severe functional limitation, comorbidity, pain, longer period of SB, and reduced life space mobility significantly contributed to the RS in older Brazilian adults during the pandemic. Longitudinal studies monitoring functional trajectories and adverse health outcomes in older patients with RS during the pandemic should be encouraged to understand the associated modifiable factors and preventive actions against this critical muscle dysfunction.

REFERENCES

1. Moro T, Paoli A. When COVID-19 affects muscle: effects of quarantine in older adults. *Eur J Transl Myol.* 2020;30(2):9069. PMID: 32782767; <https://doi.org/10.4081/ejtm.2019.9069>.
2. Kirwan R, McCullough D, Butler T, et al. Sarcopenia during COVID-19 lockdown restrictions: long-term health effects of short-term muscle loss. *Geroscience.* 2020;42(6):1547-78. PMID: 33001410; <https://doi.org/10.1007/s11357-020-00272-3>.
3. Roschel H, Artioli GG, Gualano B. Risk of Increased Physical Inactivity During COVID-19 Outbreak in Older People: A Call for Actions. *J Am Geriatr Soc.* 2020;68(6):1126-8. PMID: 32392620; <https://doi.org/10.1111/jgs.16550>.
4. Perracini MR, de Amorim JSC, Lima CA, et al. Impact of COVID-19 Pandemic on Life-Space Mobility of Older Adults Living in Brazil: REMOBILIZE Study. *Front Public Health.* 2021;9:643640. PMID: 33898378; <https://doi.org/10.3389/fpubh.2021.643640>.

5. Webber SC, Porter MM, Menec VH. Mobility in older adults: a comprehensive framework. *Gerontologist*. 2010;50(4):443-50. PMID: 20145017; <https://doi.org/10.1093/geront/gnq013>.
6. Miyashita T, Tadaka E, Arimoto A. Cross-sectional study of individual and environmental factors associated with life-space mobility among community-dwelling independent older people. *Environ Health Prev Med*. 2021;26(1):9. PMID: 33461488; <https://doi.org/10.1186/s12199-021-00936-2>.
7. Arentson-Lantz E, Galvan E, Wachter A, Fry CS, Paddon-Jones D. 2,000 Steps/Day Does Not Fully Protect Skeletal Muscle Health in Older Adults During Bed Rest. *J Aging Phys Act*. 2019;27(2):191-7. PMID: 29989486; <https://doi.org/10.1123/japa.2018-0093>.
8. Coker RH, Hays NP, Williams RH, Wolfe RR, Evans WJ. Bed rest promotes reductions in walking speed, functional parameters, and aerobic fitness in older, healthy adults. *J Gerontol A Biol Sci Med Sci*. 2015;70(1):91-6. PMID: 25122628; <https://doi.org/10.1093/gerona/glu123>.
9. Gianoudis J, Bailey CA, Daly RM. Associations between sedentary behaviour and body composition, muscle function and sarcopenia in community-dwelling older adults. *Osteoporos Int*. 2015;26(2):571-9. PMID: 25245026; <https://doi.org/10.1007/s00198-014-2895-y>.
10. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48(4):601. Erratum for: *Age Ageing*. 2019;48(1):16-31. PMID: 31081853; <https://doi.org/10.1093/ageing/afy169>.
11. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010;39(4):412-23. PMID: 20392703; <https://doi.org/10.1093/ageing/afq034>.
12. Beaudart C, Zaaria M, Pasleau F, Reginster JY, Bruyère O. Health Outcomes of Sarcopenia: A Systematic Review and Meta-Analysis. *PLoS One*. 2017;12(1):e0169548. PMID: 28095426; <https://doi.org/10.1371/journal.pone.0169548>.
13. Bruyère O, Beaudart C, Ethgen O, Reginster JY, Locquet M. The health economics burden of sarcopenia: a systematic review. *Maturitas*. 2019;119:61-9. PMID: 30502752; <https://doi.org/10.1016/j.maturitas.2018.11.003>.
14. Chen LK, Woo J, Assantachai P, et al. Asian Working Group for Sarcopenia: 2019 Consensus Update on Sarcopenia Diagnosis and Treatment. *J Am Med Dir Assoc*. 2020;21(3):300-307.e2. PMID: 32033882; <https://doi.org/10.1016/j.jamda.2019.12.012>.
15. Dent E, Morley JE, Cruz-Jentoft AJ, et al. International Clinical Practice Guidelines for Sarcopenia (ICFSR): Screening, Diagnosis and Management. *J Nutr Health Aging*. 2018;22(10):1148-61. PMID: 30498820; <https://doi.org/10.1007/s12603-018-1139-9>.
16. Maurus J, Terzer T, Benner A, et al. Validation of a proxy-reported SARC-F questionnaire for current and retrospective screening of sarcopenia-related functional impairments. *J Cachexia Sarcopenia Muscle*. 2022;13(1):264-75. PMID: 34898035; <https://doi.org/10.1002/jcsm.12871>.
17. Groll DL, To T, Bombardier C, Wright JG. The development of a comorbidity index with physical function as the outcome. *J Clin Epidemiol*. 2005;58(6):595-602. PMID: 15878473; <https://doi.org/10.1016/j.jclinepi.2004.10.018>.
18. Barbosa-Silva TG, Menezes AM, Bielemann RM, et al. Enhancing SARC-F: Improving Sarcopenia Screening in the Clinical Practice. *J Am Med Dir Assoc*. 2016;17(12):1136-41. PMID: 27650212; <https://doi.org/10.1016/j.jamda.2016.08.004>.
19. Cesari M, Marzetti E, Calvani R. Sarcopenia and SARC-F: "Perfect is the Enemy of Good". *J Am Med Dir Assoc*. 2021;22(9):1862-3. PMID: 34456011; <https://doi.org/10.1016/j.jamda.2021.07.030>.
20. Duke University Center for the Study of Aging and Human Development. Multi-dimensional Functional Assessment: The OARS methodology. Durham, NC: Duke University Center for the Study of Aging and Human Development; 1978.
21. Ramos LR, Toniolo J, Cendoroglo MS, et al. Two-year follow-up study of elderly residents in S. Paulo, Brazil: methodology and preliminary results. *Rev Saude Publica*. 1998;32(5):397-407. PMID: 10030055; <https://doi.org/10.1590/s0034-89101998000500001>.
22. Ferreira MT, Matsudo SM, Ribeiro MC, Ramos LR. Health-related factors correlate with behavior trends in physical activity level in old age: longitudinal results from a population in São Paulo, Brazil. *BMC Public Health*. 2010;10:690. PMID: 21067591; <https://doi.org/10.1186/1471-2458-10-690>.
23. Delbaere K, Hauer K, Lord SR. Evaluation of the incidental and planned activity questionnaire (IPEQ) for older people. *Br J Sports Med*. 2010;44(14):1029-34. PMID: 19474003; <https://doi.org/10.1136/bjsm.2009.060350>.
24. Simões MDSM, Garcia IF, Costa LDC, Lunardi AC. Life-Space Assessment questionnaire: Novel measurement properties for Brazilian community-dwelling older adults. *Geriatr Gerontol Int*. 2018;18(5):783-9. PMID: 29372585; <https://doi.org/10.1111/ggi.13263>.
25. Dodds RM, Murray JC, Robinson SM, Sayer AA. The identification of probable sarcopenia in early old age based on the SARC-F tool and clinical suspicion: findings from the 1946 British birth cohort. *Eur Geriatr Med*. 2020;11(3):433-41. PMID: 32297269; <https://doi.org/10.1007/s41999-020-00310-5>.
26. Kim M, Won CW. Sarcopenia in Korean Community-Dwelling Adults Aged 70 Years and Older: Application of Screening and Diagnostic Tools From the Asian Working Group for Sarcopenia 2019 Update. *J Am Med Dir Assoc*. 2020;21(6):752-8. PMID: 32386844; <https://doi.org/10.1016/j.jamda.2020.03.018>.
27. Mo YH, Zhong J, Dong X, et al. Comparison of Three Screening Methods for Sarcopenia in Community-Dwelling Older Persons. *J Am Med Dir Assoc*. 2021;22(4):746-50.e1. PMID: 32669238; <https://doi.org/10.1016/j.jamda.2020.05.041>.
28. Yang M, Hu X, Xie L, et al. SARC-F for sarcopenia screening in community-dwelling older adults: Are 3 items enough? *Medicine (Baltimore)*. 2018;97(30):e11726. PMID: 30045339; <https://doi.org/10.1097/MD.00000000000011726>.

29. Perkin O, McGuigan P, Thompson D, Stokes K. A reduced activity model: a relevant tool for the study of ageing muscle. *Biogerontology*. 2016;17(3):435-47. PMID: 26506931; <https://doi.org/10.1007/s10522-015-9613-9>.
30. Breen L, Stokes KA, Churchward-Venne TA, et al. Two weeks of reduced activity decreases leg lean mass and induces "anabolic resistance" of myofibrillar protein synthesis in healthy elderly. *J Clin Endocrinol Metab*. 2013;98(6):2604-12. PMID: 23589526; <https://doi.org/10.1210/jc.2013-1502>.
31. Malmstrom TK, Miller DK, Simonsick EM, Ferrucci L, Morley JE. SARC-F: a symptom score to predict persons with sarcopenia at risk for poor functional outcomes. *J Cachexia Sarcopenia Muscle*. 2016;7(1):28-36. PMID: 27066316; <https://doi.org/10.1002/jcsm.12048>.
32. Cruz-Jentoft AJ, Landi F, Schneider SM, et al. Prevalence of and interventions for sarcopenia in ageing adults: a systematic review. Report of the International Sarcopenia Initiative (EWGSOP and IWGS). *Age Ageing*. 2014;43(6):748-59. PMID: 25241753; <https://doi.org/10.1093/ageing/afu115>.
33. Rolland Y, Dupuy C, Abellan Van Kan G, et al. Sarcopenia Screened by the SARC-F Questionnaire and Physical Performances of Elderly Women: A Cross-Sectional Study. *J Am Med Dir Assoc*. 2017;18(10):848-52. PMID: 28629717; <https://doi.org/10.1016/j.jamda.2017.05.010>.
34. Martone AM, Marzetti E, Salini S, et al. Sarcopenia Identified According to the EWGSOP2 Definition in Community-Living People: Prevalence and Clinical Features. *J Am Med Dir Assoc*. 2020;21(10):1470-4. PMID: 32359893; <https://doi.org/10.1016/j.jamda.2020.03.007>.
35. Brennan-Olsen SL, Vogrin S, Balogun S, et al. Education, occupation and operational measures of sarcopenia: Six years of Australian data. *Australas J Ageing*. 2020;39(4):e498-e505. PMID: 32969133; <https://doi.org/10.1111/ajag.12816>.
36. Christoforetti M, Duca GFD, Benedetti TRB, Malta DC. Sociodemographic determinants of multimorbidity in Brazilian adults and older adults: a cross-sectional study. *Sao Paulo Med J*. 2022;140(1):115-22. PMID: 35043867; <https://doi.org/10.1590/1516-3180.2021.0105.R1.31052021>.
37. Lustosa LP, Tavares CCA, Vital DKJC, et al. Risco de sarcopenia em idosos com queixa de dor lombar aguda. *Fisioter Pesqui*. 2018;25(3):260-8. <https://doi.org/10.1590/1809-2950/17014525032018>.
38. Jacobs JM, Hammerman-Rozenberg R, Cohen A, Stessman J. Chronic back pain among the elderly: prevalence, associations, and predictors. *Spine (Phila Pa 1976)*. 2006;31(7):E203-7. PMID: 16582841; <https://doi.org/10.1097/01.brs.0000206367.57918.3c>.
39. Tzeng PL, Lin CY, Lai TF, et al. Daily lifestyle behaviors and risks of sarcopenia among older adults. *Arch Public Health*. 2020;78(1):113. PMID: 33292561; <https://doi.org/10.1186/s13690-020-00498-9>.
40. Smith L, Tully M, Jacob L, et al. The Association Between Sedentary Behavior and Sarcopenia Among Adults Aged ≥65 Years in Low- and Middle-Income Countries. *Int J Environ Res Public Health*. 2020;17(5):1708. PMID: 32151034; <https://doi.org/10.3390/ijerph17051708>.
41. Global Recommendations on Physical Activity for Health. Geneva: World Health Organization; 2010. PMID: 26180873.
42. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451-62. PMID: 33239350; <https://doi.org/10.1136/bjsports-2020-102955>.
43. Kitamura A, Seino S, Abe T, et al. Sarcopenia: prevalence, associated factors, and the risk of mortality and disability in Japanese older adults. *J Cachexia Sarcopenia Muscle*. 2021;12(1):30-8. PMID: 33241660; <https://doi.org/10.1002/jcsm.12651>.
44. Saraiva MD, Apolinario D, Avelino-Silva TJ, et al. The Impact of Frailty on the Relationship between Life-Space Mobility and Quality of Life in Older Adults during the COVID-19 Pandemic. *J Nutr Health Aging*. 2021;25(4):440-7. PMID: 33786560; <https://doi.org/10.1007/s12603-020-1532-z>.
45. de Lima MDCC, Perracini MR, Guerra RO, et al. Accuracy of the life-space mobility measure for discriminating frailty. *Rev Bras Geriatr Gerontol*. 2022;25(5):e210219. <https://doi.org/10.1590/1981-22562022025.210219.pt>
46. Kuspinar A, Verschoor CP, Beauchamp MK, et al. Modifiable factors related to life-space mobility in community-dwelling older adults: results from the Canadian Longitudinal Study on Aging. *BMC Geriatr*. 2020;20(1):35. PMID: 32005107; <https://doi.org/10.1186/s12877-020-1431-5>.
47. Rudnicka E, Napierała P, Podfigurna A, et al. The World Health Organization (WHO) approach to healthy ageing. *Maturitas*. 2020;139:6-11. PMID: 32747042; <https://doi.org/10.1016/j.maturitas.2020.05.018>.

Authors' contributions: Batista PP: conceptualization (equal), data curation (equal), formal analysis (equal), investigation (equal), methodology (equal), and writing-original draft (equal); Perracini MR: conceptualization (equal), data curation (equal), formal analysis (equal), investigation (equal), methodology (equal), project administration (equal), and writing-review and editing (equal); de Amorim JSC: conceptualization (equal), formal analysis (equal), investigation (equal), methodology (equal), software (equal), and writing-review and editing (equal); de Lima MDCC: formal analysis (equal), investigation (equal), methodology (equal), visualization (equal), and writing-review and editing (equal); Lima CA: data curation (equal), formal analysis (equal), investigation (equal), methodology (equal), software (equal), and writing-review and editing (equal); Pereira DS: conceptualization (equal), formal analysis (equal), investigation (equal), methodology (equal), supervision (equal), and writing-review and editing (equal); Dantas RG: data curation (equal), formal analysis (equal), investigation (equal), methodology (equal), and visualization (equal); Fittipaldi EODS: formal analysis (equal), investigation (equal), methodology (equal), and visualization (equal); Santos AD: formal analysis (equal), investigation (equal), methodology (equal), and visualization (equal); Campos HLM: formal analysis (equal), investigation (equal), methodology (equal), and

visualization (equal); and Pereira LSM: conceptualization (equal), data curation (equal), formal analysis (equal), investigation (equal), methodology (equal), supervision (equal) and writing-review and editing (lead). All authors approved the final version of the study and reserved public responsibility for its content

Remobilize Research Network Affiliations: Alexandre da Silva, Francis Trombini-Souza, Adriana Guedes Carlos, Juliana Maria Gazzola, Mirian Moreira, Paulo Henrique Silva Pelicioni, Mônica Beatriz Ferreira, Etienne Duim, Nayara Tasse de Oliveira Cirino, Renata dos Ramos Varanda, Suzana Albuquerque de Moraes, Guilherme Medeiros de Alvarenga, Cristina Cristovão Ribeiro da Silva, Renato Barbosa dos Santos, Sarah Giulia Bandeira Felipe, and Lygia Paccini Lustosa (*in memoriam*)

Sources of funding: No funding sources

Conflicts of interest: The authors declare no conflicts of interest

Date of first submission: March 24, 2022

Last received: July 18, 2022

Accepted: August 19, 2022

Address for correspondence:

Patricia Parreira Batista
Universidade Federal de Minas Gerais (UFMG)
Av. Pres. Antônio Carlos, 6.627
Pampulha — Belo horizonte (MG) — Brasil
CEP 31270-901
Tel. (+55 31) 34097395
E-mail: ppb@ufmg.br

