

UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<https://www.thieme-connect.de/products/ejournals/abstract/10.1055/s-0043-1774580>

DOI: <https://doi.org/10.1055/s-0043-1774580>

Direitos autorais / Publisher's copyright statement:

©2023 by Associação Arquivos de Neuro-Psiquiatria. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

depend on genotype. In this case, the onset of symptoms was in the first months of life with axial and appendicular hypotonia, dysphagia, early pyramidal and cerebellar signs and her survival was after early childhood. This case is classified as mild PCH after phenotype-genotype correlation and according to the report of other authors. However, it is important to note that the progression of spastic paraplegia may not have a favorable outcome.

Final comments: The phenotype of hereditary early-onset spastic paraplegia associated with the EXOSC3 gene was described in this report. Genetic tests are important for performing differential diagnosis for suspected cerebral palsy when there are no risk factors, in addition to prognostic guidance and genetic counseling.

Code: PE123

A case report of neonatal PURA syndrome

Leticia Pugim Ferreira¹, Ana Chrystina Souza Crippa¹, Liara Bohnert¹, Maytza Mayndra Côrrea¹

¹Universidade Federal do Paraná, Hospital das Clínicas, Curitiba PR, Brazil

Case presentation: G.D.V.S, a male neonate, was admitted into the neonatal intensive care unit due to respiratory insufficiency. On his sixth day of life, the patient presented with a series of tonic movements and spasm in upper and inferior limbs, followed by an approximate five-minute duration, apnea and central cyanosis. He had a term and complication-free pregnancy. On admission, could be noted global hypotonia, difficulties for nourishing, hyperreflexia, facies with cleft palate and micrognathia. He later developed an excessive hyper startle responsiveness, oculogyric crises and persistent dyskinesia. Electroencephalography has no spikes. Cerebral magnetic resonance imaging visualizes a diffuse cerebral volumetric reduction and subdural hydroma. Genetic test shows deletion of 152Kb, on heterozygous, with a pathogenic variation involving the PURA gene. During hospitalization, movements had a positive response to the use of benzodiazepines (midazolam) and was discharged after treatment of several complications (infections, chyloperitoneum, panhypopituitarism), in addition to tracheostomy, gastrostomy and continuous use of oxygen.

Discussion: PURA syndrome is caused by the mutation of the purine rich binding element protein α (PURA) gene in chromosome 5q31.2–q31.3. Neonatal patients exhibited hypotonia, feeding difficulties, apnea or primary hypoventilation, intrauterine excessive hiccups and drowsiness. The pediatric patients demonstrated moderate to severe mental retardation, epilepsy, progressive hip dysplasia, scoliosis, dysphagia, salivation and constipation. Respiratory insufficiency, including central and obstructive sleep apnea and recurrent pulmonary aspiration, were frequently observed. Early-onset feeding difficulties with moderate dysphagia and evidence of tracheal aspiration often needed nasogastric or gastric-tube feeding. Moderate to severe neurodevelopmental delays might occur, with some developing later epilepsy and nonepileptic hyperkinetic movements (dystonia, dyskinesia, and eye movement abnormalities). Most patients showed a decreased volume of white matter, a slight enlargement of lateral ventricles, and subarachnoid cysts in cerebral magnetic resonance.

Final comments: In newborns with severe hypotonia associated with respiratory abnormalities or movement disorders, further evaluation is needed since early diagnosis and inter-

vention provides a better prognosis and allows genetic counseling.

Code: PE127

Infantile neuroaxonal dystrophy (INAD): a case report

Fernanda Ferrão Antonio¹, Maria Luiza Benevides¹, Paula Thais Bandeira Elias¹, Larisse Souza Somavilla¹, Ana Carolina Piauilino Falcão¹, Isabelle Salgado Castellano¹, Katia Maria Ribeiro Schmutzler¹, Ana Carolina Coan¹, Karine Couto Sarmiento Teixeira¹

¹Universidade Estadual de Campinas, Campinas SP, Brazil

Case presentation: A previously healthy 3-year-old girl was admitted with a history of loss of developmental milestones since 18 months of age. So far, only language delay had been noticed. It evolved from then on, with frequent falls, incoordination, and truncal hypotonia. Throughout the next year, she lost the ability to walk. During the same year, she began to have episodes of tonic seizures, with partial control after the introduction of levetiracetam. When examined, there was severe global hypotonia, with strabismus and nystagmus. During the investigation, it was identified diffuse cerebellar atrophy in the MRI. In addition, there was elevated aspartate aminotransferase (AST)/alanine aminotransferase (ALT) ratio and elevated lactate dehydrogenase (DHL). At the moment of the initial investigation, there was no optic atrophy. The molecular genetic testing showed biallelic pathogenic variants in PLA2G6 in homozygosis.

Discussion: Phospholipase A2 group VI (PLA2G6)-associated neurodegeneration (PLAN) is associated with two childhood neurologic disorders: infantile neuroaxonal dystrophy (INAD) and atypical neuroaxonal dystrophy (atypical NAD). The most common presentation during the first years of life is infantile neuroaxonal dystrophy (INAD) which usually begins between the ages of six months and three years with psychomotor regression or delay, hypotonia, and progressive spastic tetraparesis. Commonly, there is strabismus, nystagmus, and optic atrophy. Disease progression is rapid, leading to loss of the ability to walk, progressive cognitive decline, and visual impairment. Typically, there is an elevated AST/ALT ratio and increased levels of DHL. The neuroimages can show cerebellar atrophy and a hypointense globus pallidus in T2 MRI, indicating iron accumulation. Before the onset of genetic testing, the establishment of the diagnosis was based on the clinical features and tissue biopsy, with the evidence of dystrophic axons. Nowadays, the use of molecular testing with the identification of biallelic pathogenic variants in PLA2G6 confirms the diagnosis.

Final comments: This case describes INAD, one of the phenotypes of PLAN. It has been associated with psychomotor regression, early truncal hypotonia, and visual abnormalities. The knowledge about its evolution contributes to the development of therapeutic possibilities in the future and the adequate management and orientation of the child and its family.

Code: PE128

Infantile neuroaxonal dystrophy associated with seizures in a patient from a teaching hospital in southern Brazil: case report

Heloisa Augusta Castralli¹, Bruna Gulate da Conceição², Antônio Diniz da Rosa Pereira²

¹Universidade Federal de Santa Maria, Santa Maria RS, Brazil

²Hospital Universitário de Santa Maria, Santa Maria RS, Brazil